
1 Trig substitutions for integration

Trig substitutions are useful for reducing two terms into one, particularly when are solving integrals with two terms under a root, such as $\int \frac{\sqrt{25x^2-4}}{x} dx$. In cases like this, we can use a trig substitution to reduce the two terms and then easily eliminate the root.

There are three situations that we can come across, and for each we form a right-angle triangle, labelling each side and then choosing a trig ratio.


1. When $x^2 + a^2$ is embedded in the integral, label the triangle like so:



From the triangle, $\tan \theta = \frac{x}{a}$, meaning $x = a \tan \theta$.

Then,
$$\frac{dx}{d\theta} = a \sec^2 \theta$$


2. When $a^2 - x^2$ is embedded in the integral, label the triangle like so:



From the triangle, $\sin \theta = \frac{x}{a}$, meaning $x = a \sin \theta$

Then,
$$\frac{dx}{d\theta} = a\cos\theta$$

3. When $x^2 - a^2$ is embedded in the integral, label the triangle like so:




1

From the triangle, $\cos \theta = \frac{a}{x}$, meaning $x = \sec \theta$

Then,
$$\frac{dx}{d\theta} = a \sec \theta \tan \theta$$

This quite a tricky concept so here are a couple of examples to illustrate:



Example 1

$$\int \frac{\sqrt{25x^2-4}}{x} \, dx$$

This is in the form $x^2 - a^2$ so we set up our triangle as so:

$$\cos \theta = \frac{2}{5x}$$
$$x = \frac{2}{5} \sec \theta$$

$$x = \frac{2}{5}\sec^{3x}\theta$$

$$dx = \frac{2}{5} \sec \theta \tan \theta \, d\theta$$

Now we can substitute everything into our integral:

$$\int \frac{\sqrt{25(\frac{2}{5}\sec\theta)^2 - 4}}{\frac{2}{5}\sec\theta} \times \frac{2}{5}\sec\theta\tan\theta\,d\theta$$

Simplifying:
$$\int \frac{\sqrt{4 \sec^2 \theta - 4}}{\frac{2}{5}} \times \frac{2}{5} \tan \theta \, d\theta$$

$$\int \frac{\sqrt{4(\sec^2 \theta - 1)}}{\frac{2}{5}} \times \frac{2}{5} \tan \theta \, d\theta$$
$$\int \frac{\sqrt{4\tan^2 \theta}}{\frac{2}{5}} \times \frac{2}{5} \tan \theta \, d\theta$$

$$\int \frac{\sqrt{4\tan^2\theta}}{\frac{2}{5}} \times \frac{2}{5} \tan\theta \, d\theta$$

$$\int 2 \tan \theta \times \tan \theta \, d\theta = 2 \int \tan^2 \theta \, d\theta$$

We can't directly integrate this, but by using the $\tan^2 \theta = \sec^2 \theta - 1$ identity, we can rewrite the integral and do it easily:

$$2\int (\sec^2 \theta - 1) d\theta = 2\tan \theta - 2\theta + c$$

Finally, we go back to our original triangle and write our solution in terms of x again:

$$\tan \theta = \frac{\sqrt{25x^2 - 1}}{1}$$

$$\tan \theta = \frac{\sqrt{25x^2 - 4}}{2}$$


$$\theta = \cos^{-1}\left(\frac{2}{5x}\right)$$

$$\int \frac{\sqrt{25x^2 - 4}}{x} dx = \sqrt{25x^2 - 4} - 2\cos^{-1}\left(\frac{2}{5x}\right) + c$$

Example 2

$$\int \frac{1}{x^2 \sqrt{x^2 + 4}} \, dx$$

 $\int \frac{1}{x^2 \sqrt{x^2 + 4}} \, dx$ This is in the form $x^2 + a^2$ so we set up our triangle like so:



$$\tan \theta = \frac{x}{2}$$

$$x = 2 \tan \theta$$

$$dx = 2\sec^2\theta \, d\theta$$

Substituting into the integral:

$$\int \frac{1}{4\tan^2\theta \sqrt{4\tan^2\theta + 4}} 2\sec^2\theta \, d\theta$$
We can simplify the root:

$$\begin{array}{l} \sqrt{4\tan^2\theta+4} = \sqrt{4(\tan^2\theta+1)} = \sqrt{4\sec^2\theta}) = 2\sec\theta \\ \int \frac{1}{4\tan^2\theta\times2\sec\theta} 2\sec^2\theta\,d\theta \\ \int \frac{\sec\theta}{4\tan^2\theta}\,d\theta \\ \text{A bit of rearranging is now required to get this into a nice integral:} \\ \frac{1}{4}\int \frac{1}{\cos\theta} \times \frac{\cos^2\theta}{\sin^2\theta}\,d\theta \\ = \frac{1}{4}\int \frac{\cos\theta}{\sin^2\theta}\,d\theta \\ = \frac{1}{4}\int \csc\theta\cot\theta\,d\theta \\ = -\frac{1}{4}\csc\theta+c \\ \text{Finally, putting it back into terms of x:} \\ \text{Remembering that } \csc\theta=\frac{1}{\sin\theta} \\ \int \frac{1}{x^2\sqrt{x^2+4}}\,dx = -\frac{1}{4}\csc\theta=-\frac{1}{4}\times\frac{\sqrt{x^2+4}}{x}=-\frac{\sqrt{x^2+4}}{4x}+c \end{array}$$

Questions

(Answers - page ??)

- $1. \int \sqrt{1-x^2} \, dx$
- $2. \int \sqrt{4 9x^2} \, dx$
- $3. \int \sqrt{1-7x^2} \, dx$
- 4. $\int \frac{\sqrt{x^2+16}}{x^4} dx$
- 5. $\int \frac{2}{x^4 \sqrt{x^2 25}} dx$
- 6. $\int x^3 (3x^2 4)^{\frac{5}{2}} dx$
- 7. $\int x^3 \sqrt{4 9x^2} \, dx$
- $8. \int \frac{\sqrt{x^2+1}}{x} dx$
- $9. \int \frac{\sqrt{1-x^2}}{x} \, dx$
- 10. $\int \frac{(x^2-1)^{\frac{3}{2}}}{x} dx$
- $11. \int \frac{1}{\sqrt{e^{2x}-1}} dx$
- $12. \int \cos x \sqrt{9 + 25\sin^2 x} \, dx$
- 13. 2022 Scholarship exam Show that $\int \frac{1}{\sqrt{1+x^2}} dx = \ln |\sqrt{1+x^2} + x| + c$