

1 Sum of roots of polynomials

The sum of the roots of any polynomial in the form $ax^n + bx^{n-1} + cx^{n-2} + \dots + z = 0$ will always be equal to $-\frac{b}{a}$.

We can see that this holds for quadratics in the form $ax^2 + bx + c = 0$ as we know from when we factorise we need to find two numbers that multiply to c and add to b . This gives us the factors, and since the roots are $(x - x_1)$, it means the sum will be $-b$ (which is $\frac{-b}{1}$ since $a = 1$ here).

We can also see this from the quadratic equation: $x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$

If we add the two roots, we get:

$$\frac{-b + \sqrt{b^2 - 4ac}}{2a} + \frac{-b - \sqrt{b^2 - 4ac}}{2a} = -\frac{2b}{2a} = -\frac{b}{a}$$

This holds for all polynomials. For example, in the polynomial $p(x) = 2x^4 - x^3 + 2x - 1 = 0$ we know the four roots will sum to $\frac{1}{2}$, since $-(-\frac{1}{2}) = \frac{1}{2}$.

Questions

1. Find the roots of the equation $z^{11} = 1$. Use this to show that:

$$\cos\left(\frac{2\pi}{11}\right) + \cos\left(\frac{4\pi}{11}\right) + \cos\left(\frac{6\pi}{11}\right) + \cos\left(\frac{8\pi}{11}\right) + \cos\left(\frac{10\pi}{11}\right) = -\frac{1}{2}$$

2. If α is a complex root of the equation $z^5 = 1$, show that $\alpha + \alpha^2 + \alpha^3 + \alpha^4 = -1$

3. The roots of the quadratic equation $ax^2 + bx + c = 0$ are $\sin \theta$ and $\cos \theta$.

Show that: $\frac{\sin \theta}{1 - \cot \theta} + \frac{\cos \theta}{1 - \tan \theta} = -\frac{b}{a}$