

# 1 Parametric integration

Just as we can differentiate parametrically, we can also evaluate definite integrals parametrically.

Suppose we want to evaluate the integral  $\int_a^b y \, dx$  but we only know the parametric form

$$\begin{cases} x = x(t) \\ y = y(t) \end{cases} \quad \text{and eliminating the parameter is not feasible.}$$

We just evaluate the values of  $t$  when  $x$  has the values  $a$  and  $b$ , creating bounds for a new definite integral. We also change the function being integrated to  $y(t) \times x'(t)$ , then evaluate the definite integral.

E.g. Evaluate  $\int_{-1}^{\frac{1}{2}} y \, dx$  for the parametric curve given by  $\begin{cases} x = \sin(t) \\ y = 2(\sin(t) + \cos(t)) \end{cases}$

First, we find  $dx$ :

$$x = \sin(t)$$

$$dx = \cos(t) \, dt$$

So now our integral is  $\int \underbrace{2(\sin(t) + \cos(t))}_{y(t)} \underbrace{\cos(t) \, dt}_{dx}$

Next, we need to rewrite the upper and lower limits in terms of  $t$ :

Upper limit of integration:

$$\sin(t) = \frac{1}{2}$$

$$t = \frac{\pi}{6}$$

Lower limit:

$$\sin(t) = -1$$

$$t = -\frac{\pi}{2}$$

Our integral becomes  $\int_{-\frac{\pi}{2}}^{\frac{\pi}{6}} 2(\sin(t) + \cos(t)) \cos(t) \, dt = \int_{-\frac{\pi}{2}}^{\frac{\pi}{6}} 2 \sin(t) \cos(t) + \cos^2(t) \, dt$

$\int_{-\frac{\pi}{2}}^{\frac{\pi}{6}} \sin(2t) + \cos(2t) + 1 \, dt$  (By the double-angle formulas)

$$\left[ \frac{-\cos(2t)}{2} + \frac{\sin(2t)}{2} + t \right]_{-\frac{\pi}{2}}^{\frac{\pi}{6}}$$


$$\left( -\frac{1}{4} + \frac{\sqrt{3}}{2} + \frac{\pi}{6} \right) - \left( \frac{1}{2} + 0 - \frac{\pi}{2} \right) = \frac{\sqrt{3}}{4} - \frac{3}{4} + \frac{2\pi}{3}$$

## Questions

(Answers - page ??)


1. Evaluate  $\int_0^1 y \, dx$  for the parametric curve given by  $\begin{cases} x = 4 - t \\ y = t^2 - 3t \end{cases}$
2. Evaluate  $\int_{-\frac{1}{2}}^1 y \, dx$  for the parametric curve given by  $\begin{cases} x = \sin(t) \\ y = 2(\cos(t) - \sin(t)) \end{cases}$
3. Evaluate  $\int_0^{\sqrt{3}} y \, dx$  for the parametric curve given by  $\begin{cases} x = \tan(t) \\ y = \sin(t) \end{cases}$
4. Use parametric integration to show that the area of a circle of radius  $r$  is  $A = \pi r^2$ , remembering that the parametric form of a circle is  $\begin{cases} x = r \cos(t) \\ y = r \sin(t) \end{cases}$
5. Find the area enclosed between a parabola and its latus rectum, the line  $x = a$ , where  $a > 0$  and the parameterised equation for the parabola is  $\begin{cases} x = at^2 \\ y = 2at \end{cases}$
6. The graph shows the parametric function  $\begin{cases} x = \cos(2t) \\ y = 2(\cos(t) + \sin(t)) \end{cases} \quad -\frac{\pi}{4} \leq t \leq \frac{3\pi}{4}$

Find the area inside the curve.



7. The astroid shown below is defined by the parametric equations

$$\begin{cases} x = \cos^3(t) \\ y = \sin^3(t) \end{cases} \quad 0 \leq t \leq 2\pi$$



By evaluating  $\int_0^1 y \, dx$ , or otherwise, calculate the exact area of the astroid.