

1 Turning equations into quadratics

A normal quadratic appears in the form $ax^2 + bx + c = 0$, whereas a hidden quadratic can be written as $a[f(x)]^2 + b[f(x)] + c = 0$

When there are three terms in an equation, we can often turn them into a quadratic, where the subject is not x but another expression $f(x)$ that we substitute in.

For example, $e^{4x} - 5e^{2x} + 6 = 0$ can be solved by making it a quadratic in terms of e^{2x} .

$$u = e^{2x}$$

$$u^2 - 5u + 6 = 0$$

$$u = 2, 3$$

Then we just back-substitute and solve:

$$e^{2x} = 2$$

$$2x = \ln 2$$

$$x = \frac{\ln 2}{2}$$

$$e^{2x} = 3$$

$$2x = \ln 3$$

$$x = \frac{\ln 3}{2}$$

If all three terms contain a variable, we can also divide the equation through by something to turn one of those into a constant, enabling us to then solve it as a quadratic.

For example, $3(2^{3x}) - 11(2^{2x}) - 2^{x+2} = 0$

If we divide each term by a common factor of 2^x , the equation changes to:

$$\frac{3(2^{3x})}{2^x} - \frac{11(2^{2x})}{2^x} - \frac{2^{x+2}}{2^x} = 0$$

$$3(2^{2x}) - 11(2^x) - 2^2 = 0$$

We can now make the substitution $u = 2^x$ to solve the equation:

$$3u^2 - 11u - 4 = 0$$

$$u = -\frac{1}{3}, 4$$

Since 2^x can clearly never be negative, we can disregard the first solution.

$$2^x = 4$$

$$x = 2$$

Questions

(Answers - page ??)

1. Solve $2^x + 4^x = 24$
2. Solve $4^x + 6^x = 9^x$
3. Solve $8(9^x) + 3(6^x) - 81(4^x) = 0$
4. Solve $25^x + 2(15^x) - 24(9^x) = 0$
5. Solve $e^x - 20e^{-x} = 1$
6. Solve $3^{2x+1} - 82 \times 3^x + 27 = 0$
7. Solve $x^{\frac{1}{4}} + 2x^{-\frac{1}{4}} = 3$
8. Solve $x^2 + 5x - 4 + \frac{5}{x} + \frac{1}{x^2} = 0$
9. Solve $x - 10\sqrt{x+2} + 24 = 0$
10. Solve $x^4 - 4x^3 - 10x^2 - 4x + 1 = 0$