

AOS Senior AP Calculus BC, Spring 2024

Cumulative, Quarter 3

(Parametric, Polar, Logistic)

_____/ 30



ACADEMIES OF LOUDOUN

HONOR CODE



Honesty and integrity are the foundations of good academic work. Whether you are working on a problem set, lab report, project, presentation, or paper, do not engage in plagiarism, unauthorized collaboration, cheating, or facilitating academic dishonesty. Our expectation is for our students to be successful while being trustworthy. The honor code is not intended to be punitive, but rather a guide for all students and faculty to follow. For these reasons, the Academies of Loudoun will uphold the following Honor Code:

On my honor, I have not accepted or provided any unauthorized aid on this test, quiz, or assignment.

As an Academies of Loudoun student, you agreed to uphold the Academies Honor Code. Please write the Honor Code Pledge below and sign this document.

Student Signature

Class

Date

Print Name:

1. If a particle in the first quadrant is moving towards the x axis then

(a) $\frac{d^2y}{dt^2} > 0$

(b) $\frac{d^2x}{dt^2} > 0$

(c) $\frac{dy}{dt} < 0$

(d) $\frac{dx}{dt} < 0$

2. $x(t) = \cos t$ and $y(t) = \sin t$ are the parametric equations for

(a) A square

(b) A hyperbola

(c) A circle

(d) A parabola

3. To find the slope of the tangent line to a parametric curve at the point where $t = c$ you should

(a) Evaluate $y'(c)/x'(c)$

(b) Evaluate $x'(c)/y'(c)$

(c) Evaluate $x'(c)$

(d) Evaluate $y'(c)$

4. If a parametric curve has a point where $x'(a) = 0$ and $y'(a) = 0$ then

(a) There is no tangent line at $t = a$

(b) The curve must cross itself

(c) There is a vertical tangent line at $t = a$

(d) There is a horizontal tangent line at $t = a$

5. The maximum growth rate for a logistic population with carrying capacity L

(a) Always occurs at $t = 0$

(b) Depends on the initial conditions

(c) Can happen more than once during a given solution

(d) Occurs when the population is $L/2$

6. The area enclosed by a polar curve between $\theta = \alpha$ and $\theta = \beta$ is always

(a) $\int_{\alpha}^{\beta} r(\theta) d\theta$

(b) $\int_{\alpha}^{\beta} \frac{1}{2} r^2(\theta) d\theta$

(c) $\int_{\alpha}^{\beta} r^2(\theta) d\theta$

(d) Dependent on if the curve intersects itself in the interval $\alpha < \theta < \beta$

7. To determine concavity of a parametric curve at the point where $t = c$

(a) Evaluate $\frac{d}{dt} \left(\frac{dy}{dt} \right)$

(b) Evaluate $x''(c)/y''(c)$

(c) Evaluate $y''(c)/x''(c)$

(d) Evaluate $\frac{\frac{d}{dt} \left(\frac{dy}{dx} \right)}{\frac{dx}{dt}}$

8. The graph of $r = a + b \sin(\theta)$

(a) Has an inner loop whenever $a > b$

(b) Has an inner loop whenever $a < b$

(c) Never intersects the y -axis

(d) Never intersects the x -axis

9. A logistic population graph $y = f(t)$ with a max population of L

(a) Has an increasing growth rate when $t > 0$

(b) Can oscillate for certain initial conditions

(c) Has a decreasing growth rate when $t > 0$

(d) Has an asymptote at $y = L$

-
10. If a polar graph is defined by $r(\theta)$ and $\frac{dr}{d\theta} > 0$ at a point where $\theta = \alpha$ then
- (a) The tangent line to the graph $\theta = \alpha$ has a positive slope
 - (b) The tangent line to the graph at $\theta = \alpha$ has a negative slope
 - (c) The graph's radius is decreasing at $\theta = \alpha$
 - (d) The graph's radius is increasing at $\theta = \alpha$
11. If $x(t)$ and $y(t)$ are the parametric equations of a curve, the curve will have a horizontal tangent line at $t = c$ if
- (a) $x(c) = 0$ and $x'(c) = 0$
 - (b) $x'(c) = 0$ and $y'(c) \neq 0$
 - (c) $y'(c) = 0$ and $x'(c) \neq 0$
 - (d) $x(c) = 0$ and $y(c) = 0$
12. A particle moves in a plane from an initial position given by the vector $\vec{r}_0 = \langle x_0, y_0 \rangle$ at time $t = 0$. The particle's velocity at any time t is described by the vector function $\vec{v}(t) = \langle v_x(t), v_y(t) \rangle$. Assuming the velocity function is integrable, which of the following expressions correctly describes the particle's position $\vec{r}(t)$ at any later time t ?
- (a) $\vec{r}(t) = \vec{r}_0 + \frac{1}{2}\vec{v}(t)t^2$
 - (b) $\vec{r}(t) = \vec{r}_0 + \int_0^t \vec{v}(t) dt$
 - (c) $\vec{r}(t) = \vec{r}_0 + \vec{v}(t)t$
 - (d) $\vec{r}(t) = \vec{r}_0 + \int \vec{v}(t)$
13. The distance traveled from $t = a$ to $t = b$ of a particle with position vector $\langle x(t), y(t) \rangle$ is given by
- (a) $\int_a^b \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} dt$
 - (b) $\int_a^b |x'(t) + y'(t)| dt$
 - (c) $\int_a^b \sqrt{x^2(t) + y^2(t)} dt$
 - (d) $\sqrt{(x(t) - x(0))^2 + (y(t) - y(0))^2}$

14. If n is a positive integer, the graph of $r = \sin(n\theta)$ always

- (a) Completes exactly one period of the graph over $0 \leq \theta < 2\pi$
- (b) Has one intercept at $(0, 0)$
- (c) Is a rose with n petals
- (d) Is a rose with $2n$ petals

15. Which of the following is **not** a polar-rectangular transformation equation?

- (a) $\tan \theta = \frac{x}{y}$
- (b) $y = r \sin \theta$
- (c) $x = r \cos \theta$
- (d) $x^2 + y^2 = r^2$

KEY

1. C
2. C
3. A
4. A
5. D
6. D
7. D
8. B
9. D
10. D
11. C
12. B
13. A
14. B
15. A