AOS Senior AP Calculus BC, Spring 2024 AP Test Review, Exam 1



ACADEMIES OF LOUDOUN HONOR CODE



Date

Honesty and integrity are the foundations of good academic work. Whether you are working on a problem set, lab report, project, presentation, or paper, do not engage in plagiarism, unauthorized collaboration, cheating, or facilitating academic dishonesty. Our expectation is for our students to be successful while being trustworthy. The honor code is not intended to be punitive, but rather a guide for all students and faculty to follow. For these reasons, the Academies of Loudoun will uphold the following Honor Code:

On my honor, I have not accepted or provided any unauthorized aid on this test, quiz, or assignment.
As an Academies of Loudoun student, you agreed to uphold the Academies Honor Code. Please write the Honor Code Pledge below and sign this document.

Class

Print Name:

Student Signature

1. Evaluate

$$\lim_{h \to 0} \frac{\sec(3(x+h)) - \sec(3x)}{h}$$

- (a) $3 \sec^2(3x)$
- (b) $3\sec(x)\tan(x)$
- (c) $3 \tan^2(3x)$
- (d) $9 \tan^2(3x)$
- (e) $3\sec(3x)\tan(3x)$
- 2. Let f be the function given by

$$f(x) = \begin{cases} 2x + 3b & \text{if } x \le 2\\ 3ax^2 & \text{if } x > 2 \end{cases}$$

What are all values of a and b for which f is differentiable at x = 2?

- (a) $a = \frac{-1}{6}$ $b = \frac{-2}{3}$
- (b) $a = \frac{1}{3}$ $b = \frac{-2}{3}$
- (c) $a = \frac{-1}{6}$ $b = \frac{1}{3}$
- (d) $a = \frac{1}{6}$ $b = \frac{-1}{3}$
- (e) $a = \frac{1}{6}$ $b = \frac{-2}{3}$
- 3. If the function f is continuous for all real numbers and if $f(x) = \frac{x^2 25}{x 5}$ when $x \neq 5$, then f(5) =
 - (a) 5
 - (b) 25
 - (c) -5
 - (d) -10
 - (e) 10
- 4. Evaluate

$$\lim_{x \to e} \frac{\ln x - 1}{x - e}$$

- (a) 1
- (b) e
- (c) 2e
- (d) $\frac{2}{e}$
- (e) $\frac{1}{e}$

$$\lim_{x \to 0} \frac{1 - \cos x}{x^2 + \sin(4x)} =$$

- (a) π
- (b) 1
- (c) -1
- (d) $\frac{\pi}{2}$
- (e) 0
- 6. If f is the function defined

$$f(x) = \begin{cases} x^2 & \text{for } x < 3\\ \frac{1}{3} & \text{for } x \ge 3 \end{cases}$$

then
$$\int_{-2}^{4} f(x)dx$$
 is

- (a) $\frac{20}{3}$
- (b) $\frac{28}{3}$
- (c) 13
- (d) $\frac{-28}{3}$
- (e) 12
- 7. Evaluate

$$\int_0^3 \frac{x^2 + 5x + 6}{x + 2} dx =$$

- (a) $4 + 2 \ln 2$
- (b) $\frac{15}{2}$
- (c) $\frac{15}{2} + 2 \ln 2$
- (d) $\frac{17}{2} 2 \ln 2$
- (e) $\frac{27}{2}$

$$\int \frac{\cos{(\sqrt{x}+1)}}{\sqrt{x}} \ dx =$$

- (a) $\frac{1}{2}\sin(\sqrt{x}+1) + C$
- (b) $\ln |\cos (\sqrt{x} + 1)| + C$
- (c) $\ln |\sin (\sqrt{x} + 1)| + C$
- (d) $\cos(\sqrt{x} + 1) + C$
- (e) $2\sin(\sqrt{x}+1) + C$
- 9. Evaluate

$$\int \frac{2x}{x^2 + 9} dx =$$

- (a) $\frac{1}{2}\ln(x^2+9) + C$
- (b) $\frac{1}{x^2+9}+C$
- (c) $\frac{2}{(x^2+9)}+C$
- (d) $\frac{x}{3} \arctan\left(\frac{x}{3}\right) + C$
- (e) $\ln(x^2+9)+C$
- 10. The function g is continuous on the closed interval [2,10]. If $\int_9^1 g(x)dx = 25$ and $\int_1^5 \frac{g(x)}{2} dx = -12$, then

$$\int_{5}^{9} g(x) \ dx =$$

- (a) 49
- (b) 1
- (c) 13
- (d) -13
- (e) -1

11. Using the substitution $u = 2x^2 + 1$, the integral $\int_2^4 2x (2x^2 + 1)^3 dx$ is equal to which of the following?

(a)
$$2\int_{9}^{33} u^3 du$$

(b)
$$\frac{1}{2} \int_9^{33} u^3 du$$

(c)
$$2\int_{9}^{33} u^3 du$$

(d)
$$\frac{1}{4} \int_{9}^{33} u^3 du$$

(e)
$$\frac{1}{2} \int_{9}^{33} u^3 du$$

12. Evaluate

$$\int \frac{9x+1}{(2x+1)(x-3)} \, dx$$

(a)
$$\ln |2x+1| + \ln |x-3| + C$$

(b)
$$\frac{1}{2} \ln |2x+1| - 2 \ln |x-3| + C$$

(c)
$$4 \ln |2x + 1| - 2 \ln |x - 3| + C$$

(d)
$$2 \ln |2x+1| - 4 \ln |x-3| + C$$

(e)
$$\frac{1}{2} \ln |2x+1| + 4 \ln |x-3| + C$$

$$\int \frac{1}{x^2 - 16x + 80} \, dx =$$

(a)
$$\frac{1}{6}\arctan\left(\frac{x-8}{6}\right) + C$$


(b)
$$\frac{x}{6} \arctan \left| \frac{x-8}{6} \right| + C$$

(c)
$$\frac{1}{\ln|x^2 - 16x + 80|} + c$$

(d)
$$2 \ln |x - 20| - 4 \ln |+4| + C$$

(e)
$$\frac{1}{4}\arctan\left(\frac{x-8}{4}\right)+C$$

14. The graph of a piecewise linear function f is given.



What is the value of $\int_{1}^{7} (4f(x) - 1) dx$?

- (a) 9.5
- (b) 7.5
- (c) 9
- (d) 10
- (e) 8

$$\int_{1}^{\infty} xe^{-(x^2-1)} dx$$

- (a) $\frac{1}{e}$
- (b) $\frac{1}{2e}$
- (c) 2
- (d) divergent
- (e) $\frac{1}{2}$

16. Integrate

$$\int x^3 e^{2x} \ dx$$

(a)
$$\frac{1}{4}e^{2x} (4x^3 - 6x^2 + 6x - 3) + C$$

(b)
$$\frac{1}{8}e^{2x}(4x^3+6x^2-x+3)+C$$

(c)
$$\frac{1}{4}e^{2x}\left(4x^3+6x^2+6x+3\right)+C$$

(d)
$$\frac{1}{8}e^{2x}(2x^3 - 3x^2 + 6x - 3) + C$$

(e)
$$\frac{1}{8}e^{2x}\left(4x^3 - 6x^2 + 6x - 3\right) + C$$