
KEY KEY D - Matching - 1 point each question



1.
$$r = 3\csc(\theta)$$
 L

2.
$$r = 4\sin(3\theta)$$
 F

3.
$$r = -2\cos(\theta)$$
 B

4.
$$r = 4 + 2\cos(\theta)$$
 E

5.
$$r = 4\sin^2(\theta)$$
 I

6.
$$r = 3 - 2\sin(\theta)$$
 D

7.
$$r = 4\cos(3\theta)$$
 G

8.
$$r = 3\sec(\theta)$$
 J

9.
$$r = \theta/2$$
 K

10.
$$r = 4\sin(\theta) + \cos(\theta)$$
 H

11.
$$r = 3 + 2\sin(\theta)$$
 C

12.
$$r = 3\sin(\theta)$$
 A

Multiple Choice - 2 pts each

Work must be shown for credit.

- 1. Convert the polar coordinate to rectangular coordinates: $(-2, 2\pi/3)$
 - A. $(1, 1/\sqrt{3})$
 - **B.** $(1, -\sqrt{3})$
 - C. $(1, \sqrt{3})$
 - D. $(-1, 1/\sqrt{3})$
 - E. $(\sqrt{3}, 1)$
- 2. Convert the polar coordinate to rectangular coordinates: $(4, -\pi/2)$
 - A. (4,0)
 - B. (0,4)
 - C. (0, -4)
 - D. (-4,0)
 - E. (4, -4)
- 3. Convert the rectangular coordinate to polar coordinates: $(15, 5\sqrt{3})$
 - A. $(10\sqrt{3}, \frac{\pi}{4})$
 - **B.** $(10\sqrt{3}, \frac{\pi}{6})$
 - C. $(10\sqrt{3}, \frac{\pi}{3})$
 - D. $(\sqrt{30}, \frac{\pi}{6})$
 - E. $(10\sqrt{3}, \frac{\pi}{3})$
- 4. Convert the rectangular coordinate to polar coordinates: (-12, -12)
 - **A.** $(12\sqrt{2}, -\frac{3\pi}{4})$
 - B. $(2\sqrt{12}, \frac{3\pi}{4})$
 - C. $(12\sqrt{2}, \frac{3\pi}{4})$
 - D. $(2\sqrt{12}, -\frac{5\pi}{4})$
 - E. $(12\sqrt{2}, -\frac{\pi}{4})$

4

5. Convert the rectangular equation to polar: $x^2+y^2=16$ A. $r=\frac{16}{2\sin\theta}$

A.
$$r = \frac{16}{2\sin\theta}$$

B.
$$r = 4\sin\theta$$

C.
$$r = 16\sin\theta$$

D.
$$r = 4$$

E.
$$r = 16$$

6. Convert the rectangular equation to polar: 2xy=1 A. $r=\frac{1}{2(\sin\theta+\cos\theta)}$

A.
$$r = \frac{1}{2(\sin\theta + \cos\theta)}$$

B.
$$r^2 = 1 - 2\sin(\theta)\cos(\theta)$$

$$\mathbf{C.} \ r^2 = \frac{1}{2\sin\theta\cos\theta}$$

D.
$$r^2 = \frac{1}{2}\sec(\theta)\cot(\theta)$$

E.
$$r = \frac{1}{2\sin(2\theta)}$$

7. Convert the polar equation to rectangular: $\theta = 2\pi/3$

A.
$$y = -\sqrt{3}x^{2}$$

B.
$$x + y = \sqrt{3}$$

C.
$$y = \sqrt{3}x$$

D.
$$y = -(1/\sqrt{3})x$$

- E. None of the above
- 8. Convert the polar equation to rectangular: $r = \frac{2}{1 + \sin \theta}$

A.
$$x^2 + xy - y^2 = 4$$

B.
$$x^2 + y^2 = (y - 2)^2$$

C.
$$x^2 - y^2 = (x + y)$$

D.
$$x^2 + y^2 + y = 4$$

E.
$$x^2 - y^2 = 2(x+y)$$

| 5. | | |
|----|--|--|
| • | | |

- 9. Find the intersection points of $r = 3\cos\theta$ and $r = \sqrt{3}\sin\theta$
 - A. $\{2\pi/3, 4\pi/3\}$
 - B. $\{4\pi/3, 7\pi/3\}$
 - C. $\{\pi/3, 5\pi/6\}$
 - D. $\{5\pi/6, 11\pi/6\}$
 - **E.** $\{\pi/3, 4\pi/3\}$

9. _____

Free Response Section

Calculator Active

A remote control drone is flying in the air and casts a shadow onto an xy plane below. Its shadow on the xy-plane has position (x(t), y(t)) at time $t \ge 0$ with

$$\frac{dx}{dt} = \sqrt{3t}$$
 and $\frac{dy}{dt} = 3\cos\left(\frac{t^2}{2}\right)$

The particle is at position (1,4) at time t=4.

1. Find the acceleration vector at time t = 6.

$$a(4) = \langle v'_x(6), v'_y(6) \rangle = \langle 0.353553, 13.5178 \rangle$$

2. Find the y-coordinate of the position of the shadow at time t=1.

$$s(1) = s(4) + \int_{4}^{1} v(t) dt = \langle -7.0829, 3.52647 \rangle$$

3. On the interval $0 \le t \le 4$, at what time does the speed of the shadow first reach 3.2 ?

Solve
$$\sqrt{v_x^2(t) + v_y^2(t)} - 3.2 = 0$$
 and find $t = 0.441822$

4. Find the total distance traveled by the shadow over the time interval $1 \le t \le 4$.

$$\int_{1}^{4} \sqrt{v_x^2(t) + v_y^2(t)} dt = 10.0099$$