AOS Math 10, Spring 2024 Cumulative, Quarter 3 (Parametric, Polar, Vectors, Complex)

/ 40



ACADEMIES OF LOUDOUN HONOR CODE



Honesty and integrity are the foundations of good academic work. Whether you are working on a problem set, lab report, project, presentation, or paper, do not engage in plagiarism, unauthorized collaboration, cheating, or facilitating academic dishonesty. Our expectation is for our students to be successful while being trustworthy. The honor code is not intended to be punitive, but rather a guide for all students and faculty to follow. For these reasons, the Academies of Loudoun will uphold the following Honor Code:

| Student Signature | Class | Date | |
|---------------------------------|----------------------------|--------------------------------|-----------------|
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| the Honor Code Pledge below and | d sign this document. | | |
| As an Academies of Loudoun stud | | the Academies Honor Code | e. Please write |
| On my honor, I have not accepte | ed or provided any unautho | orized aid on this test, quiz, | or assignment. |

Print Name:

- 1. Let $f(x) = \sqrt{x}$. What is the equation of the tangent line to f at the point (4,2)?
 - (a) $y = -\frac{1}{2}x + 3$
 - (b) $y = \frac{1}{2}x$
 - (c) y = 2x 6
 - (d) $y = \frac{1}{4}x + 1$
- 2. What is the derivative of $s(t) = \sec \sqrt{t}$?
 - (a) $\sec \frac{1}{2\sqrt{t}} \tan \frac{1}{2\sqrt{t}}$
 - (b) $\frac{\sec\sqrt{t}\tan\sqrt{t}}{2\sqrt{t}}$
 - (c) $\sec \sqrt{t} \tan \sqrt{t}$
 - (d) $\tan^2 \sqrt{t}$
- 3. If f,g, and h are nonzero differentiable functions, then the derivative of $\frac{fg}{h}$ is
 - (a) $\frac{fg'h'-fgh'}{h^2}$
 - (b) $\frac{fgh'-fg'h-f'gh}{h^2}$
 - (c) $\frac{f'gh+fg'h+fgh'}{h^2}$
 - (d) $\frac{fg'h+f'gh-fgh'}{h^2}$
 - (e) $\frac{fg'+f'g}{h'}$

| 4. | The line tangent to the curve $y=\sqrt{16-x}$ at the point $(0,4)$ has slope (a) 4 |
|----|---|
| | (b) 1/8 |
| | (c) $-1/8$ |
| | (d) -8 |
| | (e) 8 |
| 5. | At what point(s) on the curve $x^2 - y^2 + x = 2$ is the tangent line vertical? (a) $(-2,0)$ only |
| | (b) $(1,\sqrt{2})$ only |
| | (c) $(1,0)$ and $(-2,0)$ |
| | (d) The tangent line is never vertical |
| | (e) $(1,0)$ only |
| 6. | If $y = 6\cos(3x)$ then what is y' ? |
| | (a) $18\sin(x)$ |
| | (b) $18\sin(3x)$ |
| | (c) $-18\sin(3x)$ |
| | (d) $-6\sin(3x)$ |
| | |
| | |

7. What is the value of

$$\lim_{\Delta x \to 0} \frac{2(x+\Delta x)^2 - 2x^2}{\Delta x}$$

- (a) 4x
- (b) 4
- (c) 2
- (d) Does not exist
- (e) 2x
- 8. If $w(t) = \sqrt{t^2 1}$ what is the value of w'(4)?
 - (a) $\frac{2}{\sqrt{15}}$
 - (b) $\frac{1}{\sqrt{15}}$
 - (c) $\frac{1}{2\sqrt{15}}$
 - (d) $\frac{4}{\sqrt{15}}$
- 9. At which x value does the graph of $y = 3x^2 10x + 15$ have a horizontal tangent line?
 - (a) $\frac{-3}{5}$
 - (b) $\frac{5}{3}$
 - (c) $\frac{-5}{3}$
 - (d) $\frac{3}{5}$

- 10. Find $\frac{dy}{dx}$ if $x^2 + 4xy + 2y^2 = 16$
 - (a) $\frac{-2(x+y)}{x+2y}$
 - $(b) \frac{-2(x+y)}{2x+y}$
 - (c) $\frac{-x+2y}{x+y}$
 - $(d) \ \frac{-x-2y}{2x+2y}$
- 11. At which x value(s) does the graph of $y = 2x^3 24x + 16$ have a horizontal tangent line?
 - (a) 2
 - (b) 1
 - (c) 2 and -2
 - (d) 1 and -1
- 12. If $h(x) = f(x^2 + 1)$ then which of the following is true?
 - (a) h'(x) = f'(2x)
 - (b) h'(x) = 2xf'(2x)
 - (c) $h'(x) = 2xf'(x^2 + 1)$
 - (d) $h'(x) = f'(x^2 + 1)$
- 13. If $f(x) = 10x^2 5$, what is the average rate of change of f(x) over the interval

$$-1 \le x \le 2$$

- (a) 30
- (b) 20
- (c) $\frac{20}{3}$
- (d) 10

- 14. If h(x) = f(x)g(x) and f(5) = 3, f'(5) = -1, $g(5) = -\frac{1}{2}$, g'(5) = 2, then what is the value of h'(5)?
 - (a) $\frac{13}{2}$
 - (b) $-\frac{9}{2}$
 - (c) $\frac{9}{2}$
 - (d) -2
- 15. If $f(x) = \sin(x^2 + 1)$ then find f''(x)
 - (a) $-4x^2 \sin(x^2+1) + 2\cos(x^2+1)$
 - (b) $4x^2 \sin(x^2+1) 2\cos(x^2+1)$
 - (c) $-4x^2 \sin(x^2+1) 2\cos(x^2+1)$
 - (d) $4x^2 \sin(x^2+1) + 2\cos(x^2+1)$
- 16. The height (in feet) of a ball thrown vertically upward is given by

$$s(t) = -16t^2 + 32t + 64$$

where t is in seconds. What is the velocity of the ball at time t=3 seconds?

- (a) 64ft/s
- (b) -16ft/s
- (c) 16ft/s
- (d) -64 ft/s

| 17 | . At which point on the graph | is the slope of the tangent | t line closest to the | e average rate of o | change of $f(x)$ | between |
|----|-------------------------------|-----------------------------|-----------------------|---------------------|------------------|---------|
| | points X and Y ? | | | | | |

- (a) B
- (b) C
- (c) D
- (d) E
- (e) A

18. Let $f(x) = x^3 - 6x^2 + 10$. At which point(s) on the graph of f is the tangent line parallel to the line 15x - y = 11?

- (a) (2,-6) and (-2,-22)
- (b) (5, -15) and (-1, 3)
- (c) (5, -15) and (2, -6)
- (d) (2, -6) and (-2, 22)