


## Ejercicios raíces

5.7 Determine the roots of  $f(x) = -12 - 21x + 18x^2 - 2.75x^3$  graphically. In addition, determine the first root of the function with bisection (linear interpolation and Newton-Raphson)

```
f = @(x) -12 - 21*x + 18*x.^2 - 2.75*x.^3;  
fplot(f);  
grid on;
```



This plot indicates that roots are located at about  $-0.4$ ,  $2.25$  and  $4.7$ . La primera raíz está en  $[-1,0]$ .

```
[x, i] = biseccion(f, -1, 0)
```

```
x = -0.4147  
i = 54
```

```
[x, i] = interpolacionLineal(f, -1, 0)
```

```
x = -0.4147  
i = 55
```

```
[x, i] = newtonRaphson(f, -1, 0)
```

```
x = -0.4147  
i = 4
```

```
[x, i] = biseccion(f, 2, 2.5)
```

```
x = 2.2198  
i = 50
```

```
[x, i] = interpolacionLineal(f, 2, 2.5)
```

```
x = 2.2198  
i = 55
```

```
[x, i] = newtonRaphson(f, 2, 2.5)
```

```
x = 2.2198  
i = 3
```

```
[x, i] = biseccion(f, 4.5, 5)
```

```
x = 4.7403  
i = 47
```

```
[x, i] = interpolacionLineal(f, 4.5, 5)
```


```
x = 4.7403  
i = 16
```

```
[x, i] = newtonRaphson(f, 4.5, 5)
```

```
x = 4.7403  
i = 3
```

5.8 Locate the first nontrivial root of  $\sin(x) = x^2$  where  $x$  is in radians. Use a graphical technique and a numerical algorithm..

```
f = @(x) sin(x) - x.^2;  
  
fplot(f);  
grid on;
```




La raíz trivial es 0; la no trivial está en  $[0.5, 1]$ .

```
[x, i] = biseccion(f, 0.5, 1)
```

```
x = 0.8767  
i = 51
```

5.9 Determine the positive real root of  $\ln(x^2) = 0.7$  (a) graphically, (b) using a numerical method, with initial guess in  $x_l = 0.5$  and  $x_u = 2$

```
f = @(x) log(x.^2) - 0.7;  
  
fplot(f);  
grid on;
```



La solución se encuentra en  $[1, 1.5]$ .

```
[x, i] = biseccion(f, 1, 1.5)
```

```
x = 1.4191
i = 51
```

Ahora, con  $x_l = 0.5$  and  $x_u = 2$ :

```
[x, i] = biseccion(f, 0.5, 2)
```

```
x = 1.4191
i = 53
```

Sí encuentra la raíz positiva pero en más iteraciones.

Escribe aquí las funciones bisección, interpolación lineal y Newton-Raphson.

```
function [xr, i] = biseccion(f, xl, xu)
    if sign(f(xl)) * sign(f(xu)) >= 0
        error('f(a) * f(b) < 0 no se satisface.')
    end

    MAX_ITER = 55;
    TOLER = eps;
    xr = (xl + xu) / 2;
    fx = f(xr);
    i = 0;
```

```

while fx ~= 0 && abs((xu - xl) / xu) > TOLER && i < MAX_ITER
    if sign(f(xl)) == sign(fx)
        xl = xr;
    else
        xu = xr;
    end
    xr = (xl + xu) / 2;
    fx = f(xr);
    i = i + 1;
end
end

```

```

function [x, i] = interpolacionLineal(f, a, b)
    if sign(f(a)) * sign(f(b)) >= 0
        error('f(a) * f(b) < 0 no se satisface.')
    end

```

```

    MAX_ITER = 55;
    TOLER = eps;
    fa = f(a);
    x = b - f(b) * (b - a) / (f(b) - fa);
    fx = f(x);
    i = 0;

```

```

while abs((b - a) / x) > TOLER && fx ~= 0 && i < MAX_ITER
    if sign(fa) == sign(fx)
        a = x;
        fa = f(a);
    else
        b = x;
    end
    x = b - f(b) * (b - a) / (f(b) - fa);
    fx = f(x);
    i = i + 1;
end
end

```

```

function [x, i] = newtonRaphson(f, a, b)
    REL_TOL = sqrt(eps);
    MAX_ITER = 53;
    x = (a + b) / 2;
    df = matlabFunction(diff(sym(f)));

    if nargin(df) == 0
        df = @(x) df();
    end

    i = 0;
    flag = true;
    while flag
        xp = x;

```

```
x = xp - f(xp) / df(xp);  
i = i + 1;  
flag = i < MAX_ITER && abs((x - xp) / x) > REL_TOL;  
end  
end
```