



Chem1 Problem Set #1 (Lectures 1 - 12) - Solutions

Question 1:



(i)



(ii)




(iii)



enantiomers

(iv)




enantiomers

enantiomers


(v)




(vi)




(vii)




(viii)



(ix)




(x)




Question 2:


(i)




(ii)

**Question 3:**


(i)




(ii)




(iii)




(iv)




(v)

**Question 4:**


(i)




(ii)




(iii)




(iv)











(v)



(vi)


**Question 5:**

(i) a (ii) a (iii) c (iv) b (v) b (vi) c (vii) a


Question 6:

Question 7:


(i)




(ii)

both I and F *eq.*

(iii)



(iv)




*t*Bu group *eq.* in both, it acts as a *steric lock* since the bulk of the *t*Bu group can only be accommodated in the *eq.* orientation. The CH₃ group is *ax.* in (iii), *eq.* in (iv)


(ax. = axial eq. = equatorial)

Question 8:


(i)

all sp^3


(ii)




(iii)




(iv)




(v)




(vi)




(vii)




(viii)




(ix)




(x)




(xi)



(xii)



(xiii)



Question 9:

- (i) (*R*)-3-chloro-3-methylhexane
- (ii) 1-bromo-2-methyl-2-propanol
- (iii) (*E*)-1-chloropropene
- (iv) (*Z*)-3-bromo-2-hexene
- (v) 1-bromo-1-methylcyclopentane
- (vi) 3-chlorocyclohexene
- (vii) ethyl butanoate
- (viii) *m*-chlorobenzoic acid
- (ix) *p*-bromonitrobenzene
- (x) 5-bromo-2-nitrophenol