


# The impacts of ocean acidification on the distribution, abundance, species composition and shell size of thecosome pteropods



**Camille Pagniello**

Dalhousie University

*B.Sc. Co-op, Combined Honours in Marine Biology and Physics,  
with minors in Mathematics and Ocean Science (Expected May 2015)*

October 18, 2013

Image courtesy of Amy Maas

# Ocean Acidification




Figure: Rost et al. 2008

# Aragonite Saturation State

$$\Omega_{\text{aragonite}} = [\text{Ca}^{2+}] [\text{CO}_3^{2-}] / K_{\text{sp aragonite}}$$




Figure: Feely et al. 2012

# Impacts on the Plankton

- Much previous work has focused on coastal species
- Effects of experimentally decreased pH has been variable, depending on species
- Less work done on zooplankton and the open ocean

PHYSIOLOGICAL RESPONSE	MAJOR GROUP	# SPECIES STUDIED	RESPONSE TO INCREASING CO <sub>2</sub>				
			a	b	c	d	
CALCIFICATION							
	Coccolithophores	4	2	1	1	1	
	Planktonic Foraminifera	2	2	-	-	-	
	Molluscs	6	5	-	1	-	
	Echinoderms	3	2	1	-	-	
	Tropical Corals	11	11	-	-	-	
	Coralline Red Algae	1	1	1	-	-	
PHOTOSYNTHESIS <sup>1</sup>							
	Coccolithophores <sup>2</sup>	2	-	2	2	-	
	Prokaryotes	2	-	1	1	-	
	Seagrasses	5	-	5	-	-	


# Pteropods – Charismatic Microfauna

- Shell-forming planktonic gastropods aka ‘sea butterflies’
- Thecosome = Shelled
- Little known about their abundance, distribution, movements, and ecological role
- Important as prey item for many commercial fish (e.g., salmon)




Image courtesy of Amy Maas

# Ocean Acidification and Thecosome Pteropods



# Geographic Distribution of Pteropods



- highest species diversity in tropical and subtropical regions (Lalli and Gilmer, 1989; Fabry et al., 2008)
- highest abundances found at high latitudes (Lalli and Gilmer, 1989)

## Research Question

---

How will continued ocean acidification impact the ecology and role of *oceanic thecosome pteropods* within the pelagic ecosystem?

# Objective

Quantify:

- **geographic distribution**
- **numerical abundance**
- **species composition**
- **shell size**
- **biomass**

of *oceanic thecosome pteropods* in the southeast Pacific.

# Abiotic Factors

Correlate these quantities with:

- **latitude**
- **temperature**
- **surface water pH**
- **dissolved inorganic carbon (DIC)**
- **alkalinity**
- **aragonite saturation state**

## Hypotheses

---

**H1.** Species composition will vary with:

- latitude
- sea surface temperature
- pH
- alkalinity
- DIC
- aragonite saturation state

Species-specific relationships are expected.

## Hypotheses

---

H2. Abundance  $\downarrow$  with:

- $\downarrow$  latitudes
- $\uparrow$  sea surface temperatures
- $\downarrow$  pH
- $\downarrow$  alkalinity
- $\uparrow$  DIC
- $\downarrow$  aragonite saturation state

## Hypotheses

---


**H3.** Shell size  $\uparrow$  with:

- $\downarrow$  latitudes
- $\uparrow$  sea surface temperatures
- $\uparrow$  pH
- $\uparrow$  alkalinity
- $\downarrow$  DIC
- $\uparrow$  aragonite saturation state

**H4.** Biomass will vary with all abiotic factors similarly to shell size.


# Sampling Methods

- Conductivity-Temperature-Depth (CTD) to characterize temperature, salinity
- Niskin bottle sampling to measure pH, DIC, alkalinity
- 333  $\mu\text{m}$ -mesh Neuston net to collect pteropods in surface waters
- possible use of mesh meter nets to sample deeper (to 75 m)



# Shell Size and Biomass

SPECIES	Group	Equation source	Conversion	Equation name	Equation (size-weight relationship)	Equation (Davis and Wiebe, 1985)
<i>Limacina</i> spp.	Round/cylindrical/globular	GLOBEC	Diameter→DW		$WW = 0.000194 \times L^{2.5473}$	WW→DW WW×0.28
<i>Clione</i> spp.	Barel/oval-shaped (naked)	GLOBEC	Length→WW	Pteropod (naked: Clione)	$WW = 10^{(2.533 \times \log(L) - 3.89095) \times 10^3}$	WW→DW WW×0.28
<i>Styliola</i> spp.	Cone/needle/tube/bottle-shaped	GLOBEC	Length→WW	Pteropod (cone-shaped: Styliola)	$WW = \pi \times L^{3 \times 3/25}$	WW→DW WW×0.28
<i>Clio</i> spp.	Triangular/pyramidal	GLOBEC	Length→WW	Pteropod (Clio)	$WW = 0.2152 \times L^{2.293}$	WW→DW WW×0.28
<i>Diacria</i> spp.	Triangular/pyramidal	GLOBEC	Length→WW	Pteropod (Clio)	$WW = 0.2152 \times L^{2.293}$	WW→DW WW×0.28
Thecosomata	Shelled	Davis and Wiebe (1985)	Length→WW		$WW = 0.2152 \times L^{2.293}$	WW→DW WW×0.28
Gymnosomata	Naked	Davis and Wiebe (1985)	Length→WW		$WW = 10^{(2.533 \times \log(L) - 3.89095) \times 10^3}$	WW→DW WW×0.28
Pteropoda	Shelled	Davis and Wiebe (1985)	Length→WW		$WW = 0.2152 \times L^{2.293}$	WW→DW WW×0.28



Figures: Bednaršek et al. 2012; Little and Copley 2003

# References

---

- Bednaršek, N., J. Možina, M. Vogt, C. O'Brien, and G.A. Tarling. 2012. The global distribution of pteropods and their contribution to carbonate and carbon biomass in the modern ocean. *Earth System Science Data*. 4: 167-186.
- Doney, S.C., V.J. Fabry, R.A. Feely, and J.A. Kleypas. 2009. Ocean Acidification: The Other CO<sub>2</sub> Problem. *Annual Review of Marine Science*. 1: 169-192.
- Fabry, V.J., B. A. Seibel, R.A. Feely and J.C. Orr. 2008. Impacts of ocean acidification on marine fauna and ecosystem processes. *Ices Journal of Marine Science*. 65 (3): 414-432.
- Feely, R.A., C.L.. Sabine, R.H. Byrne, F.J. Millero, A.G. Dickson, R. Wanninkhof, A. Murata, L. A. Miller, and D. Greeley. 2012. Decadal changes in the aragonite and calcite saturation state of the Pacific Ocean. *Global Biogeochemical Cycles*. 26: 1-15.
- Lalli, C.M., and R.W. Gilmer. 1989. Pelagic snails: the biology of holoplanktonic gastropod molluscs. Stanford University Press: Stanford, California.
- Little, W.S., and N.J. Copley. 2003. WHOI Silhouette DIGITIZER, Version 1.0, Users Guide. WHOI Technical Report: Woods Hole Oceanographic Institution.
- Rost, B., I. Zondervan, and D. Wolf-Gladrow. 2008. Sensitivity of phytoplankton to future changes in ocean carbonate chemistry: current knowledge, contradictions and research directions. *Marine Ecology Progress Series*. 373: 227-237.

# Questions?

---



Video courtesy of Nancy Copley and Alex Bergen