

stage 2-IV

In flowering plants, female meiosis occurs in the ovule, a specialized organ in the flower. A cell in the distal region of an ovule primordium, is specified at stage 2-I to be the megaspore mother cell (MMC). Subsequently, this MMC enlarges, differentiates, and undergoes meiosis in ovules at stage 2-IV. This process distinguishes the MMC and its lineage from the remaining non-sporogenous (NS) cells of the ovule primordium.

stage 2-I

In developing ovules, a dual role for ARP6 (ACTIN-RELATED PROTEIN 6) in regulating the spatial and temporal expression of DMC1 (DISRUPTED MEIOTIC cDNA 1), a gene involved in meiotic recombination, was established (Qin et al. 2014, Plant Cell 26: 1612-1628). ARP6 inhibits *DMC1* expression in non-sporogenous cells even as it promotes DMC1 expression in the MMC that is undergoing meiosis.

images of ovules carrying a promoterreporter (pDMC1:GFP) transgene (green), and counterstained with FM4-64 to label the plasma membranes of cells (red). A wildtype ovule (left image) normally expresses DMC1 exclusively in the MMC, whereas an arp6-mutant ovule (right image) does not express DMC1 in the MMC and instead shows dramatically increased expression in the non-sporogenous ovule cells.

Methods: Ovary walls of pistils (female reproductive structure) in fresh developing buds were dissected to expose ovule primordia and stained with FM4-64 dye solution (2 µM) for 5 minutes. Stained samples were mounted in 50% glycerol and observed on a TCS SP8 confocal microscope (Leica Microsystems). Green and red fluorescence of a single optical section of ovule primordia were acquired using Leica Application Suite Advanced Fluorescence software.



YUAN QIN

Center for Genomics and Biotechnology College of Crop Science Fujian Agriculture and Forestry University Fuzhou 350002, Fujian Province, China

RAMIN YADEGARI AND RAVISHANKAR PALANIVELU

School of Plant Sciences University of Arizona Tucson, AZ 85721, USA E-mail: rpalaniv@email.arizona.edu DOI: 10.1002/mrd.22472

stage I-II