A Camel-Milk Infant Formula for Reducing Necrotizing Enterocolitis Risk

A Grant-Development Scaffold

Heavy Metal Index
Independent evidence resource · heavymetalindex.com
May 2026

This is a grant-development scaffold: a structured starting draft for a small-business research grant (NIH SBIR, or a foundation or USDA equivalent) built around your camel-milk infant-formula product and the emerging science linking dietary nickel to necrotizing enterocolitis (NEC). It was drafted from the public scientific literature compiled at heavymetalindex.com, together with the published nickel-NEC hypothesis, and is offered to your company as a courtesy starting point.

Three things to be clear about at the outset. First, your company owns and authors it: everything here is a draft for you to revise, complete, and submit under your own name. Second, heavymetalindex.com is not a partner, preparer, or sub-awardee, and has no financial stake in your product; it is an independent evidence resource you are free to cite. Third, it is written to be honest, not promotional. As you will see, the literature shows camel milk is not automatically a clean, low-metal matrix, and we have built that fact into the proposal rather than around it, because a reviewer will find it immediately. The honest framing is also the stronger one.

How to use this scaffold

1. Fill the marked gaps. Wherever you see an italic note in the form [*to supply*: ...], that is content only you can provide (product data, team, budget, facilities). The full list appears in the closing section, *What your company must add*.
2. Pick a funding route. The default here is NIH SBIR (your company is the applicant; you subcontract the laboratory and animal work). NIH STTR is an option if you bring your own academic research partner. The Gerber Foundation is a strong topical fit for infant-nutrition work on a longer timeline. The science below serves all three; only the wrapper changes.
3. Have the science reviewed. The mechanistic claims rest on a recent hypothesis paper cited throughout. Your scientific team or an academic collaborator should pressure-test the Approach before submission.

Project Summary

[*to supply*: *company name and one line on the company and product stage*] is developing a camel-milk-based infant formula intended for infants at elevated risk of necrotizing enterocolitis (NEC), including the cow-milk-protein-allergic and preterm step-down populations. NEC is among the most lethal diseases of prematurity, and formula feeding is its leading modifiable risk factor. Camel milk is an attractive base because it lacks the cow-milk allergen beta-lactoglobulin and is rich in lactoferrin, immunoglobulins, and lysozyme, properties that approximate human milk's anti-infective profile. A 2026 metallomic synthesis implicates a previously overlooked variable in formula-associated NEC: dietary nickel, the obligate cofactor for the microbial enzymes urease, [NiFe]-hydrogenase, and glyoxalase I, which are virulence determinants of the Enterobacteriaceae that bloom in NEC. Human milk is nickel-poor; soy formula is nickel-rich. Critically, the published literature shows camel milk is itself a heavy-metal accumulation matrix whose nickel, lead, and cadmium vary by sourcing and are frequently elevated. The decision-relevant quantity is therefore bioavailable nickel, total nickel offset by the metal-binding proteins that may sequester it, which has never been measured in a camel-milk formula. This project determines, for our formula, whether its metal-binding proteins hold bioavailable nickel low, tests the nickel-virulence mechanism, and delivers the sourcing-and-surveillance controls that keep a camel-milk formula within safe limits lot to lot. Phase I establishes analytical and mechanistic feasibility; Phase II confirms it in a neonatal model and finalizes the product control protocol. [*to supply: company-specific objectives and any prior product or feasibility data*]

Project Narrative

Necrotizing enterocolitis is a frequently fatal intestinal disease of premature and vulnerable infants, and the type of formula an infant is fed changes that risk. This project tests whether a camel-milk infant formula, naturally rich in the protective protein lactoferrin, can be sourced and manufactured to keep its nickel and other heavy metals at levels that do not feed the gut bacteria implicated in the disease. The work delivers both the safety evidence and the quality-control protocol needed for such a formula to reach the infants who would benefit most.

Specific Aims

We hypothesize that net bioavailable nickel, not total nickel, governs the nickel-driven dysbiosis implicated in NEC, and that a camel-milk formula's abundant metal-binding proteins can hold bioavailable nickel low if and only if the product is sourced and processed to control its heavy-metal load. We will test this and build the controls to act on it.

Aim 1 — Quantify total and bioavailable nickel, and the heavy-metal panel, in our formula and its comparators. By ISO/IEC 17025 ICP-MS with arsenic speciation, measure nickel and the priority metal panel (Pb, Cd, inorganic As, total Hg, plus Ni, Sn, Al, Cr) on an as-consumed basis in our camel-milk formula [*to supply: formula and three or more source lots per region or herd*], pooled donor human milk, and representative cow- and soy-based formulas; partition nickel into protein-bound versus bioaccessible fractions (ultrafiltration plus infant-adapted simulated digestion). Expected: a defensible bioavailable-nickel figure for our formula relative to human milk and to cow and soy formula, and the sourcing covariates that drive it.

Aim 2 — Test whether nickel availability drives NEC-pathogen virulence, and whether our matrix suppresses it. In NEC-associated isolates (*E. coli*, *Klebsiella*, *Citrobacter*), measure nickel-dependent enzyme activity, growth, and stress survival across matched nickel delivered as free ion versus within the camel-milk matrix (Phase I), then compare our formula against cow and soy formula and a human-milk reference in an established neonatal NEC model [*to supply: animal-model CRO or academic facility*] on gut nickel, dysbiosis, and NEC severity (Phase II). Expected: a direct test of whether our formula's matrix lowers nickel-enabled virulence at equal nickel.

Aim 3 — Deliver a biomarker and a certification-grade control protocol. Evaluate fecal urease and ammonia, and fecal nickel, as non-invasive readouts of the nickel-driven shift, and convert the Aim 1 and Aim 2 findings into a sourcing specification and lot-testing protocol (as-consumed limits, decision rules under measurement uncertainty, accredited-laboratory and chain-of-custody requirements), drawing on the public heavy-metal-testing methodology at heavymetalindex.com. Expected: a monitorable readout and the quality-control blueprint that keeps the finished product within safe bioavailable-nickel bounds.

This project is designed to be decision-useful regardless of outcome: a favorable bioavailability result supports our formula as a low-effective-nickel option; an unfavorable one defines the precise sourcing and remediation targets required to make it safe.

Significance

Necrotizing enterocolitis (NEC) is among the most lethal diseases of prematurity, with case-fatality of 15 to 30 percent and lifelong sequelae in survivors. Its single most consistent modifiable risk factor is enteral feeding type: preterm infants fed bovine- or soy-based formula develop NEC at substantially higher rates than those fed human milk[2]. Human milk's protection is conventionally attributed to what it contains and what formula lacks; a complementary, underexamined possibility is that formula contains a harmful constituent.

A 2026 metallomic synthesis[1] proposes that constituent is nickel. Nickel is not a human nutrient, but it is the obligate cofactor for three microbial enzyme systems, urease, [NiFe]-hydrogenase, and glyoxalase I, that are fitness and virulence determinants of the Enterobacteriaceae (*E. coli*, *Klebsiella*, *Citrobacter*) that bloom in NEC. Urease raises local pH and frees ammonia, damaging mucosa and disabling acid-producing commensals; hydrogenase lets pathogens harvest fermentative hydrogen; glyoxalase I detoxifies methylglyoxal under stress[3]. Human milk is naturally nickel-poor (5 to 16 µg/L), whereas soy formula delivers about 450 µg/L and bovine formula about 30 µg/L[1, 9, 10], an order-of-magnitude nickel bolus. The host defends with metal-sequestering proteins (calprotectin, and possibly lactoferrin) that withhold nickel from microbial enzymes; a high dietary nickel load may overwhelm that defense. Interfering with bacterial nickel uptake suppresses these virulence enzymes simultaneously[4], establishing nickel availability as a tractable lever.

Camel milk is a compelling non-human base for a hypoallergenic preterm formula. It lacks beta-lactoglobulin, the dominant cow-milk allergen; its protein is largely beta-casein, which is more readily digested;

and it carries roughly five-fold the immunoglobulin of cow milk plus elevated lactoferrin, lysozyme, and vitamin C[7, 11, 12]. These are precisely the human-milk-like, anti-infective properties a NEC-risk formula should mimic.

Camel milk is, however, a documented heavy-metal accumulation matrix. A 2024 global systematic review (20 studies, 787 samples) found camel-milk lead exceeded the Codex limit (0.020 mg/kg) in 10 of 18 studies, with Egypt and Saudi Arabia reporting the highest lead, nickel, and cadmium[5]. A five-species ICP-MS comparison found raw camel milk carried the highest nickel of any dairy species measured, 131 µg/kg, above cow (82) and goat (38)[6]. In polluted regions, camel-milk lead averaged 250 µg/kg, about 12 times the Codex limit, tracking proximity to industry and traffic[7] yet in unpolluted rural settings the same metals sat far below limits[8]. The decisive pattern is that camel-milk metal load is driven by environmental sourcing, not species physiology.

Under the nickel hypothesis, an uncontrolled camel-milk formula could deliver more, not less, bioavailable nickel than the bovine formula it is meant to improve upon, even as its metal-binding proteins may sequester that nickel. No study has measured bioavailable (versus total) nickel in a camel-milk formula. The question is consequential beyond this product: the regulatory and litigation environment for infant-food metals has hardened sharply (FDA Closer to Zero action levels[17] California AB899 mandatory testing and disclosure[18] the 2026 Supreme Court decision in Hain v. Palmquist[16] pushing heavy-metal claims into state courts), so any infant formula now needs a defensible, contemporaneous, independent contaminant record. This project generates the science and the control strategy together.

Innovation

This project is innovative on four counts. Conceptually, it operationalizes a reframing of NEC as partly a nutritional-toxicology problem (dietary nickel licensing microbial virulence), beyond probiotic and human-milk-fortifier strategies. Analytically, it produces the first characterization of total versus bioavailable nickel in a camel-milk formula, the metric the hypothesis actually implicates, which prior total-concentration surveys cannot supply. Mechanistically, it tests a specific, falsifiable proposition, that camel milk's protein-bound metal fraction suppresses nickel-dependent pathogen virulence relative to equivalent free nickel, rather than treating camel milk as generically better. Translationally, it embeds certification-grade quality control (ISO/IEC 17025 lot testing, as-consumed limits, decision rules under measurement uncertainty, sourcing qualification) as an integral deliverable, so a favorable result is immediately convertible into a defensible product.

Approach

Design and phasing. The work is structured for SBIR; your company subcontracts the analytical, bioavailability, and animal-model work to qualified laboratories. Phase I (about 12 months) executes Aim 1 in full, the in-vitro component of Aim 2, and the Aim 3 biomarker proof-of-concept. Phase II (about 24 months) executes the in-vivo NEC-model component of Aim 2 and the Aim 3 surveillance protocol. All concentrations are handled on an as-consumed basis (powder reconstituted at label dilution) so comparisons across human milk, camel formula, and bovine and soy formula are like-for-like.

Test article and matrices. Your company supplies the camel-milk formula and candidate raw-milk source lots [*to supply: formula specification, source-lot provenance*] and engages ISO/IEC 17025 laboratories and an animal-model facility [*to supply: subcontractor laboratories or CRO*]. Comparators are pooled donor human milk and representative commercial bovine and soy formulas.

Aim 1 — Total and bioavailable nickel plus the heavy-metal panel. (1a) Quantify nickel and the priority panel by ISO/IEC 17025 ICP-MS with arsenic speciation, as-consumed, across the four matrices and three or more source lots per region, with quantitation limits at or below the relevant thresholds. (1b) Partition nickel into protein-bound versus free fractions (ultrafiltration, size-exclusion, ICP-MS) and estimate gut-available nickel by an infant-adapted simulated-digestion bioaccessibility assay. Working hypothesis: the formula's abundant metal-binding proteins yield bioavailable nickel below soy formula; the magnitude is the unknown. Host nickel-sequestration is best established for calprotectin, so whether milk lactoferrin and caseins bind nickel equivalently is itself unresolved; Aim 1b therefore measures the bound fraction empirically rather than assuming the agent. (1c) Relate source-lot nickel and lead to provenance (region, industrial or traffic proximity, fodder) to define the sourcing constraints a certifiable low-nickel camel formula requires.

Aim 2 — Nickel-dependence of pathogen virulence by matrix. (2a, Phase I) In NEC isolates, measure urease, [NiFe]-hydrogenase, and glyoxalase-I activity, growth, and acid and oxidative-stress survival across defined free-nickel concentrations and across the Aim 1 matrices at matched total nickel. Decisive contrast: matched total nickel as free nickel ion versus within the camel matrix; if matrix sequestration is protective, virulence-enzyme activity will be lower at equal nickel, and selective depletion of candidate binders (lactoferrin, caseins) identifies which fraction is responsible. (2b, Phase II) In an established neonatal NEC model (rat pup or preterm piglet), compare the formula against cow and soy formula and a human-milk reference on gut nickel, nickel-enzyme expression, microbiome composition, mucosal injury, and NEC incidence and severity, powered for a meaningful difference in severity with blinded scoring.

Aim 3 — Biomarker plus control protocol. (3a) Evaluate fecal urease and ammonia, and fecal nickel, as non-invasive readouts of the nickel-driven shift. (3b) Convert findings into a sourcing-and-surveillance specification: as-consumed bioavailable-nickel and heavy-metal limits, qualifying sourcing constraints, lot-testing frequency and two-sided decision rules under measurement uncertainty, and chain-of-custody and accreditation requirements.

Pitfalls and alternatives. If the camel matrix shows no virulence suppression at matched nickel, the product rationale narrows to immunological benefit and the project pivots emphasis to Aim 1c sourcing and remediation. If the animal model is underpowered for incidence, mechanistic readouts serve as primary endpoints. Processing effects on camel lactoferrin integrity, a known variable, are controlled by characterizing the finished formula, not raw milk.

Commercialization Bridge

The commercial product is a low-bioavailable-nickel, heavy-metal-controlled camel-milk infant formula positioned for the cow-milk-protein-allergy and NEC-risk step-down segments, markets currently served by costly extensively-hydrolyzed and amino-acid formulas. The defensible contaminant record and control protocol from Aim 3 are the durable differentiator in a market under intensifying regulatory and litigation pressure. Phase I de-risks the core scientific claim; Phase II yields the product specification and surveillance protocol that support market entry. [*to supply: market sizing, pricing, distribution, intellectual property, and the full SBIR commercialization plan*]

Honest Limitations

- The nickel-NEC hypothesis is, to date, a published but preprint-stage synthesis of circumstantial evidence; this project is among the first direct tests, and is anchored by the well-established formula-NEC and camel-milk-hypoallergenic literatures so it does not stand alone.

- The sequestration rationale (that camel milk's metal-binding proteins lower bioavailable nickel) is plausible but unproven, and the specific role of milk lactoferrin in nickel binding is itself uncertain; calprotectin is the better-established nickel chelator in host defense. Aim 1b and Aim 2a measure the effect empirically and refute it cleanly if wrong.
- A camel-milk infant formula requires nutritional fortification, and any marketed US infant formula must independently satisfy FDA infant-formula requirements (Infant Formula Act; 21 CFR 106-107), including nutritional adequacy and premarket notification. This project addresses the heavy-metal axis only; nutritional adequacy, clinical efficacy, and regulatory clearance are separate steps your company owns.

What Your Company Must Add

The scientific spine above is drafted; the items below are outside what the public literature can supply and must come from your company before submission.

Component	What to supply
Company identity, product stage, prior data	Name, history, pilot or feasibility results, intellectual-property position
Test article and source lots	Finished formula specification, certificates of analysis, three or more source lots per region with provenance
Subcontractors	ISO/IEC 17025 laboratory or laboratories; animal-model CRO or academic facility
Investigators and biosketches	Principal investigator and key-personnel bios, expertise, effort
Budget and justification	Personnel, subcontracts, supplies, indirect costs
Facilities, resources, equipment	Your and your subcontractors' environment
Vertebrate-animals section	Welfare, IACUC approval, justification (Phase II)
Data-management and sharing plan	Repositories, access terms
Letters of support	Donor-milk bank, CRO, clinical collaborators
SBIR eligibility and commercialization plan	Small-business certification; full commercialization plan, market data
FDA infant-formula regulatory pathway	Infant Formula Act and 21 CFR 106-107 notification plus nutritional adequacy, a separate regulatory track

References

[1] Pendergrass K. Nickel as a Catalytic Driver of Necrotizing Enterocolitis. Microbiome Medicine, 2026. doi:10.5281/zenodo.18200348.

[2] Shulhan J, Dicken B, Hartling L, Larsen BM. Current Knowledge of NEC in Preterm Infants and the Impact of Different Types of Enteral Nutrition Products. Adv Nutr 2017;8(1):80-91. doi:10.3945/an.116.013193.

[3] Maier RJ, Benoit SL. Role of Nickel in Microbial Pathogenesis. Inorganics 2019;7(7):80. doi:10.3390/inorganics7070080.

[4] Sychantha D, Chen X, Koteva K, et al. Targeting bacterial nickel transport with aspergillomarasmine A suppresses virulence-associated Ni-dependent enzymes. Nat Commun 2024;15:4036. doi:10.1038/s41467-024-48232-1.

[5] Kerdoun MA, Djafer R. Heavy metal levels in camel milk and health risk assessment: a global systematic review. J Trace Elem Miner 2024. doi:10.1016/j.jtemin.2024.100131.

[6] Chen L, Li X, Li Z, Deng L. Analysis of 17 elements in cow, goat, buffalo, yak, and camel milk by ICP-MS. RSC Adv 2020;10:6736-6742. doi:10.1039/dora00390e.

[7] Konuspayeva G, Faye B, Loiseau G, Diacono E, Akhmetsadykova S. Pollution of Camel Milk by Heavy Metals in Kazakhstan. Open Environ Pollut Toxicol J 2010;1:112-118. doi:10.2174/1876397900901010112.

[8] Yazdanian Ghahfarokhi E, et al. Heavy metal contamination in camel milk. Vet Med Sci 2025. doi:10.1002/vms3.70114.

[9] Pandelova M, et al. Ca, Cd, Cu, Fe, Hg, Mn, Ni, Pb, Se and Zn contents in baby foods from the EU market. J Food Compos Anal 2012;27(2):120-127. doi:10.1016/j.jfca.2012.04.011.

[10] Biego GH, et al. Mineral and contaminant content of infant formulas; soy-formula nickel about 0.45 mg/L. 1998. (citation detail to confirm)

[11] El-Agamy EI, et al. Purification and characterization of lactoferrin, lactoperoxidase, lysozyme and immunoglobulins from camel milk. Int Dairy J 1996;6:129-145.

[12] Konuspayeva G, et al. Lactoferrin and immunoglobulin content in camel milk from Kazakhstan. J Dairy Res 2007.

[16] Hain Celestial Group, Inc. v. Palmquist, 607 U.S. (2026) (No. 24-724).

[17] U.S. Food and Drug Administration. Action Levels for Lead in Food Intended for Babies and Young Children (Closer to Zero), 2025.

[18] California Assembly Bill 899, 2023.

[19] Codex Alimentarius. Maximum level for lead in milk, 0.020 mg/kg, CXS 193-1995.

Prepared as a courtesy starting scaffold from the public literature at heavymetalindex.com. Your company owns and authors any resulting application. heavymetalindex.com is an independent reference and not a party to your submission. Questions about the underlying literature: karen@paleofoundation.com.