End-Stage Renal Disease (ESRD)

Dr. Umar

Microbiome Signatures, Microbiome Medicine

Abstract. ESRD is irreversible kidney failure needing dialysis or transplant, driven by nephron loss, inflammation, and microbiome-derived toxins.

Published 2025-12-02 · https://microbiomemedicine.com/conditions/end-stage-renal-disease-esrd · A Microbiome Signatures condition monograph

Overview

End-stage renal disease (ESRD) is the final, irreversible phase of chronic kidney disease (CKD), in which kidney function is insufficient to sustain life without renal replacement therapy.[1] In practice, ESRD corresponds to CKD stage 5 (usually defined by an estimated glomerular filtration rate (eGFR) below 15 mL/min/1.73 m²) and is characterized by the need for dialysis or kidney transplantation to manage life-threatening uremia.[2] This distinguishes ESRD from late-stage CKD (stage 4 and early stage 5), where eGFR is severely reduced (e.g., 15–29 mL/min) but patients may not yet require dialysis if asymptomatic. ESRD carries a grave prognosis: it is a major health burden worldwide with high morbidity and mortality. For example, in the United States over 600,000 individuals live with ESRD, yet the 5-year survival on dialysis is only ~50%[3] – a survival rate comparable to many cancers. The first year of dialysis is especially perilous (mortality up to 20–30%).[4] Kidney transplantation dramatically improves outlook, with transplanted patients achieving about 80% five-year survival (versus <50% on dialysis) as noted in clinical registries.[5] Epidemiologically, the prevalence of ESRD has steadily risen due to increasing rates of diabetes and hypertension globally, and CKD is projected to rank among the top causes of years of life lost by 2040. Overall, ESRD confers a markedly reduced life expectancy and heavy symptom burden, making prevention of progression and timely treatment crucial.

Causes

Diagnosis

Primer

The interplay between impaired metal handling, dysbiosis-driven toxin generation, and global metabolomic disruption in ESRD illustrates how the metallome, microbiome, and metabolome converge on shared pathways of inflammation, oxidative stress, and fibrosis. Loss of renal excretory capacity alters circulating metal availability, reshapes gut microbial communities, and permits accumulation of toxic microbial metabolites such as indoxyl sulfate and p-cresyl sulfate. These combined disturbances amplify endothelial dysfunction, immune activation, and intestinal barrier injury, underscoring the potential of microbiome-

based diagnostics and targeted interventions (MBTIs) to detect and modulate pathophysiologic processes that traditional renal markers cannot fully capture.

Etiology

ESRD is the common endpoint of various chronic nephropathies. Diabetic nephropathy (from long-standing diabetes mellitus) is the single leading cause of ESRD worldwide, accounting for roughly 40–50% of new ESRD cases in many regions.[6]

Pathophysiology

ESRD represents the end result of relentless nephron loss and maladaptive repair processes in the kidneys. As functioning nephron mass declines, the remaining nephrons undergo compensatory hyperfiltration: surviving glomeruli dilate and raise single-nephron GFR to uphold overall filtration. In the short term, this preserves kidney function, but chronically, the intraglomerular hypertension and stretch injury lead to capillary wall damage, podocyte loss, and ultimately segmental sclerosis of glomeruli.[7] This phenomenon (described in the Brenner hyperfiltration hypothesis) creates a vicious cycle of progressive nephron destruction. Hemodynamic maladaptations are largely mediated by the renin–angiotensin–aldosterone system (RAAS). Decreased renal perfusion triggers renin release and angiotensin II formation, which preferentially constricts efferent arterioles to maintain filtration pressure. However, angiotensin II also promotes inflammatory and fibrogenic pathways (via TGF-β1 and others), accelerating renal scarring.[8] Indeed, persistent RAAS activation is a central driver of CKD progression, and RAAS inhibitors (ACEis/ARBs) slow decline by lowering glomerular pressure and fibrosis.[9]

Metabolomic Signature of ESRD

ESRD is marked by a distinct metabolomic signature characterized by the extreme accumulation of protein-bound uremic solutes, nitrogenous waste products, organic acids, and inflammatory metabolites. Indoxyl sulfate and p-cresyl sulfate—derived from microbial metabolism of tryptophan and tyrosine—rise to levels dozens of times higher than in early CKD because damaged kidneys cannot excrete these protein-bound toxins efficiently, and dialysis removes them poorly. These metabolites drive oxidative stress, endothelial injury, vascular stiffness, left ventricular hypertrophy, and fibrotic signaling within the kidney. Parallel increases in endotoxin, or lipopolysaccharide (LPS), advanced glycation end-products, and middle molecules further promote systemic inflammation and cardiovascular risk, establishing a metabolomic milieu tightly linked to mortality and complications in dialysis patients.

Nutritional Immunity

Nutritional immunity becomes profoundly dysregulated in ESRD. Chronic inflammation elevates hepcidin, restricting iron availability for erythropoiesis while paradoxically increasing iron deposition in tissues and fostering microbial dysbiosis. Zinc deficiency, common in ESRD due to impaired absorption and inflammation-mediated sequestration, compromises host tight-junction integrity and antimicrobial defense, promoting barrier disruption and bacterial translocation. Copper handling likewise becomes erratic, contributing to oxidative stress and impaired innate immune signaling. The net effect is a disrupted metal-immune balance that heightens infection susceptibility, sustains systemic inflammation, and reinforces the gut–kidney axis disturbances characteristic of uremia.

Mismetallation in ESRD

The uremic milieu, chronic inflammation, and impaired renal excretion promote the accumulation of circulating toxins and metabolic by-products, many of which disrupt normal protein structure and enzymatic function in a manner analogous to toxic metal substitution. Protein-bound uremic solutes such as indoxyl sulfate and p-cresyl sulfate rise to extremely high concentrations because diseased kidneys cannot secrete them effectively,[10] and they bind tightly to albumin, preventing efficient dialysis clearance.[11] Their capacity to induce oxidative stress, inflammation, endothelial dysfunction, and mitochondrial injury[12][13] mirrors the biochemical consequences of mismetallation, wherein misplaced or excessive metals impair protein stability and cellular redox balance. Through oxidative injury, IS and PCS activate profibrotic pathways including TGF-β1 and epithelial-to-mesenchymal transition,[14] illustrating how metabolic overload in ESRD functionally produces effects similar to metal dysregulation: protein damage, loss of structural fidelity, and activation of maladaptive signaling cascades. Thus, while not a metal itself, the toxin-driven oxidative microenvironment in ESRD creates conditions under which mismetallation-like injury pathways become pathophysiologically relevant.

Gut–Kidney Axis

The gut–kidney axis reflects the bidirectional relationship between renal failure and microbial dysbiosis. In ESRD, high levels of uremic waste in the blood and gastrointestinal tract, combined with low-fiber dietary restrictions, shift the microbiome toward proteolytic species that generate excess nitrogenous and aromatic metabolites. This results in markedly elevated circulating levels of indoxyl sulfate and p-cresyl sulfate—often dozens-fold above normal.[15] These validated uremic toxins cause oxidative stress, inflammation, endothelial dysfunction and contribute to cardiovascular pathology, including vascular calcification and left ventricular hypertrophy.[16] IS also reduces nitric oxide and promotes fibrotic signaling in the kidney,[17] while PCS activates leukocytes and drives chronic inflammation.[18] Since both toxins bind tightly to albumin, they are poorly cleared by dialysis and accumulate further in ESRD.[19] Dysbiosis simultaneously weakens the intestinal barrier: uremia induces mucosal atrophy and disrupts tight junction proteins, facilitating the translocation of endotoxin and microbial fragments into the bloodstream.[20] This “leaky gut” syndrome amplifies systemic inflammation and contributes to malnutrition, atherosclerosis, and inflammatory anemia. The resulting feedback loop—renal failure worsening dysbiosis, dysbiosis generating toxins and endotoxin, and these factors intensifying systemic and cardiovascular injury—reinforces the central role of the gut–kidney axis in ESRD pathophysiology.[21]

Interventions

ESRD requires more than replacing filtration—it demands direct intervention against the systemic offenders driving inflammation, oxidative stress, toxin accumulation, and metallomic imbalance. Microbiome-derived uremic toxins, dysfunctional RAAS signaling, chronic acidosis, dysregulated iron metabolism, zinc depletion, copper elevation, endotoxin translocation, and toxic solute retention collectively maintain a biochemical environment that accelerates cardiovascular and renal injury. Interventions must therefore combine dialysis adequacy, pharmacologic therapy, nutritional strategies, and targeted management of metallomic and microbiome-toxin burdens, all supported by the mechanistic pathways described in the referenced literature.

FAQs

Research Feed

KDIGO 2012 Clinical Practice Guideline for the Evaluation and Management of Chronic Kidney Disease
Chronic Kidney Disease (CKD) •
Chronic Kidney Disease (CKD)

Did you know?
The bidirectional relationship between diabetes and CKD creates a vicious cycle in which each condition perpetuates the other. Diabetic patients who develop CKD show accelerated progression of kidney disease and markedly increased cardiovascular event rates compared to nondiabetic CKD populations.

Alias iure reprehenderit aut accusantium. Molestiae dolore suscipit. Necessitatibus eum quaerat. Repudiandae suscipit quo necessitatibus. Voluptatibus ullam nulla temporibus nobis. Atque eaque sed totam est assumenda. Porro modi soluta consequuntur veritatis excepturi minus delectus reprehenderit est. Eveniet labore ut quas minima aliquid quibusdam. Vitae possimus fuga praesentium eveniet debitis exercitationem deleniti.

Create a free account to unlock this study summary.
Microbiome Insiders can read two study summaries for any topic on Microbiome.
(Get started with your free account)

Get started with Microbiome+ today.
Causes of Death in End-Stage Kidney Disease: Comparison Between the United States Renal Data System and a Large Integrated Health Care System
Chronic Kidney Disease (CKD) •
Chronic Kidney Disease (CKD)

Did you know?
The bidirectional relationship between diabetes and CKD creates a vicious cycle in which each condition perpetuates the other. Diabetic patients who develop CKD show accelerated progression of kidney disease and markedly increased cardiovascular event rates compared to nondiabetic CKD populations.

End-Stage Renal Disease (ESRD) •
End-Stage Renal Disease (ESRD)

Did you know?
*Some uremic toxins produced by gut bacteria, like indoxyl sulfate, can reach blood concentrations in ESRD that are more than **50 times higher** than in healthy individuals — yet dialysis removes almost none of them because they bind so tightly to albumin.*

Alias iure reprehenderit aut accusantium. Molestiae dolore suscipit. Necessitatibus eum quaerat. Repudiandae suscipit quo necessitatibus. Voluptatibus ullam nulla temporibus nobis. Atque eaque sed totam est assumenda. Porro modi soluta consequuntur veritatis excepturi minus delectus reprehenderit est. Eveniet labore ut quas minima aliquid quibusdam. Vitae possimus fuga praesentium eveniet debitis exercitationem deleniti.

Create a free account to unlock this study summary.

Microbiome Insiders can read two study summaries for any topic on Microbiome.
(Get started with your free account)

Get started with Microbiome+ today.
Risk of rapid progression to dialysis in patients with type 2 diabetes mellitus with and without diabetes-related complications at diagnosis

Alias iure reprehenderit aut accusantium. Molestiae dolore suscipit. Necessitatibus eum quaerat. Repudiandae suscipit quo necessitatibus. Voluptatibus ullam nulla temporibus nobis. Atque eaque sed totam est assumenda. Porro modi soluta consequuntur veritatis excepturi minus delectus reprehenderit est. Eveniet labore ut quas minima aliquid quibusdam. Vitae possimus fuga praesentium eveniet debitis exercitationem deleniti.

Create a free account to unlock this study summary.

Microbiome Insiders can read two study summaries for any topic on Microbiome.
(Get started with your free account)

Get started with Microbiome+ today.
End-stage renal disease

Alias iure reprehenderit aut accusantium. Molestiae dolore suscipit. Necessitatibus eum quaerat. Repudiandae suscipit quo necessitatibus. Voluptatibus ullam nulla temporibus nobis. Atque eaque sed totam est assumenda. Porro modi soluta consequuntur veritatis excepturi minus delectus reprehenderit est. Eveniet labore ut quas minima aliquid quibusdam. Vitae possimus fuga praesentium eveniet debitis exercitationem deleniti.

Create a free account to unlock this study summary.

Microbiome Insiders can read two study summaries for any topic on Microbiome.
(Get started with your free account)

Get started with Microbiome+ today.
Uremic toxins induce kidney fibrosis by activating intrarenal renin–angiotensin–aldosterone system associated epithelial-to-mesenchymal transition

Alias iure reprehenderit aut accusantium. Molestiae dolore suscipit. Necessitatibus eum quaerat. Repudiandae suscipit quo necessitatibus. Voluptatibus ullam nulla temporibus nobis. Atque eaque sed totam est assumenda. Porro modi soluta consequuntur veritatis excepturi minus delectus reprehenderit est. Eveniet labore ut quas minima aliquid quibusdam. Vitae possimus fuga praesentium eveniet debitis exercitationem deleniti.

Create a free account to unlock this study summary.

Microbiome Insiders can read two study summaries for any topic on Microbiome.
(Get started with your free account)

Get started with Microbiome+ today.
Chronic kidney disease and the gut microbiome

Alias iure reprehenderit aut accusantium. Molestiae dolore suscipit. Necessitatibus eum quaerat. Repudiandae suscipit quo necessitatibus. Voluptatibus ullam nulla temporibus nobis. Atque eaque sed

totam est assumenda. Porro modi soluta consequuntur veritatis excepturi minus delectus reprehenderit est. Eveniet labore ut quas minima aliquid quibusdam. Vitae possimus fuga praesentium eveniet debitis exercitationem deleniti.

Create a free account to unlock this study summary.

Microbiome Insiders can read two study summaries for any topic on Microbiome.
(Get started with your free account)

Get started with Microbiome+ today.

Modular Strategies for Nephron Replacement and Clinical Translation

Alias iure reprehenderit aut accusantium. Molestiae dolore suscipit. Necessitatibus eum quaerat. Repudiandae suscipit quo necessitatibus. Voluptatibus ullam nulla temporibus nobis. Atque eaque sed totam est assumenda. Porro modi soluta consequuntur veritatis excepturi minus delectus reprehenderit est. Eveniet labore ut quas minima aliquid quibusdam. Vitae possimus fuga praesentium eveniet debitis exercitationem deleniti.

Create a free account to unlock this study summary.

Microbiome Insiders can read two study summaries for any topic on Microbiome.
(Get started with your free account)

Get started with Microbiome+ today.

Intestinal microbiota in pediatric patients with end stage renal disease: a Midwest Pediatric Nephrology Consortium study

Alias iure reprehenderit aut accusantium. Molestiae dolore suscipit. Necessitatibus eum quaerat. Repudiandae suscipit quo necessitatibus. Voluptatibus ullam nulla temporibus nobis. Atque eaque sed totam est assumenda. Porro modi soluta consequuntur veritatis excepturi minus delectus reprehenderit est. Eveniet labore ut quas minima aliquid quibusdam. Vitae possimus fuga praesentium eveniet debitis exercitationem deleniti.

Create a free account to unlock this study summary.

Microbiome Insiders can read two study summaries for any topic on Microbiome.
(Get started with your free account)

Get started with Microbiome+ today.

Update History

2025-12-02 11:00:28

End-Stage Renal Disease (ESRD) major
published

1
Chronic Kidney Disease (CKD)
Dysbiosis in chronic kidney disease (CKD) reflects a shift toward reduced beneficial taxa and increased pathogenic, uremic toxin-producing species, driven by a bidirectional interaction in which the uremic environment disrupts microbial composition and dysbiotic metabolites accelerate renal deterioration.

Microbiome-Targeted Interventions (MBTIs)
Microbiome Targeted Interventions (MBTIs) are cutting-edge treatments that utilize information from Microbiome Signatures to modulate the microbiome, revolutionizing medicine with unparalleled precision and impact.

Lipopolysaccharide (LPS)
Lipopolysaccharide (LPS), a potent endotoxin present in the outer membrane of Gram-negative bacteria that causes chronic immune responses associated with inflammation.

Nutritional Immunity
Nutritional immunity restricts metal access to pathogens, leveraging sequestration, transport, and toxicity to control infections and immunity.

Zinc
Zinc is an essential trace element vital for cellular functions and microbiome health. It influences immune regulation, pathogen virulence, and disease progression in conditions like IBS and breast cancer. Pathogens exploit zinc for survival, while therapeutic zinc chelation can suppress virulence, rebalance the microbiome, and offer potential treatments for inflammatory and degenerative diseases.

Chronic Kidney Disease (CKD)
Dysbiosis in chronic kidney disease (CKD) reflects a shift toward reduced beneficial taxa and increased pathogenic, uremic toxin-producing species, driven by a bidirectional interaction in which the uremic environment disrupts microbial composition and dysbiotic metabolites accelerate renal deterioration.

Chronic Kidney Disease (CKD)
Dysbiosis in chronic kidney disease (CKD) reflects a shift toward reduced beneficial taxa and increased pathogenic, uremic toxin-producing species, driven by a bidirectional interaction in which the uremic environment disrupts microbial composition and dysbiotic metabolites accelerate renal deterioration.

End-Stage Renal Disease (ESRD)
End-stage renal disease is the irreversible loss of kidney function marked by uremic toxin accumulation, systemic complications, and the need for dialysis or transplantation. Its pathophysiology involves nephron loss, inflammation, metabolic disruption, and microbiome-derived toxins that accelerate cardiovascular and immune dysfunction.

References

1. KDIGO 2012 Clinical Practice Guideline for the Evaluation and Management of Chronic Kidney Disease. Kidney Disease: Improving Global Outcomes (KDIGO) CKD Work Group.. (Kidney International Supplements. 2013)
2. KDIGO 2012 Clinical Practice Guideline for the Evaluation and Management of Chronic Kidney Disease. Kidney Disease: Improving Global Outcomes (KDIGO) CKD Work Group.. (Kidney International Supplements. 2013)
3. Causes of Death in End-Stage Kidney Disease: Comparison Between the United States Renal Data System and a Large Integrated Health Care System.. Bhandari SK, Zhou H, Shaw SF, Shi J, Tilluckdharry NS, Rhee CM, Jacobsen SJ, Sim JJ.. (American Journal of Nephrology. 2022)
4. Causes of Death in End-Stage Kidney Disease: Comparison Between the United States Renal Data System and a Large Integrated Health Care System.. Bhandari SK, Zhou H, Shaw SF, Shi J, Tilluckdharry NS, Rhee CM, Jacobsen SJ, Sim JJ.. (American Journal of Nephrology. 2022)

5. Causes of Death in End-Stage Kidney Disease: Comparison Between the United States Renal Data System and a Large Integrated Health Care System.. Bhandari SK, Zhou H, Shaw SF, Shi J, Tilluckdharry NS, Rhee CM, Jacobsen SJ, Sim JJ.. (American Journal of Nephrology. 2022)
6. Risk of rapid progression to dialysis in patients with type 2 diabetes mellitus with and without diabetes-related complications at diagnosis.. Shih HM, Tsai WC, Wu PY, Chiu LT, Kung PT.. (Scientific Reports. 2023)
7. End-stage renal disease.. Rout P, Aslam A.. (In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2025 Jun 22)
8. Uremic toxins induce kidney fibrosis by activating intrarenal renin–angiotensin–aldosterone system associated epithelial-to-mesenchymal transition.. Sun CY, Chang SC, Wu MS.. (PLoS One. 2012)
9. Uremic toxins induce kidney fibrosis by activating intrarenal renin–angiotensin–aldosterone system associated epithelial-to-mesenchymal transition.. Sun CY, Chang SC, Wu MS.. (PLoS One. 2012)
10. Chronic kidney disease and the gut microbiome. Hobby GP, Karaduta O, Dusio GF, Singh M, Zybailov BL, Arthur JM.. (Am J Physiol Renal Physiol. 2019)
11. Chronic kidney disease and the gut microbiome. Hobby GP, Karaduta O, Dusio GF, Singh M, Zybailov BL, Arthur JM.. (Am J Physiol Renal Physiol. 2019)
12. Uremic toxins induce kidney fibrosis by activating intrarenal renin–angiotensin–aldosterone system associated epithelial-to-mesenchymal transition.. Sun CY, Chang SC, Wu MS.. (PLoS One. 2012)
13. Chronic kidney disease and the gut microbiome. Hobby GP, Karaduta O, Dusio GF, Singh M, Zybailov BL, Arthur JM.. (Am J Physiol Renal Physiol. 2019)
14. Uremic toxins induce kidney fibrosis by activating intrarenal renin–angiotensin–aldosterone system associated epithelial-to-mesenchymal transition.. Sun CY, Chang SC, Wu MS.. (PLoS One. 2012)
15. Chronic kidney disease and the gut microbiome. Hobby GP, Karaduta O, Dusio GF, Singh M, Zybailov BL, Arthur JM.. (Am J Physiol Renal Physiol. 2019)
16. Chronic kidney disease and the gut microbiome. Hobby GP, Karaduta O, Dusio GF, Singh M, Zybailov BL, Arthur JM.. (Am J Physiol Renal Physiol. 2019)
17. Uremic toxins induce kidney fibrosis by activating intrarenal renin–angiotensin–aldosterone system associated epithelial-to-mesenchymal transition.. Sun CY, Chang SC, Wu MS.. (PLoS One. 2012)
18. Chronic kidney disease and the gut microbiome. Hobby GP, Karaduta O, Dusio GF, Singh M, Zybailov BL, Arthur JM.. (Am J Physiol Renal Physiol. 2019)
19. Chronic kidney disease and the gut microbiome. Hobby GP, Karaduta O, Dusio GF, Singh M, Zybailov BL, Arthur JM.. (Am J Physiol Renal Physiol. 2019)
20. Chronic kidney disease and the gut microbiome. Hobby GP, Karaduta O, Dusio GF, Singh M, Zybailov BL, Arthur JM.. (Am J Physiol Renal Physiol. 2019)
21. Chronic kidney disease and the gut microbiome. Hobby GP, Karaduta O, Dusio GF, Singh M, Zybailov BL, Arthur JM.. (Am J Physiol Renal Physiol. 2019)

Kidney Disease: Improving Global Outcomes (KDIGO) CKD Work Group.

KDIGO 2012 Clinical Practice Guideline for the Evaluation and Management of Chronic Kidney Disease Kidney International Supplements. 2013

Read Review
Kidney Disease: Improving Global Outcomes (KDIGO) CKD Work Group.

KDIGO 2012 Clinical Practice Guideline for the Evaluation and Management of Chronic Kidney Disease Kidney International Supplements. 2013

Read Review
Bhandari SK, Zhou H, Shaw SF, Shi J, Tilluckdharry NS, Rhee CM, Jacobsen SJ, Sim JJ.

Causes of Death in End-Stage Kidney Disease: Comparison Between the United States Renal Data System and a Large Integrated Health Care System.
American Journal of Nephrology. 2022

Read Review
Bhandari SK, Zhou H, Shaw SF, Shi J, Tilluckdharry NS, Rhee CM, Jacobsen SJ, Sim JJ.

Causes of Death in End-Stage Kidney Disease: Comparison Between the United States Renal Data System and a Large Integrated Health Care System.
American Journal of Nephrology. 2022

Read Review
Bhandari SK, Zhou H, Shaw SF, Shi J, Tilluckdharry NS, Rhee CM, Jacobsen SJ, Sim JJ.

Causes of Death in End-Stage Kidney Disease: Comparison Between the United States Renal Data System and a Large Integrated Health Care System.
American Journal of Nephrology. 2022

Read Review
Shih HM, Tsai WC, Wu PY, Chiu LT, Kung PT.

Risk of rapid progression to dialysis in patients with type 2 diabetes mellitus with and without diabetes-related complications at diagnosis.
Scientific Reports. 2023

Read Review
Rout P, Aslam A.

End-stage renal disease.
In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2025 Jun 22

Read Review
Sun CY, Chang SC, Wu MS.

Uremic toxins induce kidney fibrosis by activating intrarenal renin–angiotensin–aldosterone system associated epithelial-to-mesenchymal transition.
PLoS One. 2012

Read Review
Sun CY, Chang SC, Wu MS.

Uremic toxins induce kidney fibrosis by activating intrarenal renin–angiotensin–aldosterone system associated epithelial-to-mesenchymal transition.
PLoS One. 2012

Read Review
Hobby GP, Karaduta O, Dusio GF, Singh M, Zybailov BL, Arthur JM.

Chronic kidney disease and the gut microbiome
Am J Physiol Renal Physiol. 2019

Read Review

Hobby GP, Karaduta O, Dusio GF, Singh M, Zybailov BL, Arthur JM.

Chronic kidney disease and the gut microbiome
Am J Physiol Renal Physiol. 2019

Read Review
Sun CY, Chang SC, Wu MS.

Uremic toxins induce kidney fibrosis by activating intrarenal renin–angiotensin–aldosterone system associated epithelial-to-mesenchymal transition.
PLoS One. 2012

Read Review
Hobby GP, Karaduta O, Dusio GF, Singh M, Zybailov BL, Arthur JM.

Chronic kidney disease and the gut microbiome
Am J Physiol Renal Physiol. 2019

Read Review
Sun CY, Chang SC, Wu MS.

Uremic toxins induce kidney fibrosis by activating intrarenal renin–angiotensin–aldosterone system associated epithelial-to-mesenchymal transition.
PLoS One. 2012

Read Review
Hobby GP, Karaduta O, Dusio GF, Singh M, Zybailov BL, Arthur JM.

Chronic kidney disease and the gut microbiome
Am J Physiol Renal Physiol. 2019

Read Review
Hobby GP, Karaduta O, Dusio GF, Singh M, Zybailov BL, Arthur JM.

Chronic kidney disease and the gut microbiome
Am J Physiol Renal Physiol. 2019

Read Review
Sun CY, Chang SC, Wu MS.

Uremic toxins induce kidney fibrosis by activating intrarenal renin–angiotensin–aldosterone system associated epithelial-to-mesenchymal transition.
PLoS One. 2012

Read Review
Hobby GP, Karaduta O, Dusio GF, Singh M, Zybailov BL, Arthur JM.

Chronic kidney disease and the gut microbiome
Am J Physiol Renal Physiol. 2019

Read Review
Hobby GP, Karaduta O, Dusio GF, Singh M, Zybailov BL, Arthur JM.

Chronic kidney disease and the gut microbiome
Am J Physiol Renal Physiol. 2019

Read Review
Hobby GP, Karaduta O, Dusio GF, Singh M, Zybailov BL, Arthur JM.

Chronic kidney disease and the gut microbiome
Am J Physiol Renal Physiol. 2019

Read Review
Hobby GP, Karaduta O, Dusio GF, Singh M, Zybailov BL, Arthur JM.

Chronic kidney disease and the gut microbiome
Am J Physiol Renal Physiol. 2019

Read Review

Join the Roundtable
Contribute to published consensus reports, connect with top clinicians and researchers, and receive exclusive invitations to roundtable conferences.

Yes, I want to help shape the future of microbiome medicine.

Increased Info
Decreased Info
Increased
Decreased
A microbiome signature is defined through a comprehensive approach that integrates differential abundance analysis across multiple datasets, followed by interpretation for biological relevance. This rigorous process ensures that the identified signature is both statistically robust and biologically meaningful. Associations that do not meet statistical significance or lack mechanistic relevance are excluded from the final signature.
Relative Abundance (Class)
A microbiome signature is defined through a comprehensive approach that integrates differential abundance analysis across multiple datasets, followed by interpretation for biological relevance. This rigorous process ensures that the identified signature is both statistically robust and biologically meaningful. Associations that do not meet statistical significance or lack mechanistic relevance are excluded from the final signature.
Relative Abundance (Family)
A microbiome signature is defined through a comprehensive approach that integrates differential abundance analysis across multiple datasets, followed by interpretation for biological relevance. This rigorous process ensures that the identified signature is both statistically robust and biologically meaningful. Associations that do not meet statistical significance or lack mechanistic relevance are excluded from the final signature.
Relative Abundance (Genus)
A microbiome signature is defined through a comprehensive approach that integrates differential abundance analysis across multiple datasets, followed by interpretation for biological relevance. This rigorous process ensures that the identified signature is both statistically robust and biologically

meaningful. Associations that do not meet statistical significance or lack mechanistic relevance are excluded from the final signature.
Relative Abundance (Kingdom)
A microbiome signature is defined through a comprehensive approach that integrates differential abundance analysis across multiple datasets, followed by interpretation for biological relevance. This rigorous process ensures that the identified signature is both statistically robust and biologically meaningful. Associations that do not meet statistical significance or lack mechanistic relevance are excluded from the final signature.
Relative Abundance (Order)
A microbiome signature is defined through a comprehensive approach that integrates differential abundance analysis across multiple datasets, followed by interpretation for biological relevance. This rigorous process ensures that the identified signature is both statistically robust and biologically meaningful. Associations that do not meet statistical significance or lack mechanistic relevance are excluded from the final signature.
Relative Abundance (Phylum)
A microbiome signature is defined through a comprehensive approach that integrates differential abundance analysis across multiple datasets, followed by interpretation for biological relevance. This rigorous process ensures that the identified signature is both statistically robust and biologically meaningful. Associations that do not meet statistical significance or lack mechanistic relevance are excluded from the final signature.
Relative Abundance (Species)

▶ **Studies**

References

1. KDIGO 2012 Clinical Practice Guideline for the Evaluation and Management of Chronic Kidney Disease. Kidney Disease: Improving Global Outcomes (KDIGO) CKD Work Group.. (Kidney International Supplements. 2013)
2. KDIGO 2012 Clinical Practice Guideline for the Evaluation and Management of Chronic Kidney Disease. Kidney Disease: Improving Global Outcomes (KDIGO) CKD Work Group.. (Kidney International Supplements. 2013)
3. Causes of Death in End-Stage Kidney Disease: Comparison Between the United States Renal Data System and a Large Integrated Health Care System.. Bhandari SK, Zhou H, Shaw SF, Shi J, Tilluckdharry NS, Rhee CM, Jacobsen SJ, Sim JJ.. (American Journal of Nephrology. 2022)
4. Causes of Death in End-Stage Kidney Disease: Comparison Between the United States Renal Data System and a Large Integrated Health Care System.. Bhandari SK, Zhou H, Shaw SF, Shi J, Tilluckdharry NS, Rhee CM, Jacobsen SJ, Sim JJ.. (American Journal of Nephrology. 2022)
5. Causes of Death in End-Stage Kidney Disease: Comparison Between the United States Renal Data System and a Large Integrated Health Care System.. Bhandari SK, Zhou H, Shaw SF, Shi J, Tilluckdharry NS, Rhee CM, Jacobsen SJ, Sim JJ.. (American Journal of Nephrology. 2022)
6. Risk of rapid progression to dialysis in patients with type 2 diabetes mellitus with and without diabetes-related complications at diagnosis.. Shih HM, Tsai WC, Wu PY, Chiu LT, Kung PT.. (Scientific Reports. 2023)
7. End-stage renal disease.. Rout P, Aslam A.. (In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2025 Jun 22)
8. Uremic toxins induce kidney fibrosis by activating intrarenal renin–angiotensin–aldosterone system associated epithelial-to-mesenchymal transition.. Sun CY, Chang SC, Wu MS.. (PLoS One. 2012)

9. Uremic toxins induce kidney fibrosis by activating intrarenal renin–angiotensin–aldosterone system associated epithelial-to-mesenchymal transition.. Sun CY, Chang SC, Wu MS.. (PLoS One. 2012)
10. Chronic kidney disease and the gut microbiome. Hobby GP, Karaduta O, Dusio GF, Singh M, Zybailov BL, Arthur JM.. (Am J Physiol Renal Physiol. 2019)
11. Chronic kidney disease and the gut microbiome. Hobby GP, Karaduta O, Dusio GF, Singh M, Zybailov BL, Arthur JM.. (Am J Physiol Renal Physiol. 2019)
12. Uremic toxins induce kidney fibrosis by activating intrarenal renin–angiotensin–aldosterone system associated epithelial-to-mesenchymal transition.. Sun CY, Chang SC, Wu MS.. (PLoS One. 2012)
13. Chronic kidney disease and the gut microbiome. Hobby GP, Karaduta O, Dusio GF, Singh M, Zybailov BL, Arthur JM.. (Am J Physiol Renal Physiol. 2019)
14. Uremic toxins induce kidney fibrosis by activating intrarenal renin–angiotensin–aldosterone system associated epithelial-to-mesenchymal transition.. Sun CY, Chang SC, Wu MS.. (PLoS One. 2012)
15. Chronic kidney disease and the gut microbiome. Hobby GP, Karaduta O, Dusio GF, Singh M, Zybailov BL, Arthur JM.. (Am J Physiol Renal Physiol. 2019)
16. Chronic kidney disease and the gut microbiome. Hobby GP, Karaduta O, Dusio GF, Singh M, Zybailov BL, Arthur JM.. (Am J Physiol Renal Physiol. 2019)
17. Uremic toxins induce kidney fibrosis by activating intrarenal renin–angiotensin–aldosterone system associated epithelial-to-mesenchymal transition.. Sun CY, Chang SC, Wu MS.. (PLoS One. 2012)
18. Chronic kidney disease and the gut microbiome. Hobby GP, Karaduta O, Dusio GF, Singh M, Zybailov BL, Arthur JM.. (Am J Physiol Renal Physiol. 2019)
19. Chronic kidney disease and the gut microbiome. Hobby GP, Karaduta O, Dusio GF, Singh M, Zybailov BL, Arthur JM.. (Am J Physiol Renal Physiol. 2019)
20. Chronic kidney disease and the gut microbiome. Hobby GP, Karaduta O, Dusio GF, Singh M, Zybailov BL, Arthur JM.. (Am J Physiol Renal Physiol. 2019)
21. Chronic kidney disease and the gut microbiome. Hobby GP, Karaduta O, Dusio GF, Singh M, Zybailov BL, Arthur JM.. (Am J Physiol Renal Physiol. 2019)

Generated from microbiomemedicine.com/conditions/end-stage-renal-disease-esrd. © Microbiome Signatures.