Postpartum Depression (PPD)

Divine Aleru

Microbiome Signatures, Microbiome Medicine

Abstract. Postpartum depression affects many new mothers. The gut microbiome, diet, exercise, and interventions like probiotics can help manage and alleviate PPD symptoms.

Published 2025-06-10 · https://microbiomemedicine.com/conditions/postpartum-depression-ppd · A Microbiome Signatures condition monograph

Overview

Postpartum depression (PPD) is a significant mental health issue affecting 13-19% of women globally within the first year after childbirth.[1][2] It is characterized by symptoms such as persistent sadness, anxiety, fatigue, and irritability. PPD not only impacts the mother's mental health but also poses risks to infant development, including attachment issues, growth impairment, and behavioral problems.[3] Factors such as hormonal changes, a history of depression, lack of social support, and stressful life events have been shown to increase the likelihood of PPD.[4] The microbiome also plays a role in PPD, as research into the microbiome's influence on mental health has gained momentum in recent years. While the direct link between the microbiome and PPD remains under investigation, alterations in the gut-brain axis and microbial diversity are believed to influence mood and cognitive function, providing a potential avenue for non-invasive diagnostic and treatment strategies.[5]

Associated Conditions

Postpartum depression is often linked with a range of associated conditions, both psychological and physiological. Beyond the obvious co-occurrence with anxiety and obsessive-compulsive disorder, PPD has been associated with maternal suicidality, a serious consequence that underscores the importance of early detection and intervention.[6] Studies indicate that untreated PPD can also lead to longer-term emotional and behavioral problems in children, affecting their social and cognitive development.[7] In addition to mental health disorders, PPD has a strong connection with obstetric complications. Women who experience preterm birth or cesarean section deliveries are at an increased risk for developing PPD.[8] Likewise, women with gestational diabetes or those who experience complications during childbirth are more likely to report depressive symptoms postpartum.

Causes

The exact causes of postpartum depression are multifactorial and involve a complex interplay of biological, psychological, and sociocultural factors. Biological theories suggest that hormonal fluctuations after childbirth, particularly the rapid drop in estrogen and progesterone, can trigger mood disturbances.[9] Neurotransmitter imbalances, particularly in serotonin and dopamine systems, also play a crucial role in

the pathogenesis of PPD.[10] Psychological theories emphasize the stress of childbirth, particularly in the absence of social support or with a history of mental illness. Women with a prior history of depression or those experiencing stressful life events, such as financial instability or marital conflicts, are at greater risk.[11] Cultural theories suggest that postpartum depression may be viewed and experienced differently across cultures. For instance, in some non-Western cultures where there is stronger social support and family involvement, PPD prevalence is lower.[12] In contrast, cultures with less support for new mothers or where mental health stigma is prevalent, such as in some parts of Africa or Asia, show higher rates of PPD.[13] These cultural differences highlight the importance of incorporating cultural competence into the management and treatment of PPD.

Diagnosis

The diagnosis of postpartum depression typically involves clinical assessment and screening tools. The Edinburgh Postnatal Depression Scale (EPDS) is the most commonly used screening tool, with sensitivity ranging from 75% to 100% and specificity from 87% to 98%, making it an effective measure in a variety of cultural contexts.[14][15] However, cultural differences in the expression of depressive symptoms can sometimes affect the accuracy of diagnosis. In some regions, the stigma surrounding mental health issues may also prevent women from seeking treatment, which can lead to underreporting and misdiagnosis.[16] Recent research has also explored the role of microbiome signatures in diagnosing PPD.[17] Although this area is still developing, changes in gut microbiota have been implicated in mood disorders, including depression, by influencing the gut-brain axis.

Primer

Postpartum depression emerges from a complex network of physiological, psychological and environmental changes that extend far beyond hormonal fluctuations.[18] Increasingly, evidence points to the gut microbiome and trace metal dynamics as key biological layers influencing postpartum mental health.[19] Shifts in microbial composition after childbirth can disrupt immune signaling and neurotransmitter production, subtly reshaping the brain's stress and mood regulation systems.[20] At the same time, imbalances in essential metals, such as deficiencies in zinc or altered copper levels, may interfere with neurological function, amplifying vulnerability to depressive symptoms.[21] These biological changes don't occur in isolation; they interact in ways that heighten or buffer risk depending on the individual's baseline health and exposures during pregnancy and postpartum. Recognizing how these microbial and metallomic factors intersect offers a more nuanced view of PPD, not just as a psychological condition, but as one shaped by deeper biological currents. This perspective opens the door to new forms of prevention, early detection, and intervention that are grounded in biology and tailored to the postpartum experience.

Metallomic Signatures

Emerging evidence suggests that the metallomic signature of women with postpartum depression is notably disrupted, particularly involving key trace elements such as zinc, magnesium, iron, and copper. These trace metals participate in neurodevelopmental processes, neurotransmitter synthesis, immune modulation, and oxidative stress regulation, all of which are implicated in the pathophysiology of depression.[22] Several studies have demonstrated altered serum levels of these metals in individuals with depressive symptoms

during the postpartum period, implicating a distinct metallomic imbalance in PPD.[23][24] Investigating the metallomic profile of women with PPD offers potential for early diagnosis and targeted treatments.

▶ **What is the metallomic signature of postpartum depression?**

Metabolomic Signatures

The role of gut microbiota and their associated metabolites in postpartum depression is increasingly being recognized as a significant factor influencing its onset and progression. Microbial species like *Prevotellaceae* and *Veillonellaceae* are inversely associated with PPD risk, with *Prevotellaceae* known for its production of short-chain fatty acids (SCFAs) such as butyrate, which has been shown to alleviate depressive symptoms by enhancing serotonin (5-HT) levels, boosting brain-derived neurotrophic factor (BDNF), and improving blood-brain barrier integrity.[35]*Prevotellaceae* can reduce the risk of PPD by elevating Xanthine levels, a novel pathway not previously explored in PPD. This finding suggests that *Xanthine*, a metabolite linked to purine metabolism, may play a role in mitigating depressive symptoms.[36]*Ruminococcaceae UCG011* has been linked to a reduced risk of PPD, suggesting that increasing dietary fiber intake, which promotes SCFA production, could mitigate depressive behaviors.[37]*Bifidobacterium* has shown protective effects against PPD, supported by studies indicating reduced levels of *Bifidobacterium* in major depressive disorder patients.[38] This genus has been found to modulate the HPA axis and influence serotonin reuptake, further emphasizing its role in mental health.[39] Interestingly, there is a novel association between *Bifidobacterium* and LysoPI (lysophosphatidylinositol), a metabolite involved in immune and inflammatory responses, linking it to conditions such as asthma and COVID-19. The significance of LysoPI in PPD suggests that dietary interventions, like ϵ-polylysine supplementation, will offer therapeutic potential for modulating both the gut microbiota and blood metabolites.[40] Microbial metabolites in PPD pathophysiology suggests that gut microbiota-targeted interventions are promising strategies for preventing and treating postpartum depression.

Interventions

Microbiome-targeted interventions (MBTIs) offer a promising approach to managing postpartum depression by addressing both microbial imbalances and the underlying biological mechanisms of the condition. These interventions aim to restore the gut microbiota to a healthier balance, thus potentially modulating inflammation, neurotransmitter synthesis, and immune responses, all of which are implicated in PPD. By aligning therapeutic effects with clinical outcomes and key biomarkers, MBTIs can help correct the dysbiosis observed in PPD patients. This approach not only strengthens the role of microbiome signatures in diagnosis but also supports the development of effective, personalized treatments.

Intervention	Classification	Mechanism of Action	MBTI Status
Omega-3 Fatty Acids	Diet	Omega-3 fatty acids are integral to reducing inflammation and supporting neurogenesis. EPA and DHA modulate the synthesis of pro-inflammatory cytokines, improve serotonin and dopamine levels, and enhance neuroplasticity.[41][42] These actions help to reduce the neuroinflammatory pathways that contribute to postpartum depression.[43]	Promising Candidate
Probiotics	Diet	Probiotics influence the production of SCFAs, such as butyrate, which improves mood and reduces depressive symptoms.[44] Probiotics help reduce the dysregulated immune responses associated with postpartum depression.[45]	Promising Candidate
Fecal Microbiota Transplantation (FMT)	Microbiota-based therapy	FMT may potentially restore microbial diversity, modulate neuroinflammation, and influence gut-brain axis pathways. This intervention is being explored for its potential to reduce depressive-like behaviors by altering microbial composition, particularly increasing beneficial microbiota and decreasing pro-inflammatory bacteria.[46]	Experimental
Dietary Fiber	Diet	Dietary fiber promotes the production of SCFAs, like butyrate, propionate, and acetate, through microbial fermentation. Butyrate, in particular, has been shown to improve brain function by increasing brain-derived neurotrophic factor (BDNF), modulating serotonin levels, and enhancing blood-brain barrier integrity.[47]	Under Investigation
Vitamin D	Supplement	Vitamin D deficiency has been linked to increased depressive symptoms, and supplementation helps normalize the immune response, improving mood and alleviating depression in postpartum women, particularly those with low vitamin D levels.[48][49]	Experimental
Bright Light Therapy	Psychosocial	Bright light therapy helps regulate circadian rhythms by exposing individuals to bright light, which influences melatonin and serotonin levels. Light therapy may alter gut microbiota composition, indirectly influencing the gut-brain axis. [50][51]	Experimental

Intervention	Classification	Mechanism of Action	MBTI Status
Exercise and Physical Activity	Behavioral Therapy	Exercise improves mood by increasing the release of endorphins, which boost serotonin levels and help reduce inflammation.[52] Regular physical activity has been shown to positively affect gut microbiota diversity, enhancing the growth of bacteria that produce SCFAs.[53] Exercise also regulates the HPA axis and modulates neuroplasticity.[54][x]	Validated
Iron Supplementation	Supplement	Iron supplementation corrects iron deficiency anemia, which is often linked to fatigue, cognitive impairment, and depressive symptoms.[55] Iron supports neurotransmitter synthesis, particularly dopamine and serotonin, and enhances mitochondrial function.[56]	Validated
Trace elements (Zn, Cu, Mg) Supplementation	Supplement	Supplementing these metals helps restore balance in neurochemical systems disrupted in postpartum depression.[57] These metals influence gut microbiota, providing potential therapeutic benefits by regulating gut health and mood.[58]	Promising Candidate

References

1. Postpartum Depression in The Arab Region: A Systematic Literature Review. Ayoub K, Shaheen A, Hajat S.. (Clin Pract Epidemiol Ment Health, 2020; 16:)
2. A Comprehensive Review on Postpartum Depression. Suryawanshi O 4th, Pajai S.. (Cureus. 2022 Dec 20;14(12):e3274)
3. A Comprehensive Review on Postpartum Depression. Suryawanshi O 4th, Pajai S.. (Cureus. 2022 Dec 20;14(12):e3274)
4. Risk Factors of Postpartum Depression. Agrawal I, Mehendale AM, Malhotra R.. (Cureus. 2022 Oct 31;14(10):e30898)
5. The role of gut microbiota in the pathogenesis and treatment of postpartum depression.. Zhang, S., Lu, B. & Wang, G.. (Ann Gen Psychiatry 22, 36 (2023))
6. Diagnosis of peripartum depression disorder: A state-of-the-art approach from the COST Action Riseup-PPD. Radoš, S. N., Akik, B. K., Žutić, M., Rodriguez-Muñoz, M. F., Uriko, K., Motrico, E., Moreno-Peral, P., Apter, G., & Den Berg, M. L. V. (2024). (Comprehensive Psychiatry, 130, 152456)
7. A Comprehensive Review on Postpartum Depression. Suryawanshi O 4th, Pajai S.. (Cureus. 2022 Dec 20;14(12):e3274)
8. Risk Factors of Postpartum Depression. Agrawal I, Mehendale AM, Malhotra R.. (Cureus. 2022 Oct 31;14(10):e30898)
9. Risk Factors of Postpartum Depression. Agrawal I, Mehendale AM, Malhotra R.. (Cureus. 2022 Oct 31;14(10):e30898)
10. A Comprehensive Review on Postpartum Depression. Suryawanshi O 4th, Pajai S.. (Cureus. 2022 Dec 20;14(12):e3274)
11. The Rate and Risk Factors of Postpartum Depression in Vietnam From 2010 to 2020: A Literature Review. Nguyen HTT, Hoang AP, Do LTK, Schiffer S and Nguyen HTH (2021). (Front. Psychol. 12:731306)
12. The Impact of Cultural Factors Upon Postpartum Depression: A Literature Review. Bina, R. (2008). (Health Care for Women International, 29(6), 568–592.)

13. The Rate and Risk Factors of Postpartum Depression in Vietnam From 2010 to 2020: A Literature Review. Nguyen HTT, Hoang AP, Do LTK, Schiffer S and Nguyen HTH (2021). (Front. Psychol. 12:731306)
14. A Comprehensive Review on Postpartum Depression. Suryawanshi O 4th, Pajai S.. (Cureus. 2022 Dec 20;14(12):e3274)
15. The magnitude of postpartum depression among mothers in Africa: a literature review. Catherine Atuhaire et al.. (Pan African Medical Journal. 2020;37(89).)
16. Factors Related to Seeking Help for Postpartum Depression: A Secondary Analysis of New York City PRAMS Data.. Pickering, S., Ortega, M. A., Asúnsolo, Á., & Romero, D. (2019). (International Journal of Environmental Research and Public Health, 17(24), 9328)
17. The role of gut microbiota in the pathogenesis and treatment of postpartum depression.. Zhang, S., Lu, B. & Wang, G.. (Ann Gen Psychiatry 22, 36 (2023))
18. Perinatal Depression. Carlson K, Mughal S, Azhar Y, et al.. ([Updated 2025 Jan 22]. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2025 Jan-)
19. Postpartum depression and role of serum trace elements. Etebary S, Nikseresht S, Sadeghipour HR, Zarrindast MR.. (Iran J Psychiatry. 2010 Spring;5(2):40-6)
20. The role of gut microbiota in the pathogenesis and treatment of postpartum depression.. Zhang, S., Lu, B. & Wang, G.. (Ann Gen Psychiatry 22, 36 (2023))
21. Postpartum depression and role of serum trace elements. Etebary S, Nikseresht S, Sadeghipour HR, Zarrindast MR.. (Iran J Psychiatry. 2010 Spring;5(2):40-6)
22. Postpartum depression and role of serum trace elements. Etebary S, Nikseresht S, Sadeghipour HR, Zarrindast MR.. (Iran J Psychiatry. 2010 Spring;5(2):40-6)
23. Postpartum depression and role of serum trace elements. Etebary S, Nikseresht S, Sadeghipour HR, Zarrindast MR.. (Iran J Psychiatry. 2010 Spring;5(2):40-6)
24. Joint effects of traffic-related air pollution and hypertensive disorders of pregnancy on maternal postpartum depressive and anxiety symptoms. Hu, Y., Chavez, T., Eckel, S.P. et al.. (J Expo Sci Environ Epidemiol 35, 278–287 (2025))
25. Postpartum depression and role of serum trace elements. Etebary S, Nikseresht S, Sadeghipour HR, Zarrindast MR.. (Iran J Psychiatry. 2010 Spring;5(2):40-6)
26. Correlation of Serum Zinc Levels with Postpartum Depression-A Case- control Study in North Karnataka. Hiremath, Kavitha & Dharambhat, Shravya & Mutalik, Narayan & Chandrashekaraya, SH & Kashinakunti, Sangappa. (2021).. (Journal of Clinical and Diagnostic Research. 15.)
27. Correlation of Serum Zinc Levels with Postpartum Depression-A Case- control Study in North Karnataka. Hiremath, Kavitha & Dharambhat, Shravya & Mutalik, Narayan & Chandrashekaraya, SH & Kashinakunti, Sangappa. (2021).. (Journal of Clinical and Diagnostic Research. 15.)
28. Antidepressant activity of zinc and magnesium in view of the current hypotheses of antidepressant action. Szewczyk B, Poleszak E, Sowa-Kućma M, Siwek M, Dudek D, Ryszewska-Pokraśniewicz B, Radziwoń-Zaleska M, Opoka W, Czekaj J, Pilc A, Nowak G.. (Pharmacol Rep. 2008 Sep-Oct;60(5):588-9)
29. Postpartum depression and role of serum trace elements. Etebary S, Nikseresht S, Sadeghipour HR, Zarrindast MR.. (Iran J Psychiatry. 2010 Spring;5(2):40-6)
30. The association between anemia and postpartum depression: A systematic review and meta-analysis. Azami, M., Badfar, G., Khalighi, Z., Qasemi, P., Shohani, M., Soleymani, A., & Abbasalizadeh, S.. (Caspian Journal of Internal Medicine, 10(2), 115.)
31. Maternal Iron Deficiency Anemia Affects Postpartum Emotions and Cognition. Beard, J. L., Hendricks, M. K., Perez, E. M., Murray-Kolb, L. E., Berg, A., Vernon-Feagans, L., Irlam, J., Isaacs, W., Sive, A., & Tomlinson, M. (2005).. (The Journal of Nutrition, 135(2), 267-272.)
32. Gestational iron supplementation reverses depressive-like behavior in post-partum Sprague Dawley rats: Evidence from behavioral and neurohistological studies. Kukuia, K. K. E., Torbi, J., Amoateng, P., Adutwum-Ofosu, K. K., Koomson, A. E., Appiah, F., Tagoe, T. A., Mensah, J. A., Ameyaw, E. O., Adi-Dako, O., & Amponsah, S. K. (2022). (IBRO Neuroscience Reports, 12, 280-296.)
33. Postpartum depression and role of serum trace elements. Etebary S, Nikseresht S, Sadeghipour HR, Zarrindast MR.. (Iran J Psychiatry. 2010 Spring;5(2):40-6)

34. Elevated serum copper levels in women with a history of post-partum depression. Crayton, J. W., & Walsh, W. J. (2007).. (Journal of Trace Elements in Medicine and Biology, 21(1), 17-21.)

35. The microbiota-gut-brain axis and central nervous system diseases: From mechanisms of pathogenesis to therapeutic strategies. Xu, J., & Lu, Y. (2025).. (Frontiers in Microbiology, 16, 1583562)

36. The role of gut microbiota and blood metabolites in postpartum depression: A Mendelian randomization analysis. Cui, J., Zhai, Q., Yang, Z., & Liu, Y. (2024).. (Frontiers in Cellular and Infection Microbiology, 14, 1416298.)

37. High-dietary fiber intake alleviates antenatal obesity-induced postpartum depression: roles of gut microbiota and microbial metabolite short-chain fatty acid involved. Liu, Z., Li, L., Ma, S., Ye, J., Zhang, H., Li, Y., et al. (2020). (J. Agric. Food Chem. 68, 13697–13710)

38. Possible association of Bifidobacterium and Lactobacillus in the gut microbiota of patients with major depressive disorder. Aizawa, E., Tsuji, H., Asahara, T., Takahashi, T., Teraishi, T., Yoshida, S., et al. (2016).. (J. Affect. Disord. 202, 254–257. doi:)

39. The effect of exercise on depression and gut microbiota: Possible mechanisms. Yao, M., Qu, Y., Zheng, Y., & Guo, H. (2024).. (Brain Research Bulletin, 220, 111130)

40. The role of gut microbiota and blood metabolites in postpartum depression: A Mendelian randomization analysis. Cui, J., Zhai, Q., Yang, Z., & Liu, Y. (2024).. (Frontiers in Cellular and Infection Microbiology, 14, 1416298.)

41. The Role of Nutrient Supplementation to Prevent Perinatal Depression. A Narrative Review.. Aly HAI and New KJ.. (Journal of Psychiatry and Psychiatric Disorders. 7 (2023):228-247)

42. Treatment of postpartum depression: Clinical, psychological and pharmacological options. Fitelson, E., Kim, S., Baker, A. S., & Leight, K. (2010).. (International Journal of Women's Health, 3, 1.)

43. Dietary interventions for perinatal depression and anxiety: A systematic review and meta-analysis of randomized controlled trials. Tsai, Z., Shah, N., Tahir, U., Mortaji, N., Owais, S., Perreault, M., & Van Lieshout, R. J. (2023). (The American Journal of Clinical Nutrition, 117(6), 1130-1142.)

44. Effect of Lactobacillus rhamnosus HN001 in Pregnancy on Postpartum Symptoms of Depression and Anxiety: A Randomised Double-blind Placebo-controlled Trial. Slykerman, R., Hood, F., Wickens, K., Thompson, J., Barthow, C., Murphy, R., Kang, J., Rowden, J., Stone, P., Crane, J., Stanley, T., Abels, P., Purdie, G., Maude, R., Mitchell, E., & Group, P. S. (2017).. (EBioMedicine, 24, 159.)

45. The role of gut microbiota in the pathogenesis and treatment of postpartum depression. Zhang, S., Lu, B. & Wang, G.. (Ann Gen Psychiatry 22, 36 (2023).)

46. Gut microbiota dysbiosis contributes to depression-like behaviors via hippocampal NLRP3-mediated neuroinflammation in a postpartum depression mouse model. Xu, Q., Sun, L., Chen, Q., Jiao, C., Wang, Y., Li, H., Xie, J., Zhu, F., Wang, J., Zhang, W., Xie, L., Wu, H., Zuo, Z., & Chen, X. (2024). (Brain, Behavior, and Immunity, 119, 220-235.)

47. Gut microbiota: Linking nutrition and perinatal depression. Song, J., Zhou, B., Kan, J., Liu, G., Zhang, S., Si, L., Zhang, X., Yang, X., Ma, J., Cheng, J., Liu, X., & Yang, Y. (2022).. (Frontiers in Cellular and Infection Microbiology, 12, 932309.)

48. Gut microbiota: Linking nutrition and perinatal depression. Song, J., Zhou, B., Kan, J., Liu, G., Zhang, S., Si, L., Zhang, X., Yang, X., Ma, J., Cheng, J., Liu, X., & Yang, Y. (2022).. (Frontiers in Cellular and Infection Microbiology, 12, 932309.)

49. The role of gut microbiota in the pathogenesis and treatment of postpartum depression. Zhang, S., Lu, B. & Wang, G.. (Ann Gen Psychiatry 22, 36 (2023).)

50. "Photobiomics": Can Light, Including Photobiomodulation, Alter the Microbiome?. Liebert, A., Bicknell, B., Johnstone, D. M., Gordon, L. C., Kiat, H., & Hamblin, M. R. (2019).. (Photobiomodulation, Photomedicine, and Laser Surgery, 37(11), 681.)

51. Treatment of postpartum depression: Clinical, psychological and pharmacological options. Fitelson, E., Kim, S., Baker, A. S., & Leight, K. (2010).. (International Journal of Women's Health, 3, 1.)

52. Treatment of postpartum depression: Clinical, psychological and pharmacological options. Fitelson, E., Kim, S., Baker, A. S., & Leight, K. (2010).. (International Journal of Women's Health, 3, 1.)

53. Physical Exercise and the Gut Microbiome: A Bidirectional Relationship Influencing Health and Performance. Varghese, S., Rao, S., Khattak, A., Zamir, F., & Chaari, A. (2024).. (Nutrients, 16(21), 3663.)

54. Exercise and the hypothalamic–pituitary–adrenal axis: A special focus on acute cortisol and growth hormone responses. Anderson, T., Berry, N. T., & Wideman, L. (2019). (Current Opinion in Endocrine and Metabolic Research, 9, 74-77.)

55. Trace Elements Levels in Major Depressive Disorder—Evaluation of Potential Threats and Possible Therapeutic Approaches. Baj, J., Bargieł, J., Cabaj, J., Skierkowski, B., Hunek, G., Portincasa, P., Flieger, J., & Smoleń, A. (2023).. (International Journal of Molecular Sciences, 24(20), 15071.)

56. Iron and Mechanisms of Emotional Behavior. Kim, J., & Wessling-Resnick, M. (2014).. (The Journal of Nutritional Biochemistry, 25(11), 1101.)

57. Toxic and essential metals: Metabolic interactions with the gut microbiota and health implications. Zhu, Q., Chen, B., Zhang, F., Zhang, B., Guo, Y., Pang, M., Huang, L., & Wang, T. (2024).. (Frontiers in Nutrition, 11, 1448388.)

58. Toxic and essential metals: Metabolic interactions with the gut microbiota and health implications. Zhu, Q., Chen, B., Zhang, F., Zhang, B., Guo, Y., Pang, M., Huang, L., & Wang, T. (2024).. (Frontiers in Nutrition, 11, 1448388.)

Generated from microbiomemedicine.com/conditions/postpartum-depression-ppd. © Microbiome Signatures.