

# Contents

<b>1</b>	<b>Analysis</b>	<b>3</b>
1.1	Introduction . . . . .	3



# Chapter 1

## Analysis

blabla

### 1.1 Introduction

Discovering the mechanism responsible for electroweak symmetry-breaking and the origin of mass for elementary particles has been one of the major goals of the physics program at the Large Hadron Collider (LHC) [1]. In the Standard Model (SM) this mechanism requires the existence of a single scalar particle, the Higgs boson [2, 3, 4, 5, 6]. In the Minimal Supersymmetric extension of the Standard Model (MSSM) [7, 8] the Higgs sector is composed of two Higgs doublets of opposite hyper-charge, resulting in five observable Higgs bosons. Two of these Higgs bosons are neutral and  $CP$ -even ( $h, H$ ), one is neutral and  $CP$ -odd ( $A$ ) and two are charged ( $H^\pm$ ). At tree level their properties such as masses, widths and branching ratios can be predicted in terms of only two parameters, often chosen to be the mass of the  $CP$ -odd Higgs boson  $m_A$ , and the ratio of the vacuum expectation values of the two Higgs doublets  $\tan\beta$ . For relatively large values of  $\tan\beta$  one of the  $CP$ -even Higgs bosons is almost degenerate in mass with  $A$ . Moreover, Higgs couplings to down (up) type fermions are enhanced (suppressed) by  $\tan\beta$ , meaning that for large  $\tan\beta$  bottom-quark and  $\tau$  lepton will play a more important role than in the SM case either for production and decay.

The production of the neutral  $CP$ -even MSSM Higgs bosons at hadron colliders proceeds via the same processes as for the SM Higgs production. However, the pseudoscalar  $A$  instead cannot be produced in association with gauge bosons or in vector boson fusion (VBF) at tree-level, as this coupling is forbidden due to  $CP$ -invariance. At the LHC one of the most relevant production mechanisms for the MSSM Higgs bosons is gluon-gluon fusion,  $gg \rightarrow A/H/h$ . In addition, the production in association with  $b$ -quarks becomes important for large value of  $\tan\beta$ . The decays of the neutral MSSM Higgs bosons (in the assumption that all supersymmetric particle

are heavy enough) are the same as for the SM one with the already cited exception of  $A$ , however the decay rates depend on a large extent to the couplings with fermions and gauge bosons.

Searches for neutral MSSM Higgs bosons have been performed at LEP [9], the Tevatron [10, 11, 12, 13, 14, 15] and the LHC [16, 17]. In this note a search for neutral MSSM Higgs bosons with the ATLAS experiment at CERN is presented, using proton-proton collisions at centre-of-mass energy of 8 TeV, with a recorded integrated luminosity of  $20.3 fb^{-1}$ . The results of this search are interpreted in a model independent fashion, as limits on the product of the cross section and branching ratio for such a new particle, as well as limits on the MSSM in the  $m_h^{max}$  scenario [34]. Only  $b$ -quark associated and gluon fusion are considered as production mode for the Higgs bosons, the search then focuses on the subsequent decay into a  $\tau^+\tau^-$  pair. Furthermore, only cases in which both  $\tau$  decays leptonically, with one decaying to an electron and the other to a muon, are considered. This final state corresponds to a total  $\tau^+\tau^-$  branching ratio of approximately 6%. The analysis strategy is to split the selected events in two categories by requesting the presence ( $b$ -tag) or absence ( $b$ -veto) of a jet coming from a  $b$ -quark. This solution helps to separate the contribution of the two production modes and allows for optimisation of selections due to the different backgrounds for the different final states..

The signal topology is characterised by a final state with an electron, a muon, and missing transverse energy due to the presence of four neutrinos from the  $\tau$  decays. Furthermore, the final state may be split by the presence or absence of a  $b$ -quark initiated jet, depending on the production process. The background processes which are considered in this study are the production of  $W$  and  $Z$  bosons in association with jets ( $W/Z$ +jets), pairs of top quarks ( $t\bar{t}$ ), single top quark (the so-called single-top) and pairs of electroweak gauge bosons ( $WW, WZ, ZZ$ ). Finally QCD multi-jet also forms a non-negligible background due to its large production cross-section. Where possible these backgrounds are estimated using data driven methods.

This note is structured as follows: The ATLAS detector is briefly described in Section ???. In Section ??? the collision data set, the Monte Carlo-simulated event samples as well as hybrid (tau-embedded) data samples used in this study are described. The reconstruction of physics objects, the trigger requirements and the offline event selection are discussed in Sections ??? and ???. Background estimation methods are described in Section ???, and the systematic uncertainties are discussed in Section ???. A statistical analysis and resulting exclusion limits are described in Section ???, followed by conclusions in Section ???.

# Bibliography

- [1] L. Evans and P. Bryant, *LHC Machine*, JINST **3** (2008) S08001.
- [2] F. Englert and R. Brout, *Broken Symmetry and the Mass of Gauge Vector Mesons*, Phys. Rev. Lett. **13** (1964) 321.
- [3] P. W. Higgs, *Broken symmetries, massless particles and gauge fields*, Phys. Lett. **12** (1964) 132.
- [4] P. W. Higgs, *Broken Symmetries and the Masses of Gauge Bosons*, Phys. Rev. Lett. **13** (1964) 508.
- [5] P. W. Higgs, *Spontaneous Symmetry Breakdown without Massless Bosons*, Phys. Rev. **145** (1966) 1156.
- [6] G. S. Guralnik, C.R. Hagen and T. W. B. Kibble Phys.Rev.Lett. **13** (1964) 585.
- [7] N. P. Nilles, *Supersymmetry, supergravity and particle physics*, Phys. Rep. **110** (1984) 1.
- [8] H. E. Haber and G. L. Kane, *The search for supersymmetry: Probing physics beyond the standard model*, Phys. Rep. **117** (1985) 75.
- [9] ALEPH, DELPHI, L3 and OPAL Collaboration, *Search for neutral MSSM Higgs bosons at LEP*, Eur. Phys. J. **C47** (2006) 547.
- [10] *Combined CDF and D0 upper limits on MSSM Higgs boson production in tau-tau final states with up to  $2.2\text{ fb}^{-1}$  of data*, arXiv:1003.3363 [hep-ex].
- [11] CDF Collaboration, T. Aaltonen et al. Phys. Rev. Lett. **103** (2009) 201801.
- [12] D0 Collaboration, V. Abazov et al. Phys. Rev. Lett. **101** (2008) 071804.
- [13] TNPWG (Tevatron New Physics Higgs Working Group), CDF and D0 Collaborations, *Search for Neutral Higgs Bosons in Events with Multiple Bottom Quarks at the Tevatron*, arXiv:1207.2757 [hep-ex].

- [14] CDF Collaboration, T. Aaltonen et al., *Search for Higgs Bosons Produced in Association with  $b$ -quarks*, Phys.Rev. **D85** (2012) 032005, [arXiv:1106.4782 \[hep-ex\]](#).
- [15] D0 Collaboration, V.M. Abazov et al., *Search for neutral Higgs bosons in the multi- $b$ -jet topology in  $5.2\text{fb}^{-1}$  of  $p\bar{p}$  collisions at  $\sqrt{s} = 1.96\text{ TeV}$* , Phys.Lett. **B698** (2011) 97–104, [arXiv:1011.1931 \[hep-ex\]](#).
- [16] The CMS Collaboration, S. Chatrchyan et al., [arXiv:1104.1619 \[hep-ex\]](#) [hep-ex].
- [17] The ATLAS Collaboration, *Search for the neutral Higgs bosons of the Minimal Supersymmetric Standard Model in  $pp$  collisions at  $\sqrt{s} = 7\text{ TeV}$  with the ATLAS detector*, [arXiv:1211.6956 \[hep-ex\]](#).
- [18] T. A. Collaboration, *Observation of a new particle in the search for the Standard Model Higgs boson with the ATLAS detector at the LHC*, Physics Letters B **716** (2012) 1–29.
- [19] T. C. Collatoration, *Observation of a new boson at a mass of  $125\text{ GeV}$  with the CMS experiment at the LHC*, Physics Letters B **716** (2012) 30–61.
- [20] S. Heinemeyer, O. Stål and G. Weiglein, *Interpreting the LHC Higgs search results in the MSSM*, Phys.Lett. **B710** (2012) 201–206, [arXiv:1112.3026 \[hep-ph\]](#).
- [21] A. Arbey, M. Battaglia, A. Djouadi and F. Mahmoudi, *The Higgs sector of the phenomenological MSSM in the light of the Higgs boson discovery*, JHEP **1209** (2012) 107, [arXiv:1207.1348 \[hep-ph\]](#).
- [22] The ATLAS Collaboration, G. Aad et al., *The ATLAS Experiment at the CERN Large Hadron Collider*, JINST **3** (2008) S08003.
- [23] M. L. Mangano et al., *ALPGEN, a generator for hard multiparton processes in hadronic collisions*, JHEP **07** (2003) 001.
- [24] J. Alwall et al., *Comparative study of various algorithms for the merging of parton showers and matrix elements in hadronic collisions*, Eur. Phys. J. **C53** (2008) 473, [arXiv:0706.2569](#).
- [25] S. Frixione and B. R. Webber, *Matching NLO QCD computations and parton shower simulations*, JHEP **06** (2002) 029, [hep-ph/0204244](#).
- [26] B. P. Kersevan and E. Richter-Was, *The Monte Carlo Event Generator AcerMC 2.0 with Interfaces to PYTHIA 6.2 and HERWIG 6.5*, [arXiv:0405247v1 \[hep-ph\]](#).

- [27] G. Corcella et al., *HERWIG 6: an event generator for hadron emission reactions with interfering gluons (including supersymmetric processes)*, JHEP **01** (2001) 010.
- [28] J. M. Butterworth, J. R. Forshaw, and M. H. Seymour, *Multiparton Interactions in Photoproduction at HERA*, Z. Phys. **C72** (1996) 637.
- [29] T. Binoth, M. Ciccolini, N. Kauer, and M. Kramer, *Gluon-induced W-boson pair production at the LHC*, JHEP **12** (2006) 046.
- [30] A. S. et al., *Higgs boson production in gluon fusion*, JHEP **02** (2009) 029.
- [31] T. Gleisberg et al., *Event generation with SHERPA 1.1*, JHEP **02** (2009) 007.
- [32] J. Pumplin, D. R. Stump, J. Huston, H. L. Lai, P. M. Nadolsky and W. K. Tung, “New generation of parton distributions with uncertainties from global QCD analysis,” JHEP **0207** (2002) 012 [hep-ph/0201195].
- [33] H. -L. Lai, M. Guzzi, J. Huston, Z. Li, P. M. Nadolsky, J. Pumplin and C. -P. Yuan, “New parton distributions for collider physics,” Phys. Rev. D **82** (2010) 074024 [arXiv:1007.2241 [hep-ph]].
- [34] M. Carena, S. Heinemeyer, C. E. M. Wagner, and G. Weiglein, *Suggestions for benchmark scenarios for MSSM Higgs boson searches at hadron colliders*, Eur. Phys. J. **C26** (2003) 601–607, hep-ph/0202167.
- [35] The ATLAS Collaboration, *ATLAS Monte Carlo Tunes for MC09*, ATL-PHYS-PUB-2010-002.
- [36] S. Jadach, J. H. Kuhn and Z. Was, *TAUOLA - a library of Monte Carlo programs to simulate decays of polarized  $\tau$  leptons*, Comput. Phys. Commun. **64** (1990) 275.
- [37] E. Barberio, B. V. Eijk and Z. Was, *Photos - a universal Monte Carlo for QED radiative corrections in decays*, Comput. Phys. Commun. **66** (1991) 115.
- [38] The GEANT4 Collaboration, S. Agostinelli et al., *GEANT4 - a simulation toolkit*, Nucl. Instrum. Meth. **A506** (2003) 250.
- [39] The ATLAS Collaboration, G. Aad et al., *The ATLAS Simulation Infrastructure*, ATLAS-SOFT-2010-01-004, submitted to Eur. Phys. J. C., arXiv:1005.4568.
- [40] The ATLAS Collaboration, *Estimation of  $Z \rightarrow \tau\tau$  Background in VBF  $H \rightarrow \tau\tau$  Searches from  $Z \rightarrow \mu\mu$  Data using an Embedding Technique*, ATL-PHYS-INT-2009-109.

- [41] The ATLAS Collaboration, *Search for the Standard Model Higgs boson in the  $H \rightarrow \tau\tau$  decay mode with 4.7 fb of ATLAS detector*, Tech. Rep. ATLAS-CONF-2012-014, CERN, Geneva, Mar, 2012.
- [42] The ATLAS Collaboration, *Search for the Standard Model Higgs boson  $H \rightarrow \tau\tau$  decays with the ATLAS detector*, ATL-COM-PHYS-2013-722.
- [43] T. S. et al., *Z physics at LEP 1*, CERN 89-08 **3** (1989) 143.
- [44] The ATLAS Collaboration, *Expected Performance of the ATLAS Experiment - Detector, Trigger and Physics*, CERN-OPEN-2008-020, [arXiv:0901.0512](https://arxiv.org/abs/0901.0512).
- [45] The ATLAS Collaboration, *ATLAS Muon Momentum Resolution in the First Pass Reconstruction of the 2010 p-p Collision Data at  $\sqrt{s} = 7$  TeV*, ATLAS-CONF-2011-046.
- [46] The ATLAS Collaboration, *Muon reconstruction efficiency in reprocessed 2010 LHC p-p collision data recorded with the ATLAS detector*, ATLAS-CONF-2011-063.
- [47] The ATLAS Collaboration, *Expected electron performance in the ATLAS experiment*, ATLAS-PUB-2011-006.
- [48] ATLAS egamma WG, *Electron efficiency measurements*, <https://twiki.cern.ch/twiki/bin/view/AtlasProtected/EfficiencyMeasurements>.
- [49] M. Cacciari, G. P. Salam, and G. Soyez, *The anti- $k_t$  jet clustering algorithm*, JHEP **04** (2008) 063.
- [50] W. Lampl et al., *Calorimeter Clustering Algorithms : Description and Performance*, ATL-LARG-PUB-2008-002.
- [51] T. Barillari et al., *Local Hadron Calibration*, ATL-LARG-PUB-2009-001.
- [52] The ATLAS Collaboration, *Jet energy scale and its systematic uncertainty in proton-proton collisions at  $\sqrt{s} = 7$  TeV in ATLAS 2010 data*, ATLAS-CONF-2011-032.
- [53] The ATLAS Collaboration, *Performance of the Reconstruction and Identification of Hadronic tau Decays in ATLAS with 2011 Data*, ATLAS-CONF-2012-142.
- [54] The ATLAS Collaboration, *Reconstruction and Calibration of Missing Transverse Energy and Performance in Z and W events in ATLAS Proton-Proton Collisions at  $\sqrt{s}=7$  TeV*, ATLAS-CONF-2011-080.



- [55] A. Elagin, P. Murat, A. Pranko, and A. Safonov, *A New Mass Reconstruction Technique for Resonances Decaying to di-tau*, [arXiv:1012.4686 \[hep-ex\]](#). \* Temporary entry \*.
- [56] ATLAS Jet/EtMiss Combined Performance  
Group, Jet Energy Resolution Provider,  
<https://twiki.cern.ch/twiki/bin/view/Main/JetEnergyResolutionProvider>.
- [57] The ATLAS Collaboration, *Data-Quality Requirements and Event Cleaning for Jets and Missing Transverse Energy Reconstruction with the ATLAS Detector in Proton-Proton Collisions at a Center-of-Mass Energy of  $\sqrt{s} = 7$  TeV*, ATLAS-CONF-2010-038.
- [58] T. A. Collaboration, *Search for neutral MSSM Higgs bosons decaying to  $\tau\tau$  pairs in proton-proton collisions at with the ATLAS detector*, Physics Letters B **705** (2011) no. 3, 174 – 192.
- [59] The ATLAS Collaboration, *Data-driven estimation of the background to charged Higgs boson searches using hadronically-decaying tau final states in ATLAS*, ATLAS-CONF-2011-051.
- [60] The ATLAS Collaboration, *Measurement of the  $Z \rightarrow \tau\tau$  cross section with the ATLAS detector*, Phys. Rev. D **84** (2011) 112006.
- [61] T. A. Collaboration, *Search for the neutral Higgs bosons of the Minimal Supersymmetric Standard Model in pp collisions at  $\sqrt{s} = 7$  TeV with the ATLAS detector*, JHEP , [arXiv:1211.6956](#).
- [62] Atlas statistics forum, *ABCD method in searches*, [link](#)
- [63] The ATLAS Collaboration, *Search for Neutral MSSM Higgs Bosons  $H$  to  $\tau\tau$  to  $l\tau_h$  with the ATLAS Detector in 7 TeV Collisions*, ATL-COM-PHYS-2012-094.
- [64] The ATLAS Collaboration, *Search for neutral Higgs Bosons in the decay mode  $H \rightarrow \tau\tau \rightarrow ll + 4\nu$  in proton proton collision at  $\sqrt{7}$  TeV with the ATLAS Detector*, ATL-COM-PHYS-2011-758.
- [65] The ATLAS Collaboration, *Measurement of the b-tag Efficiency in a Sample of Jets Containing Muons with 5  $fb^{-1}$  of Data from the ATLAS Detector*, ATLAS-CONF-2012-043.
- [66] The ATLAS Collaboration, *Luminosity Determination in pp Collisions at  $\sqrt{s} = 7$  TeV using the ATLAS Detector in 2011*, ATLAS-CONF-2011-116.
- [67] T. Sjostrand, S. Mrenna and P. Skands, *PYTHIA 6.4 physics and manual*, JHEP **05** (2006) 026.

- [68] A. B. et al., *Rivet user manual*, [arXiv:1003.0694](#) [hep-ph].
- [69] E. G. G. Cowan, K. Cranmer and O. Vitells, *Asymptotic formulae for likelihood-based tests of new physics*, [arXiv:1007.1727](#) [hep-ex].
- [70] LHC Higgs Cross Section Working Group, S. Dittmaier, C. Mariotti, G. Passarino, R. Tanaka (Eds.), et al., *Handbook of LHC Higgs Cross Sections: 1. Inclusive Observables*, [arXiv:1101.0593](#) [hep-ph].
- [71] LHC Higgs Cross Section Working Group, S. Dittmaier, C. Mariotti, G. Passarino, and R. Tanaka (Eds.), *Handbook of LHC Higgs Cross Sections: 2. Differential Distributions*, CERN-2012-002 (CERN, Geneva, 2012) , [arXiv:1201.3084](#) [hep-ph].
- [72] D. de Florian, G. Ferrera, M. Grazzini and D. Tommasini, *Transverse-momentum resummation: Higgs boson production at the Tevatron and the LHC*, JHEP **1111** (2011) , [arXiv:1109.2109](#) [hep-ph].
- [73] Statistical twiki, NuisanceCheck. <https://twiki.cern.ch/twiki/bin/view/AtlasProtected/Nu>