

# Contents

<b>1</b>	<b>The Higgs Bosons and The MSSM</b>	<b>3</b>
1.1	The Standard Model of Particle Physics and its Limitations . . . .	4
1.2	The Minimal Supersymmetric Standard Model . . . . .	5
1.3	The Higgs Sector in The MSSM . . . . .	7
1.4	Higgs Phenomenology in The MSSM . . . . .	7



# Chapter 1

## The Higgs Bosons and The MSSM

This chapter is devoted to introduce the the Minimal Supersymmetric extension of the Standard Model (MSSM) whit focus on its Higgs sector. In section... introduction is given based on [3],

## 1.1 The Standard Model of Particle Physics and its Limitations

The Standard Model (SM) of particle physics is a theory aimed to describe and quantitatively predict the phenomenology of fundamental interaction. At “microscopic” level all the phenomenology of matter and radiation can be understood in terms of three classes of fundamental interactions, which are the strong, the electromagnetic and the weak interactions, the gravitational force is instead negligible in atomic and nuclear physics (quantum effects in gravity are expected to become important at energies corresponding to the Planck mass  $E \sim M_{\text{planck}}c^2 \sim 10^{19}$  GeV). All these interactions (with the exception of gravity) are described by a local relativistic quantum field theory. To each particle, described as pointlike, is associated a field with suitable (depending on the particle spin) transformation properties under the Lorentz group (the relativistic space-time coordinate transformations). The description of all these particle interactions is based on a common principle: “gauge” invariance. A gauge symmetry is invariance under transformations that rotate the basic internal degrees of freedom but with rotation angles that depend on the space-time point.

The SM is a gauge field theory based on the symmetry group  $SU(3)_c \otimes SU(2)_L \otimes U(1)_Y$ . This group has  $8 + 3 + 1 = 12$  generators with a non trivial commutator algebra. The electromagnetic and weak interactions (EW) are described by the  $SU(2)_L \otimes U(1)_Y$  symmetry group, while the  $SU(3)_c$  is the colour group of the theory of strong interactions (QCD). To each generator of the symmetry group is associated a vector boson which act as mediator of the corresponding interactions. Eight gluons are associated to the  $SU(3)_c$  colour generators, while for  $SU(2)_L \otimes U(1)_Y$  there are four gauge bosons  $W^\pm$ ,  $Z^0$  and  $\gamma$ . Of these, only the gluons and the photon are massless since the symmetry induced by the other three generators is spontaneously broken. In the SM the spontaneous symmetry breaking is realized by the Higgs mechanism, the Higgs particle has been recently found at the LHC with  $m_H \sim 125$  GeV representing one of the major milestones of particle physics. The Higgs boson acts as mediator of a new class of interactions that, at tree level, are coupled in proportion to the particle masses.

The fermionic (all of spin 1/2) matter fields of the SM are quarks and leptons. Quarks are subject to all SM interactions, each type of quark is a colour triplet and carries electroweak charges, in particular electric charges  $+2/3$  for up-type quarks and  $-1/3$  for down-type quarks. Leptons are colourless but have electroweak charges, in particular electric charges  $-1$  for charged leptons  $e$ ,  $\mu$  and  $\tau$  (opposite sign charge is intended for respective anti-particle) and charge 0 for neutrinos  $\nu_e$ ,  $\nu_\mu$  and  $\nu_\tau$ . Quarks and leptons are grouped in three “generations” with equal quantum numbers but different masses.

Precision tests of the SM have been performed over a wide range of energies in experiment in the last several decades, Precision tests [8] of the standard electroweak theory performed at LEP, SLC and Tevatron [7], has confirmed the couplings of quark and leptons to the weak gauge bosons  $W^\pm$  and  $Z$  are indeed precisely those prescribed by the gauge symmetry. The accuracy of a few per-mille for these tests implies that, not only the tree level, but also the structure of quantum corrections

has been verified. Several other experimental results [6] rare decays and the anomalous magnetic moment of the muon provide a test for low-energies of the standard model.

## 1.2 The Minimal Supersymmetric Standard Model

Supersymmetry (SUSY) was first introduced since it offer a natural way to solve the hierarchy problem....[] This is achieved by introducing a symmetry transformation that relates bosons with fermions. The SUSY generators  $\mathcal{Q}$  transforms fermion into bosons and vice versa:

$$\mathcal{Q}|\text{Fermion}\rangle = |\text{Boson}\rangle, \quad \mathcal{Q}|\text{Boson}\rangle = |\text{Fermion}\rangle \quad (1.1)$$

This is suggesting that in a supersymmetric extension of the SM [?] each of the known fundamental particles is in either a chiral or gauge “supermultiplet” and must have a superpartner with spin differing by  $1/2$  unit. Fermions since the left-handed and right-handed component transform differently under gauge transformations also their superpartner should maintain this property, the name of the superpartner of the quarks and leptons are made by adding an “s” to the SM name, standing for scalar. Accordingly, the gauge bosons related to the generator of the group  $SU(3)_c \otimes SU(2)_L \otimes U(1)_Y$  should also have a spin  $1/2$  partner, whose name will be made by adding a “ino” at the end of the SM name. The symbol of superpartners is defined by adding a  $(\tilde{\phantom{x}})$  to the SM symbol.

The Minimal Supersymmetric extension of the Standard Model (MSSM) [?], is defined by requiring the minimal gauge group (i.e., the SM one) and the minimal particle content: three generation of fermions (without right-handed neutrinos), gauge bosons and two Higgs doublet, each with its superpartners. Tables 1.1 and 1.2 summarize chiral and gauge supermultiplets in the MSSM. This guys will mix to form neutralino and chargino....

### ***R*-parity conservation**

The MSSM also requires a discrete and multiplicative symmetry called *R*-parity [?], this symmetry assures baryon and lepton number conservation and it is defined as follows:

$$R_p = (-1)^{2s+3B=L} \quad (1.2)$$

where  $L$  and  $B$  are lepton and baryon numbers and  $s$  stands for the spin quantum number. *R*-parity quantum numbers has value  $+1$  for ordinary SM particles and  $-1$  for their superpartners. This symmetry was first introduced to overcome the problem of instability of the proton, lepton and baryon number violation leads, in many cases, to unstable proton with life-time shorter than the experimental lower limit. However the conservation of *R*-parity has also other important phenomenological consequences, SUSY particle are always produced in pairs and in their decay there is always an odd number of SUSY particle, and the lightest SUSY particle is stable, providing a good candidate for dark matter.

Names	Supermultiplets	Spin 1/2	Spin 0
quark, squarks ( $\times 3$ families)	$Q$	$(u_L \ d_L)$	$(\tilde{u}_L \ \tilde{d}_L)$
	$\bar{u}$	$u_R^\dagger$	$\tilde{u}_R^*$
	$\bar{d}$	$d_R^\dagger$	$\tilde{d}_R^*$
leptons, sleptons ( $\times 3$ families)	$L$	$(\nu \ e_L)$	$(\tilde{\nu} \ \tilde{e}_L)$
	$\bar{e}$	$e_R^\dagger$	$\tilde{e}_R^*$
higgsinos, Higgs	$H_1$	$(\tilde{H}_1^0 \ \tilde{H}_1^-)$	$(H_1^0 \ H_1^-)$
	$H_1$	$(\tilde{H}_2^+ \ \tilde{H}_2^0)$	$(H_2^+ \ H_2^0)$

Table 1.1: This table is based on [4] and summarize the chiral supermultiplets in the Minimal Supersymmetric Standard Model. The spin-0 fields are complex scalars and the spin-1/2 are left-handed two-component Weyl fermions.

Names	Spin 1	Spin 1/2
gluon, gluino	$g$	$\tilde{g}$
W bosons, winos	$W^\pm \ W^0$	$\tilde{W}^\pm \ \tilde{W}^0$
B boson, bino	$B^0$	$\tilde{B}^0$

Table 1.2: This table is based on [4] and summarize the gauge supermultiplets in the Minimal Supersymmetric Standard Model.

### The Soft SUSY Breaking

In case the Supersymmetry is an exact symmetry of nature, the bosonic fields and the corresponding fermion fields should have the same mass and quantum numbers, except for the spin. However, the particle spectrum of SUSY has not yet been observed, suggesting that these particles should have a higher mass than their SM superpartners. To achieve SUSY-breaking in a way which does not reintroduce the quadratic divergences to the Higgs mass squared, a so called “soft-SUSY-breaking” term is introduced [?], this term explicitly breaks SUSY introducing ad hoc the mass terms for Higgs, gauginos and sfermions, furthermore trilinear coupling terms between sfermions and Higgs bosons are introduced. In general, if intergenerational mixing and complex phases are allowed, the soft-SUSY-breaking terms will introduce a huge number of unknown parameters  $\mathcal{O}(100)$  [?]. However, in absence of phases and mixing, and if obey to a set of boundary conditions [?], only few new parameters are introduced.

## 1.3 The Higgs Sector in The MSSM

In the MSSM two doublets of complex scalar field of opposite hypercharge are required to break the electroweak symmetry, this requirement is motivated by the needs to generate masses separately to isospin up-type fermion and down-type fermions [?] and to cancel chiral anomalies that otherwise would spoil the renormalizability of the theory [?]. The two Higgs doublets then are:

$$H_1 = \begin{pmatrix} H_1^0 \\ H_1^- \end{pmatrix} \text{ with } Y_{H_1} = -1, \quad H_2 = \begin{pmatrix} H_2^+ \\ H_2^0 \end{pmatrix} \text{ with } Y_{H_2} = +1 \quad (1.3)$$

In analogy with the SM, a similar Higgs mechanism is employed in the MSSM [?] to break electroweak symmetry. The neutral components of the two Higgs fields acquire vacuum expectation values:

$$\langle H_1^0 \rangle = \frac{v_1}{\sqrt{2}}, \quad \langle H_2^0 \rangle = \frac{v_2}{\sqrt{2}} \quad (1.4)$$

Three of the original eight degrees of freedom of the scalar fields are absorbed by the  $W^\pm$  and  $Z$  bosons, building their longitudinal polarizations and acquire masses. The remaining degrees of freedom correspond to five scalar Higgs bosons: two CP-even and neutral  $h$  and  $H$ , a neutral pseudoscalar boson  $A$  and a pair of charged bosons  $H^\pm$ . At tree level, besides the masses of these particles, two additional parameters define the system: the mixing angle in the neutral CP-even sector  $\alpha$  and the ratio between the two vacuum expectation values  $\tan \beta = v_1/v_2$ .

## 1.4 Higgs Phenomenology in The MSSM





# Bibliography

- [1] W. Hollik, “*Electroweak theory*,” hep-ph/9602380.
- [2] Michael E. Peskin, Dan V. Schroeder, *An Introduction To Quantum Field Theory*, Westview Press, 1995.
- [3] G. Altarelli, “*Collider Physics within the Standard Model: a Primer*,” arXiv:1303.2842 .
- [4] S. P. Martin, “*A Supersymmetry primer*,” In \*Kane, G.L. (ed.): Perspectives on supersymmetry II\* 1-153 [hep-ph/9709356].
- [5] A. Djouadi, *The Anatomy of Electroweak Symmetry Breaking Tome II: The Higgs Bosons in the Minimal Supersymmetric Model*, Phys. Rep. 459 (2008) 1.
- [6] The Review of Particle Physics, J. Beringer et al. (Particle Data Group), Phys. Rev. D86, 010001 (2012) and 2013 partial update for the 2014 edition.
- [7] The ALEPH, CDF, D0, DELPHI, L3, OPAL and SLD Collaborations, the LEP Electroweak Working Group, Tevatron Working Group and SLD Electroweak and Heavy Flavour Working Groups, *Precision Electroweak Measurements and Constraints on the Standard Model*, arXiv:1012.2367. Updated for 2012 winter conferences, March 2012.
- [8] G. Altarelli and M. W. Grunewald, “*Precision electroweak tests of the standard model*,” Phys. Rept. **403-404** (2004) 189 [hep-ph/0404165].
- [9] L. Evans and P. Bryant, *LHC Machine*, JINST **3** (2008) S08001.
- [10] F. Englert and R. Brout, *Broken Symmetry and the Mass of Gauge Vector Mesons*, Phys. Rev. Lett. **13** (1964) 321.
- [11] P. W. Higgs, *Broken symmetries, massless particles and gauge fields*, Phys. Lett. **12** (1964) 132.
- [12] P. W. Higgs, *Broken Symmetries and the Masses of Gauge Bosons*, Phys. Rev. Lett. **13** (1964) 508.
- [13] P. W. Higgs, *Spontaneous Symmetry Breakdown without Massless Bosons*, Phys. Rev. **145** (1966) 1156.
- [14] G. S. Guralnik, C.R. Hagen and T. W. B. Kibble Phys.Rev.Lett. **13** (1964) 585.

- [15] N. P. Nilles, *Supersymmetry, supergravity and particle physics*, Phys. Rep. **110** (1984) 1.
- [16] H. E. Haber and G. L. Kane, *The search for supersymmetry: Probing physics beyond the standard model*, Phys. Rep. **117** (1985) 75.
- [17] ALEPH, DELPHI, L3 and OPAL Collaboration, *Search for neutral MSSM Higgs bosons at LEP*, Eur. Phys. J. **C47** (2006) 547.
- [18] *Combined CDF and D0 upper limits on MSSM Higgs boson production in tau-tau final states with up to  $2.2 \text{ fb}^{-1}$  of data*, arXiv:1003.3363 [hep-ex].
- [19] The CMS Collaboration, S. Chatrchyan et al., arXiv:1104.1619 [hep-ex] [hep-ex].
- [20] The ATLAS Collaboration, *Search for the neutral Higgs bosons of the Minimal Supersymmetric Standard Model in  $pp$  collisions at  $\sqrt{s} = 7 \text{ TeV}$  with the ATLAS detector*, arXiv:1211.6956 [hep-ex].
- [21] S. Heinemeyer, O. Stål and G. Weiglein, *Interpreting the LHC Higgs search results in the MSSM*, Phys.Lett. **B710** (2012) 201–206, arXiv:1112.3026 [hep-ph].
- [22] A. Arbey, M. Battaglia, A. Djouadi and F. Mahmoudi, *The Higgs sector of the phenomenological MSSM in the light of the Higgs boson discovery*, JHEP **1209** (2012) 107, arXiv:1207.1348 [hep-ph].
- [23] The ATLAS Collaboration, G. Aad et al., *The ATLAS Experiment at the CERN Large Hadron Collider*, JINST **3** (2008) S08003.
- [24] M. L. Mangano et al., *ALPGEN, a generator for hard multiparton processes in hadronic collisions*, JHEP **07** (2003) 001.
- [25] J. Alwall et al., *Comparative study of various algorithms for the merging of parton showers and matrix elements in hadronic collisions*, Eur. Phys. J. **C53** (2008) 473, arXiv:0706.2569.
- [26] S. Frixione and B. R. Webber, *Matching NLO QCD computations and parton shower simulations*, JHEP **06** (2002) 029, hep-ph/0204244.
- [27] B. P. Kersevan and E. Richter-Was, *The Monte Carlo Event Generator AcerMC 2.0 with Interfaces to PYTHIA 6.2 and HERWIG 6.5*, arXiv:0405247v1 [hep-ph].
- [28] G. Corcella et al., *HERWIG 6: an event generator for hadron emission reactions with interfering gluons (including supersymmetric processes)*, JHEP **01** (2001) 010.
- [29] J. M. Butterworth, J. R. Forshaw, and M. H. Seymour, *Multiparton Interactions in Photoproduction at HERA*, Z. Phys. **C72** (1996) 637.

- [30] T. Binoth, M. Ciccolini, N. Kauer, and M. Kramer, *Gluon-induced W-boson pair production at the LHC*, JHEP **12** (2006) 046.
- [31] A. S. et al., *Higgs boson production in gluon fusion*, JHEP **02** (2009) 029.
- [32] T. Gleisberg et al., *Event generation with SHERPA 1.1*, JHEP **02** (2009) 007.
- [33] J. Pumplin, D. R. Stump, J. Huston, H. L. Lai, P. M. Nadolsky and W. K. Tung, “New generation of parton distributions with uncertainties from global QCD analysis,” JHEP **0207** (2002) 012 [hep-ph/0201195].
- [34] H. -L. Lai, M. Guzzi, J. Huston, Z. Li, P. M. Nadolsky, J. Pumplin and C. - P. Yuan, “New parton distributions for collider physics,” Phys. Rev. D **82** (2010) 074024 [arXiv:1007.2241 [hep-ph]].
- [35] S. Heinemeyer *et al.* [LHC Higgs Cross Section Working Group Collaboration], “Handbook of LHC Higgs Cross Sections: 3. Higgs Properties,” arXiv:1307.1347 [hep-ph].
- [36] M. Carena, S. Heinemeyer, C. E. M. Wagner, and G. Weiglein, *Suggestions for benchmark scenarios for MSSM Higgs boson searches at hadron colliders*, Eur. Phys. J. **C26** (2003) 601–607, hep-ph/0202167.
- [37] The ATLAS Collaboration, *ATLAS Monte Carlo Tunes for MC09*, ATL-PHYS-PUB-2010-002.
- [38] S. Jadach, J. H. Kuhn and Z. Was, *TAUOLA - a library of Monte Carlo programs to simulate decays of polarized  $\tau$  leptons*, Comput. Phys. Commun. **64** (1990) 275.
- [39] E. Barberio, B. V. Eijk and Z. Was, *Photos - a universal Monte Carlo for QED radiative corrections in decays*, Comput. Phys. Commun. **66** (1991) 115.
- [40] The GEANT4 Collaboration, S. Agostinelli et al., *GEANT4 - a simulation toolkit*, Nucl. Instrum. Meth. **A506** (2003) 250.
- [41] The ATLAS Collaboration, G. Aad et al., *The ATLAS Simulation Infrastructure*, ATLAS-SOFT-2010-01-004, submitted to Eur. Phys. J. C., arXiv:1005.4568.
- [42] The ATLAS Collaboration, *Estimation of  $Z/\gamma^* \rightarrow \tau\tau$  Background in VBF  $H \rightarrow \tau\tau$  Searches from  $Z \rightarrow \mu\mu$  Data using an Embedding Technique*, ATL-PHYS-INT-2009-109.
- [43] The ATLAS Collaboration, *Search for the Standard Model Higgs boson in the  $H \rightarrow \tau\tau$  decay mode with  $4.7\text{ fb}^{-1}$  of ATLAS detector*, Tech. Rep. ATLAS-CONF-2012-014, CERN, Geneva, Mar, 2012.
- [44] The ATLAS Collaboration, *Search for the Standard Model Higgs boson  $H \rightarrow \tau\tau$  decays with the ATLAS detector*, ATL-COM-PHYS-2013-722.

- [45] T. S. et al., *Z physics at LEP 1*, CERN 89-08 **3** (1989) 143.
- [46] The ATLAS Collaboration, Inner Detector: Technical Design Report, CERN/LHCC/97-016/017 (1997).
- [47] The ATLAS Collaboration, G. Aad et al., The ATLAS Experiment at the CERN Large Hardon Collider, 2008 JINST 3 S08003.
- [48] A. Bazan, T. Bouedo, P. Ghez, M. Marino and C. Tull, “The Athena data dictionary and description language,” eConf C **0303241** (2003) MOJT010 [cs/0305049 [cs-se]].
- [49] The ATLAS Collaboration, *Expected Performance of the ATLAS Experiment - Detector, Trigger and Physics*, CERN-OPEN-2008-020, arXiv:0901.0512.
- [50] T. Cornelissen et al., Concepts, Design and Implementation of the ATLAS New Tracking, ATLAS Note ATL-SOFT-PUB-2007-007 (2007).
- [51] Kalman, R. E. (1960). “*A New Approach to Linear Filtering and Prediction Problems*”. Journal of Basic Engineering 82 (1): 3545. doi:10.1115/1.3662552
- [52] The ATLAS Collaboration, Performance of primary vertex reconstruction in proton-proton collisions at  $s = \sqrt{s}(7)$  TeV in the ATLAS experiment. ATLAS-CONF-2010-069.
- [53] R. Fruhwirth, W. Waltenberger, P. Vanlaer, *Adaptive vertex fitting*, J. Phys. G34 (2007).
- [54] The ATLAS Collaboration, *Characterization of Interaction-Point Beam Parameters Using the pp Event-Vertex Distribution Reconstructed in the ATLAS Detector at the LHC*, ATL-CONF-2010-027.
- [55] The ATLAS collaboration, *Expected electron performance in the ATLAS experiment*, ATL-PHYS-PUB-2011-006
- [56] The ATLAS Collaboration, *Electron reconstruction and identification efficiency measurements with the ATLAS detector using the 2011 LHC proton-proton collision data*, CERN-PH-EP-2014-040, arXiv:1404.2240
- [57] The ATLAS Collaboration, G. Aad et al., *Electron performance measurements with the ATLAS detector using the 2010 LHC proton-proton collision data*, Eur.Phys.J. C72 (2012) 1909.
- [58] S. Hassini, et al., *A muon identification and combined reconstruction procedure for the ATLAS detector at the LHC using the (MUONBOY, STACO, MuTag) reconstruction packages*, NIM A572 (2007) 7779.
- [59] The ATLAS Collaboration, G. Aad et al., *Preliminary results on the muon reconstruction efficiency, momentum resolution, and momentum scale in ATLAS 2012 pp collision data*, ATLAS-CONF-2013-088, CERN, 2013,

- [60] M. Cacciari, G. P. Salam, and G. Soyez, *FastJet user manual*, Eur.Phys.J. C72 (2012) 1896.
- [61] W. Lampl et al., *Calorimeter Clustering Algorithms : Description and Performance*, ATL-LARG-PUB-2008-002.
- [62] M. Cacciari, G. P. Salam, and G. Soyez, *The anti-kt jet clustering algorithm*, JHEP 04 (2008) 63.
- [63] E. Abat, J. Abdallah, T. Addy, P. Adragna, et al., *Combined performance studies for electrons at the 2004 ATLAS combined test-beam*, JINST 5 (2010) P11006.
- [64] ATLAS Collaboration, *Jet energy measurement with the ATLAS detector in proton-proton collisions at  $\sqrt{s} = 7$  TeV*, Submitted to EPJ (2011) , arXiv:1112.6426
- [65] The ATLAS Collaboration, *Pile-up corrections for jets from proton-proton collisions at ATLAS in 2011*, ATLAS-CONF-2012-064, July, 2012.
- [66] M. Cacciari and G. P. Salam, *Pileup subtraction using jet areas*, Phys.Lett. B659 (2008) 119.
- [67] The ATLAS Collaboration, G. Aad et al., *Jet energy resolution in proton-proton collisions at  $\sqrt{s} = 7$  TeV recorded in 2010 with the ATLAS detector*, Eur.Phys.J. C73 (2013) 2306
- [68] The ATLAS collaboration, *Jet energy scale and its systematic uncertainty in proton-proton collisions at  $\sqrt{s} = 7$  TeV with ATLAS 2011 data*, ATLAS-CONF-2013-004
- [69] The ATLAS Collaboration, *Data-Quality Requirements and Event Cleaning for Jets and Missing Transverse Energy Reconstruction with the ATLAS Detector in Proton-Proton Collisions at a Center-of-Mass Energy of  $\sqrt{s} = 7$  TeV*, ATLAS-CONF-2010-038.
- [70] G. Piacquadio, C. Weiser, *A new inclusive secondary vertex algorithm for b-jet tagging in ATLAS*, JPCS 119 (2008) 032032
- [71] The ATLAS Collaboration, G. Aad et al., *Commissioning of the ATLAS high-performance b-tagging algorithms in the 7 TeV collision data*, ATLAS-CONF-2011-102, CERN, 2011, ATLAS-CONF-2011-102.
- [72] The ATLAS Collaboration, *Measuring the b-tag efficiency in a  $t\bar{t}$  sample with  $4.7\text{ fb}^{-1}$  of data from the ATLAS detector* ATLAS-CONF-2012-097.
- [73] The ATLAS Collaboration, *Calibration of b-tagging using dileptonic top pair events in a combinatorial likelihood approach with the ATLAS experiment* ATLAS-CONF-2014-004.

- [74] The ATLAS Collaboration, *Reconstruction and Calibration of Missing Transverse Energy and Performance in Z and W events in ATLAS Proton-Proton Collisions at  $\sqrt{s}=7$  TeV*, ATLAS-CONF-2011-080.
- [75] ATLAS Collaboration, G. Aad et al., Performance of missing transverse momentum reconstruction in proton-proton collisions at 7 TeV with ATLAS, Eur.Phys.J. C72 (2012) 1844.
- [76] TheATLAS Collaboration, *Performance of the Reconstruction and Identification of Hadronic tau Decays in ATLAS with 2011 Data*, ATLAS-CONF-2012-142.
- [77] The ATLAS Collaboration, G. Aad et al., *Performance of the ATLAS trigger system in 2010*, Eur.Phys.J. C72 (2012) 1849.
- [78] The ATLAS Collaboration, G. Aad et al., *Performance of the ATLAS muon trigger in 2011*, ATLAS-CONF-2012-099, CERN, 2012.
- [79] The ATLAS Collaboration, G. Aad et al., *Performance of the ATLAS electron and photon trigger in p-p collisions at  $\sqrt{s} = 7$  TeV in 2011*, ATLAS-CONF-2012-048, CERN, 2012.
- [80] M. Dobbs and J.B. Hansen, *The HepMC C++ Monte Carlo Event Record for High Energy Physics*, Computer Physics Communications, ATL-SOFT-2000-001.
- [81] The ATLAS Collaboration, *Observation of a new particle in the search for the Standard Model Higgs boson with the ATLAS detector at the LHC*, Physics Letters B **716** (2012) 1–29.
- [82] The CMS Collatoration, *Observation of a new boson at a mass of 125 GeV with the CMS experiment at the LHC*, Physics Letters B **716** (2012) 30–61.
- [83] The ATLAS Collaboration, *Evidence for the spin-0 nature of the Higgs boson using ATLAS data*, Phys. Lett. B 726 (2013), pp. 120-144.
- [84] The ATLAS Collaboration, *Measurements of Higgs boson production and couplings in diboson final states with the ATLAS detector at the LHC*, Phys. Lett. B 726 (2013), pp. 88-119.
- [85] The CMS Collaboration, *“Evidence for the direct decay of the 125 GeV Higgs boson to fermions,”* arXiv:1401.6527 [hep-ex].
- [86] The CMS Collaboration, *Higgs boson width from on- vs. off-shell production and decay to Z-boson pairs*, arXiv:1405.3455.
- [87] A. Elagin, P. Murat, A. Pranko, and A. Safonov, *A New Mass Reconstruction Technique for Resonances Decaying to di-tau*, arXiv:1012.4686 [hep-ex]. \* Temporary entry \*.
- [88] T. A. Collaboration, *Search for neutral MSSM Higgs bosons decaying to  $\tau\tau$  pairs in proton-proton collisions at with the ATLAS detector*, Physics Letters B **705** (2011) no. 3, 174 – 192.

- [89] The ATLAS Collaboration, *Data-driven estimation of the background to charged Higgs boson searches using hadronically-decaying tau final states in ATLAS*, ATLAS-CONF-2011-051.
- [90] The ATLAS Collaboration, *Measurement of the  $Z \rightarrow \tau\tau$  cross section with the ATLAS detector*, Phys. Rev. D **84** (2011) 112006.
- [91] T. A. Collaboration, *Search for the neutral Higgs bosons of the Minimal Supersymmetric Standard Model in pp collisions at  $\sqrt{s} = 7$  TeV with the ATLAS detector*, JHEP , [arXiv:1211.6956](#).
- [92] Atlas statistics forum, *ABCD method in searches*, [link](#)
- [93] The ATLAS Collaboration, *Search for Neutral MSSM Higgs Bosons  $H$  to  $\tau\tau$  to  $l\tau_h$  with the ATLAS Detector in 7 TeV Collisions*, ATL-COM-PHYS-2012-094.
- [94] The ATLAS Collaboration, *Search for neutral Higgs Bosons in the decay mode  $H \rightarrow \tau\tau \rightarrow ll+4\nu$  in proton proton collision at  $\sqrt{7}$  TeV with the ATLAS Detector*, ATL-COM-PHYS-2011-758.
- [95] The Atlas Collaboration, *Measurement of the  $t\bar{t}$  production cross-section in pp collisions at  $\sqrt{s} = 8$  TeV using e-mu events with b-tagged jets* . ATLAS-CONF-2013-097.
- [96] The ATLAS Collaboration, *Luminosity Determination in pp Collisions at  $\sqrt{s} = 7$  TeV using the ATLAS Detector in 2011*, ATLAS-CONF-2011-116.
- [97] T. Sjostrand, S. Mrenna and P. Skands, *PYTHIA 6.4 physics and manual*, JHEP **05** (2006) 026.
- [98] A. B. et al., *Rivet user manual*, [arXiv:1003.0694](#) [hep-ph].
- [99] The ATLAS and CMS Collaborations, *Procedure for the LHC Higgs boson search combination in summer 2011*, ATL-PHYS-PUB-2011-011, ATL-COM-PHYS-2011-818, CMS-NOTE-2011-005.
- [100] W. Verkerke and D. P. Kirkby, *The RooFit Toolkit for Data Modeling*, eConf C0303241 (2003) MOLT007, [arXiv:physics/0306116](#)
- [101] K. S. Cranmer et al., *The Roostats Project*, PoS ACAT2010 (2010) 57.
- [102] K. Cranmer, G. Lewis, L. Moneta, A. Shibata, and W. Verkerke, *Histfactory: A Tool for Creating Statistical Models for use with RooFit and Roostats*, CERNOPEN2012016 (2012)
- [103] E. G. G. Cowan, K. Cranmer and O. Vitells, *Asymptotic formulae for likelihood-based tests of new physics*, [arXiv:1007.1727](#) [hep-ex].
- [104] A. L. Read. *Presentation of search results: the CLs technique*. J. Phys. G: Nucl. Part. Phys., 28, 2002.

- [105] A. L. Read. *Modified frequentist analysis of search results (the CLs method)*. In Proceedings of the First Workshop on Confidence Limits, CERN, Geneva, Switzerland, 2000.
- [106] LHC Higgs Cross Section Working Group, S. Dittmaier, C. Mariotti, G. Passarino, R. Tanaka (Eds.), et al., *Handbook of LHC Higgs Cross Sections: 1. Inclusive Observables*, arXiv:1101.0593 [hep-ph].
- [107] LHC Higgs Cross Section Working Group, S. Dittmaier, C. Mariotti, G. Passarino, and R. Tanaka (Eds.), *Handbook of LHC Higgs Cross Sections: 2. Differential Distributions*, CERN-2012-002 (CERN, Geneva, 2012) , arXiv:1201.3084 [hep-ph].
- [108] ATLAS collaboration *Performance of the ATLAS Silicon Pattern Recognition Algorithm in Data and Simulation at  $\sqrt{s} = 7$  TeV*, ATLAS-CONF-2010-072
- [109] The ATLAS Collaboration, *A measurement of the material in the ATLAS inner detector using secondary hadronic interactions*, arXiv:1110.6191, JINST 7 (2012) P01013
- [110] The ATLAS Collaboration, *Validation of the ATLAS jet energy scale uncertainties using tracks in proton-proton collision  $\sqrt{s} = 7$  TeV*, ATLAS-CONF-2011-067
- [111] The ATLAS Collaboration, *Track Reconstruction Efficiency in  $\sqrt{s} = 7$  TeV Data for Tracks with  $P_T > 100$  MeV* , ATL-PHYS-INT-2010-112
- [112] D. de Florian, G. Ferrera, M. Grazzini and D. Tommasini, *Transverse-momentum resummation: Higgs boson production at the Tevatron and the LHC*, JHEP **1111** (2011) , arXiv:1109.2109 [hep-ph].
- [113] Statistical twiki, NuisanceCheck. <https://twiki.cern.ch/twiki/bin/view/AtlasProtected/NuisanceCheck>