

Contents

1	Prospects for Neutral MSSM Higgs Search Improvement	3
1.1	Introduction to Trackjets	4
1.2	Trackjet Performance	5
1.2.1	B-tagging Performance	5
1.2.2	Comparison Calo-jet Track-jet	5
1.2.3	Impact of Trackjet to the Analysis	5
1.2.4	B-Discriminant	5
1.3	Systematic Uncertainties on Trackjets	5
1.3.1	General discussion	5
1.3.2	Track Subtraction Method	5
1.3.3	Track Subtraction Validation	5
1.3.4	Track Subtraction Results	5

Chapter 1

Prospects for Neutral MSSM Higgs Search Improvement

The neutral MSSM Higgs search, described in the previous chapter, suffers strongly of poor jet reconstruction efficiency and b-tagging performance due to the particular phase space required, this bound the potential of this search, improving b-tagging would result in a major improvement of the search sensitivity. This chapter investigates an alternative to the commonly used calorimeter jets in ATLAS, which is trackjets b-tagging. The prospects for successfully use trackjets b-tagging in the future neutral MSSM Higgs searches are reported, b-tagging on trackjets was never attempted before. Section?? describes this and that...




Figure 1.1: Comparison of simulated b-hadron distribution for signal b-associated production events (left) and $t\bar{t}$ events (right). The red line in the figure shows the acceptance region due to calibrated jet p_t requirements.

1.1 Introduction to Trackjets

This problematic has two sources: - The ATLAS calorimeter is not a sampling calorimeter, this means that responses differently for Hadrons and for leptons, has different responses to electromagnetic and hadronic shower. The Calorimeter cells are calibrated in energy using response to electromagnetic showers, to know the energy of the original parton that initiated the jet there are different procedure to calibrate the Jets offline which are called in short Jet Energy Scale (JES) corrections [], which make use of MC simulation. Due to the high amount of pileup and ambient energy density in the events, jets are calibrated from 20 GeV in p_t , this means that currently is not possible with calorimeter jets to access the low transverse momentum phase space.

The neutral MSSM Higgs search, as described in chapter ??, splits the dataset in two category by means of the presence or the absence of a b-tagged jet, the b-tagged category is optimized for the b-associated production mechanism, in which the Higgs is produced in association with two b-jets. Figure 1.1 shows a comparison between p_t spectrum of simulated b-hadron in b-associated Higgs and $t\bar{t}$ events, the signal prefers b-hadron with relatively low transverse momentum, jet calibration involve jet $p_t > 20$ GeV removing a large fraction of possible signal candidate, many of the b-associated production signal events falls in fact in the b-veto category, making the separation not so effective. The low p_t spectrum is actually quite challenging, jet reconstruction efficiency and calibration set then a lower limit to the signal sensitivity in the b-tag category. Another challenge to this search are the poor b-tagging performance at low transverse momentum, for the a fixed tagging point of the MV1 tagger the b-tagging efficiency drops, in fact, rapidly with jet p_t reaching a minimum of 50% at 20 GeV [65, 66] (using as tagging point the 70% point).

A solution to the jet reconstruction efficiency is to use, instead of calorimetric jets (calo-jet), track-jet, which are as well anti-kt object (see chapter??) but constructed using inner detector tracks as building blocks, not calorimeter cells. Jets in the ATLAS reconstruction software are reconstructed by clustering four vector

objects (calorimeter energy cluster, tracks, truth particle, etc.) in the $\eta - \phi$ plane. In the case of clustering tracks, however, it is possible to take advantage of the longitudinal (z) impact parameter information provided by the inner detector and build track-jets in three dimensions $\eta - \phi - z$. Track-jets will then contain only tracks originating from the same interaction point (reconstructed vertex). Even though for calorimeter jet it is possible to use the JVF, track-jets result to be more resistant to decrease in performance in the presence of pile-up, thus particularly important in b-tagging, which depends on the determination of the jet-axis. B-tagging has been never tested before on track-jets, in the following, the first study of b-tagging over track-jets performances is reported.

1.2 Trackjet Performance

1.2.1 B-tagging Performance

1.2.2 Comparison Calo-jet Track-jet

1.2.3 Impact of Trackjet to the Analysis

1.2.4 B-Discriminant

1.3 Systematic Uncertainties on Trackjets

1.3.1 General discussion

1.3.2 Track Subtraction Method

1.3.3 Track Subtraction Validation

1.3.4 Track Subtraction Results

Bibliography

- [1] L. Evans and P. Bryant, *LHC Machine*, JINST **3** (2008) S08001.
- [2] F. Englert and R. Brout, *Broken Symmetry and the Mass of Gauge Vector Mesons*, Phys. Rev. Lett. **13** (1964) 321.
- [3] P. W. Higgs, *Broken symmetries, massless particles and gauge fields*, Phys. Lett. **12** (1964) 132.
- [4] P. W. Higgs, *Broken Symmetries and the Masses of Gauge Bosons*, Phys. Rev. Lett. **13** (1964) 508.
- [5] P. W. Higgs, *Spontaneous Symmetry Breakdown without Massless Bosons*, Phys. Rev. **145** (1966) 1156.
- [6] G. S. Guralnik, C.R. Hagen and T. W. B. Kibble Phys.Rev.Lett. **13** (1964) 585.
- [7] N. P. Nilles, *Supersymmetry, supergravity and particle physics*, Phys. Rep. **110** (1984) 1.
- [8] H. E. Haber and G. L. Kane, *The search for supersymmetry: Probing physics beyond the standard model*, Phys. Rep. **117** (1985) 75.
- [9] ALEPH, DELPHI, L3 and OPAL Collaboration, *Search for neutral MSSM Higgs bosons at LEP*, Eur. Phys. J. **C47** (2006) 547.
- [10] *Combined CDF and D0 upper limits on MSSM Higgs boson production in tau-tau final states with up to 2.2 fb^{-1} of data*, arXiv:1003.3363 [hep-ex].
- [11] CDF Collaboration, T. Aaltonen et al. Phys. Rev. Lett. **103** (2009) 201801.
- [12] D0 Collaboration, V. Abazov et al. Phys. Rev. Lett. **101** (2008) 071804.
- [13] TNPWG (Tevatron New Physics Higgs Working Group), CDF and D0 Collaborations, *Search for Neutral Higgs Bosons in Events with Multiple Bottom Quarks at the Tevatron*, arXiv:1207.2757 [hep-ex].
- [14] CDF Collaboration, T. Aaltonen et al., *Search for Higgs Bosons Produced in Association with b-quarks*, Phys.Rev. **D85** (2012) 032005, arXiv:1106.4782 [hep-ex].

- [15] D0 Collaboration, V.M. Abazov et al., *Search for neutral Higgs bosons in the multi-b-jet topology in 5.2fb^{-1} of $p\bar{p}$ collisions at $\sqrt{s} = 1.96\text{ TeV}$* , Phys.Lett. **B698** (2011) 97–104, [arXiv:1011.1931](#) [hep-ex].
- [16] The CMS Collaboration, S. Chatrchyan et al., [arXiv:1104.1619](#) [hep-ex] [hep-ex].
- [17] The ATLAS Collaboration, *Search for the neutral Higgs bosons of the Minimal Supersymmetric Standard Model in pp collisions at $\sqrt{s} = 7\text{ TeV}$ with the ATLAS detector*, [arXiv:1211.6956](#) [hep-ex].
- [18] T. A. Collaboration, *Observation of a new particle in the search for the Standard Model Higgs boson with the ATLAS detector at the LHC*, Physics Letters B **716** (2012) 1–29.
- [19] T. C. Collatoration, *Observation of a new boson at a mass of 125 GeV with the CMS experiment at the LHC*, Physics Letters B **716** (2012) 30–61.
- [20] S. Heinemeyer, O. Stål and G. Weiglein, *Interpreting the LHC Higgs search results in the MSSM*, Phys.Lett. **B710** (2012) 201–206, [arXiv:1112.3026](#) [hep-ph].
- [21] A. Arbey, M. Battaglia, A. Djouadi and F. Mahmoudi, *The Higgs sector of the phenomenological MSSM in the light of the Higgs boson discovery*, JHEP **1209** (2012) 107, [arXiv:1207.1348](#) [hep-ph].
- [22] The ATLAS Collaboration, G. Aad et al., *The ATLAS Experiment at the CERN Large Hadron Collider*, JINST **3** (2008) S08003.
- [23] M. L. Mangano et al., *ALPGEN, a generator for hard multiparton processes in hadronic collisions*, JHEP **07** (2003) 001.
- [24] J. Alwall et al., *Comparative study of various algorithms for the merging of parton showers and matrix elements in hadronic collisions*, Eur. Phys. J. **C53** (2008) 473, [arXiv:0706.2569](#).
- [25] S. Frixione and B. R. Webber, *Matching NLO QCD computations and parton shower simulations*, JHEP **06** (2002) 029, [hep-ph/0204244](#).
- [26] B. P. Kersevan and E. Richter-Was, *The Monte Carlo Event Generator AcerMC 2.0 with Interfaces to PYTHIA 6.2 and HERWIG 6.5*, [arXiv:0405247v1](#) [hep-ph].
- [27] G. Corcella et al., *HERWIG 6: an event generator for hadron emission reactions with interfering gluons (including supersymmetric processes)*, JHEP **01** (2001) 010.
- [28] J. M. Butterworth, J. R. Forshaw, and M. H. Seymour, *Multiparton Interactions in Photoproduction at HERA*, Z. Phys. **C72** (1996) 637.

- [29] T. Binoth, M. Ciccolini, N. Kauer, and M. Kramer, *Gluon-induced W -boson pair production at the LHC*, JHEP **12** (2006) 046.
- [30] A. S. et al., *Higgs boson production in gluon fusion*, JHEP **02** (2009) 029.
- [31] T. Gleisberg et al., *Event generation with SHERPA 1.1*, JHEP **02** (2009) 007.
- [32] J. Pumplin, D. R. Stump, J. Huston, H. L. Lai, P. M. Nadolsky and W. K. Tung, “New generation of parton distributions with uncertainties from global QCD analysis,” JHEP **0207** (2002) 012 [hep-ph/0201195].
- [33] H. -L. Lai, M. Guzzi, J. Huston, Z. Li, P. M. Nadolsky, J. Pumplin and C. - P. Yuan, “New parton distributions for collider physics,” Phys. Rev. D **82** (2010) 074024 [arXiv:1007.2241 [hep-ph]].
- [34] M. Carena, S. Heinemeyer, C. E. M. Wagner, and G. Weiglein, *Suggestions for benchmark scenarios for MSSM Higgs boson searches at hadron colliders*, Eur. Phys. J. **C26** (2003) 601–607, hep-ph/0202167.
- [35] The ATLAS Collaboration, *ATLAS Monte Carlo Tunes for MC09*, ATL-PHYS-PUB-2010-002.
- [36] S. Jadach, J. H. Kuhn and Z. Was, *TAUOLA - a library of Monte Carlo programs to simulate decays of polarized τ leptons*, Comput. Phys. Commun. **64** (1990) 275.
- [37] E. Barberio, B. V. Eijk and Z. Was, *Photos - a universal Monte Carlo for QED radiative corrections in decays*, Comput. Phys. Commun. **66** (1991) 115.
- [38] The GEANT4 Collaboration, S. Agostinelli et al., *GEANT4 - a simulation toolkit*, Nucl. Instrum. Meth. **A506** (2003) 250.
- [39] The ATLAS Collaboration, G. Aad et al., *The ATLAS Simulation Infrastructure*, ATLAS-SOFT-2010-01-004, submitted to Eur. Phys. J. C., arXiv:1005.4568.
- [40] The ATLAS Collaboration, *Estimation of $Z \rightarrow \tau\tau$ Background in VBF $H \rightarrow \tau\tau$ Searches from $Z \rightarrow \mu\mu$ Data using an Embedding Technique*, ATL-PHYS-INT-2009-109.
- [41] The ATLAS Collaboration, *Search for the Standard Model Higgs boson in the $H \rightarrow \tau\tau$ decay mode with 4.7 fb of ATLAS detector*, Tech. Rep. ATLAS-CONF-2012-014, CERN, Geneva, Mar, 2012.
- [42] The ATLAS Collaboration, *Search for the Standard Model Higgs boson $H \rightarrow \tau\tau$ decays with the ATLAS detector*, ATL-COM-PHYS-2013-722.
- [43] T. S. et al., *Z physics at LEP 1*, CERN 89-08 **3** (1989) 143.
- [44] The ATLAS Collaboration, *Expected Performance of the ATLAS Experiment - Detector, Trigger and Physics*, CERN-OPEN-2008-020, arXiv:0901.0512.

- [45] The ATLAS Collaboration, *ATLAS Muon Momentum Resolution in the First Pass Reconstruction of the 2010 p-p Collision Data at $\sqrt{s} = 7$ TeV*, ATLAS-CONF-2011-046.
- [46] The ATLAS Collaboration, *Muon reconstruction efficiency in reprocessed 2010 LHC p-p collision data recorded with the ATLAS detector*, ATLAS-CONF-2011-063.
- [47] The ATLAS Collaboration, *Expected electron performance in the ATLAS experiment*, ATLAS-PUB-2011-006.
- [48] ATLAS egamma WG, *Electron efficiency measurements*, <https://twiki.cern.ch/twiki/bin/view/AtlasProtected/EfficiencyMeasurements>.
- [49] M. Cacciari, G. P. Salam, and G. Soyez, *The anti- k_t jet clustering algorithm*, JHEP **04** (2008) 063.
- [50] W. Lampl et al., *Calorimeter Clustering Algorithms : Description and Performance*, ATL-LARG-PUB-2008-002.
- [51] T. Barillari et al., *Local Hadron Calibration*, ATL-LARG-PUB-2009-001.
- [52] The ATLAS Collaboration, *Jet energy scale and its systematic uncertainty in proton-proton collisions at $\sqrt{s} = 7$ TeV in ATLAS 2010 data*, ATLAS-CONF-2011-032.
- [53] The ATLAS Collaboration, *Performance of the Reconstruction and Identification of Hadronic tau Decays in ATLAS with 2011 Data*, ATLAS-CONF-2012-142.
- [54] The ATLAS Collaboration, *Reconstruction and Calibration of Missing Transverse Energy and Performance in Z and W events in ATLAS Proton-Proton Collisions at $\sqrt{s}=7$ TeV*, ATLAS-CONF-2011-080.
- [55] A. Elagin, P. Murat, A. Pranko, and A. Safonov, *A New Mass Reconstruction Technique for Resonances Decaying to di-tau*, arXiv:1012.4686 [hep-ex]. * Temporary entry *.
- [56] ATLAS Jet/EtMiss Combined Performance Group, *Jet Energy Resolution Provider*, <https://twiki.cern.ch/twiki/bin/view/Main/JetEnergyResolutionProvider>.
- [57] The ATLAS Collaboration, *Data-Quality Requirements and Event Cleaning for Jets and Missing Transverse Energy Reconstruction with the ATLAS Detector in Proton-Proton Collisions at a Center-of-Mass Energy of $\sqrt{s} = 7$ TeV*, ATLAS-CONF-2010-038.
- [58] T. A. Collaboration, *Search for neutral MSSM Higgs bosons decaying to $\tau\tau$ pairs in proton-proton collisions at with the ATLAS detector*, Physics Letters B **705** (2011) no. 3, 174 – 192.

- [59] The ATLAS Collaboration, *Data-driven estimation of the background to charged Higgs boson searches using hadronically-decaying tau final states in ATLAS*, ATLAS-CONF-2011-051.
- [60] The ATLAS Collaboration, *Measurement of the $Z \rightarrow \tau\tau$ cross section with the ATLAS detector*, Phys. Rev. D **84** (2011) 112006.
- [61] T. A. Collaboration, *Search for the neutral Higgs bosons of the Minimal Supersymmetric Standard Model in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector*, JHEP , [arXiv:1211.6956](#).
- [62] Atlas statistics forum, *ABCD method in searches*, [link](#)
- [63] The ATLAS Collaboration, *Search for Neutral MSSM Higgs Bosons H to $\tau\tau$ to $b\tau_h$ with the ATLAS Detector in 7 TeV Collisions*, ATL-COM-PHYS-2012-094.
- [64] The ATLAS Collaboration, *Search for neutral Higgs Bosons in the decay mode $H \rightarrow \tau\tau \rightarrow ll+4\nu$ in proton proton collision at $\sqrt{7}$ TeV with the ATLAS Detector*, ATL-COM-PHYS-2011-758.
- [65] The ATLAS Collaboration, *Measuring the b -tag efficiency in a $t\bar{t}b\bar{b}$ sample with 4.7 fb^{-1} of data from the ATLAS detector* ATLAS-CONF-2012-097.
- [66] The ATLAS Collaboration, *Calibration of b -tagging using dileptonic top pair events in a combinatorial likelihood approach with the ATLAS experiment* ATLAS-CONF-2014-004.
- [67] The ATLAS Collaboration, *Luminosity Determination in pp Collisions at $\sqrt{s} = 7$ TeV using the ATLAS Detector in 2011*, ATLAS-CONF-2011-116.
- [68] T. Sjostrand, S. Mrenna and P. Skands, *PYTHIA 6.4 physics and manual*, JHEP **05** (2006) 026.
- [69] A. B. et al., *Rivet user manual*, [arXiv:1003.0694](#) [[hep-ph](#)].
- [70] E. G. G. Cowan, K. Cranmer and O. Vitells, *Asymptotic formulae for likelihood-based tests of new physics*, [arXiv:1007.1727](#) [[hep-ex](#)].
- [71] LHC Higgs Cross Section Working Group, S. Dittmaier, C. Mariotti, G. Passarino, R. Tanaka (Eds.), et al., *Handbook of LHC Higgs Cross Sections: 1. Inclusive Observables*, [arXiv:1101.0593](#) [[hep-ph](#)].
- [72] LHC Higgs Cross Section Working Group, S. Dittmaier, C. Mariotti, G. Passarino, and R. Tanaka (Eds.), *Handbook of LHC Higgs Cross Sections: 2. Differential Distributions*, CERN-2012-002 (CERN, Geneva, 2012) , [arXiv:1201.3084](#) [[hep-ph](#)].
- [73] D. de Florian, G. Ferrera, M. Grazzini and D. Tommasini, *Transverse-momentum resummation: Higgs boson production at the Tevatron and the LHC*, JHEP **1111** (2011) , [arXiv:1109.2109](#) [[hep-ph](#)].
- [74] Statistical twiki, NuisanceCheck. <https://twiki.cern.ch/twiki/bin/view/AtlasProtected/NuisanceCheck>