

Time spent and left of transient states in stationary populations

Tim Riffe^{*1} and Francisco Villavicencio^{2,3}

¹Max Planck Institute for Demographic Research, Rostock, Germany

²Max-Planck Odense Center on the Biodemography of Aging, Department of Public Health,
University of Southern Denmark, Odense, Denmark

October 25, 2017

Abstract

The Brouard-Carey equality establishes equality between the distributions of years-lived and years-left in perfectly stationary populations. This equality can be generalized to account for time spent and time left in states within multistate stationary populations. We provide an intuitive proof that the distribution of time spent and left in transient states is equal in multistate stationary populations. We speculate that this equality may be helpful under certain constraints to estimate the distribution of otherwise unobserved onset timing for health or other states.

1 Background

The Brouard-Carey equality establishes that the distributions of years-lived and years-left are identical in perfectly stationary populations (Brouard 1989, Vaupel 2009, Rao and Carey 2015, Villavicencio and Riffe 2016). A perfectly stationary population is either perfectly stationary because it is of infinite size or because it is finite and deterministically repeating, and in either case vital rate schedules must be fixed and the intrinsic growth rate constant at null.

Let's say that individuals in the stationary population can obtain different states over the life course. If vital rates do not vary by states, then it does not matter whether state transition rates are fixed or not, for in the aggregate the population remains stationary in the traditional sense. However, if vital rates depend on one's state, transition rates must also be fixed in order for stationarity to hold in the aggregate— This is the situation that we entertain in the following. The standard set of invariant quantities of course remains: birth cohorts and death cohorts are of fixed and equal size. Further, the Brouard-Carey equality still holds in the aggregate. Under these conditions, one may take for granted that the age-state structure of the stationary population is also invariant over time. Further, the distribution of state-specific tenures for individuals entering a state at a given age is a fixed attribute, by consequence of fixed transition and vital rate schedules.

^{*}riffe@demogr.mpg.de

This setup is not entirely contrived, for it corresponds with the assumptions of common multistate markov models, often used to calculate state expectancies. In the present, however, we are not bound to memoryless transition probabilities. Instead, we simply require the result of invariant age-state structure and invariant conditional age-state tenure distributions, which may result from either memoryless transitions or from arbitrarily linked dependencies. It is the fact of a fixed age-state structure on which we base the following observations. In essence, we end up with a fixed distribution of lifelines, each consisting in a sequence of state-specific durations.

Under fixed vital and transition rates, and zero growth, we introduce a new theorem, which is a more general version of the Brouard-Carey equality. Namely, it turns out that if there is some state s in the population, a randomly drawn individual from state s has equal probabilities of having entered s x years ago and exiting s in x years. This proposition is probably not intuitive at first glance, so we prove it, and then speculate as to how this equality may be put to good use.

2 An intuitive proof of transient symmetry

Theorem 2.1. Given a stationary population and fixed transition rates, the probability that a randomly selected individual is in state s and entered s x years ago is equal to the probability of being in state s and exiting in x years.

Proof. The proof of this statement follows five intuitive steps.

1. A duration can be represented as a line segment, potentially a subset of a life-line. Points along a single within-person duration can be sampled over time in arbitrarily fine time steps, δ . Each time step can then be used to bisect the segment, collecting two sets of values: 1) time spent in the state prior to the sample point, and 2) time left until exiting the state. If observations are evenly spaced, and values are truncated to the nearest δ , by way of complements these two sets will consist of the same values.

If the duration of the i^{th} individual is called d^i , the age of entry is a_L^i and the age of exit is a_R^i , such that $d^i = a_R^i - a_L^i$. The set of time spent values, A^i is defined as:


$$A^i = \left\{ \delta \cdot k \mid k \in \mathbb{Z}, 0 \leq k \leq \left\lfloor \frac{d_i}{\delta} \right\rfloor \right\} \quad , \quad (1)$$

where $\lfloor \dots \rfloor$ denotes truncation to the lower integer bound. The set of time left values, T^i is also:

$$T^i = \left\{ \delta \cdot k \mid k \in \mathbb{Z}, \left\lfloor \frac{d_i}{\delta} \right\rfloor \geq k \geq 0 \right\} \quad . \quad (2)$$

Figure 1 illustrates the construction of A^i and T^i in expressions (1) and (2). The central cut-point moves along the duration from the time of entry until the time of exit, creating two lists of values.


Figure 1: A lifeline of individual i showing the construction of the sets A^i and T^i .



2. If this individual is reborn every δ time step, destined to relive the exact same life course as the first, we end up with a set of identically aligned and identically long segments placed side-by-side, spaced apart by δ . This potentially large set of segments may be imagined as a plane, although this would only hold in the limit. One could in this setting take a census at a single point in time, collecting a set of time spent and time left values, each from a unique individual in sequence. It is clear that the two sets observed at a single time point but drawn from the population of perfect clones, will be identical to the first two sets that were observed of a single duration over its entire length, if the same δ -truncation operation is followed.

Figure 2 illustrates this notion with uniformly-spaced lifelines in a Lexis configuration. The vertical line indicates a hypothetical census at time t of the population of this cloned individual. At time t , the blue-highlighted segments indicate the set of time-spent values in A^i , and red-highlighted segments are the time-left elements of T^i .

Figure 2: The life of individual i repeated in δ time steps. A census with followup now constructs the sets A^i and T^i with values identical to the within-individual sets.



Formally, sets A^i and T^i consist of the same values as the previous, but coming from individuals born in a uniform series from between $t - a_R^i$ years ago until as recently as $t - a_L^i$ years ago. This demonstrates both period-cohort equality and time spent-left equality. The blue and red triangles in Figure 2 are simple rotations

of one another.

3. Assume we have a second individual from the same birth cohort as individual i that enters the same state as the first, but at a different time and for a different total duration. We could demonstrate time spent and time left equality in the same way for this individual, by sampling in δ time steps. Since this individual has a different life course timing than the first individual, their sets of time-spent, A^2 , and time left T^2 will be distinct. A^1 and T^1 range from 0 to d^1 , whereas A^2 and T^2 range from 0 to d^2 . However, their concatenations are identical:

$$\{A^1, A^2\} = \{T^1, T^2\} \quad (3)$$

4. If the second individual is also perfectly cloned in δ time steps as in step 2, then our census at a single point in time also yields identical sets of time spent and time left values, A^2 and T^2 . Also from this census, the concatenation of the first and second time-spent sets and the concatenation of the first and second time-left sets are guaranteed to be identical, as in equation (3).
5. By induction we can keep adding durations in this state, infinitely if we please, and the collection of all resultant time-spent sets and the collection of time-left sets will continue being identical. Therefore the probability of selecting a particular value from the time-spent set is identical to the probability of selecting the same value from the time-left set.

□

3 Discussion

The approach from this proof is equally valid to prove the original statement of the Brouard-Carey equality, but it is more general. The statement and proof is so flexible as to hold for irreversible and reversible states. It is also applicable to repeatable states, whether time spent in the state is kept in cumulative fashion over spells, or whether the clock resets to zero on each entry into the state. The equality may also be conditionable in curious ways: for example, the distribution of time-spent in a state conditional on having entered at age a must also be equal to the distribution of time-left in the state, conditional on having entered at age a . Likewise, one may condition statements on exit age.

Prior to proof, this equality is probably less intuitive than the original Brouard-Carey equality, because state entry is not necessarily aligned on age zero. It is therefore less visible in commonly-produced plots. It might be tempting to think that due to state-varying vital rates, that the equality simply ought not hold. However, the basis of the proof is the observation that if each individual duration is symmetrical by complements, then so are aggregations of durations, irrespective of alignment. Since each cohort in a perfectly stationary population is an identical copy of the previous, census-like cross-sections are also equally-composed.

Empirical applications of the presently-described transient tenure equality may be easy to conjure up. For example, imagine a hypothetical health state that shows no noticeable symptoms, but that is medically measurable. One may take a census with regular follow-ups, until eventually the state is exited by each individual, whether by

absorption into death or entry into another state. Then, if the assumption of stationarity is acceptable, one may be able to say something about onset timing in the aggregate, itself unobserved. We think that the present equality will come as good news to researchers in similar settings.

The relationship between years lived and left established by the Brouard-Carey equality is itself amenable to non-zero growth rates (Riffe 2015). We speculate that the transient tenure equality we describe here may also be tractable in stable populations, but we have not investigated this possibility in detail.

References

- Brouard, N. (1989). *Mouvements et modèles de population*. Institut de formation et de recherche démographiques.
- Rao, A. S. S. and J. R. Carey (2015). Generalization of Carey’s equality and a theorem on stationary population. *Journal of Mathematical Biology* 71(3), 583–594.
- Riffe, T. (2015). Renewal and stability in populations structured by remaining years of life. *MPIDR Working Papers* (WP-2015-007).
- Vaupel, J. W. (2009). Life lived and left: Carey’s equality. *Demographic Research* 20, 7–10.
- Villavicencio, F. and T. Riffe (2016). Symmetries between life lived and left in finite stationary populations. *Demographic Research* 35(14), 381–398.