

# Symmetry in the forward and backward tenure of transient states in stationary populations

Tim Riffe<sup>\*1</sup> and Francisco Villavicencio<sup>2,3</sup>

<sup>1</sup>Max Planck Institute for Demographic Research, Rostock, Germany

<sup>2</sup>Max-Planck Odense Center on the Biodemography of Aging, Odense, Denmark

<sup>3</sup>Department of Public Health, University of Southern Denmark, Odense, Denmark

September 20, 2017

## Abstract

Guess what? The Brouard-Carey equality also applies to transient states in multistate stationary populations. That's even less intuitive than the original theorem because entries to and exits from states aren't aligned on either end. This property even holds if the transient state affects mortality, because age and TTD are beside the point. It'd only be practical if a census were at large.

## 1 sketch of proof

(1): A duration can be represented as a line segment, potentially a subset of a life-line. Points along a single within-person duration could be continuously sampled over time, used to bisect the segment, collecting two infinite sets of values: time spent in the duration and time until exiting the duration. If continuously collected, these two sets will have the same values, with time spent in ascending order and time left in descending order. If sorted identically these two sets will therefore be identical. This is so by way of complements.

(2): If this individual is reborn in each instant, destined to relive the exact same life course as the first, we end up with an infinite number of identically aligned identically long segments placed side-by-side. This infinite set of segments in effect forms a plane. One could in this setting take a census at a single point in time, collecting an infinite set of time spent and time left values, each from a unique cohort in sequence. It is clear that the two sets observed at a single time point will be identical to the first two sets that were observed of a single duration over its entire length.

(3): Assume we have another individual from the same birth cohort that enters the same state as the first, but at a different time and for a different duration. We could do (1) in the same way, sampling at an infinite set of points along the duration, and

---

<sup>\*</sup>riffe@demogr.mpg.de

constructing two sets, one of time spent and another of time left in the state. These too would be identical sets, but they would not be identical to those of the first individual, as the range of values would be different. The union of the first and second time-spent sets and the union of the first and second time-left sets are guaranteed to be identical.

(4): If the second individual repeats over time as in (2), then our comprehensive sample at a single point in time also yields identical sets of time spent and time left values. Also from this census, the union of the first and second time-spent sets and the union of the first and second time-left sets are guaranteed to be identical.

(5): By induction we can keep adding durations in this state, infinitely if we please, and the union of all resultant time spent sets and the union of time left sets will continue being identical. It is therefore true that the probability of selecting a particular value from the time-spent set is identical to the probability of selecting the same value from the time-left set. This constitutes a proof of symmetry of time spent and left in transient states in stationary populations.