

# Understanding Analysis Attempt/Solution

pand4monium

December 28, 2025



# Contents

<b>1</b>	<b>The Real Numbers</b>	<b>1</b>
1.2	Some Preliminaries . . . . .	1
1.3	The Axiom of Completeness . . . . .	6
1.4	Consequences of Completeness . . . . .	10
1.5	Cardinality . . . . .	12
1.6	Cantor's Theorem . . . . .	17
<b>2</b>	<b>Sequences and Series</b>	<b>21</b>
2.2	The Limit of a Sequence . . . . .	21
2.3	The Algebraic and Order Limit Theorem . . . . .	24
2.4	The Monotone Convergence Theorem and a First Look at Infinite Series . . . . .	29
2.5	Subsequences and the Bolzano-Weierstrass Theorem . . . . .	33
2.6	The Cauchy Criterion . . . . .	33
2.7	Properties of Infinite Series . . . . .	33
2.8	Double Summations and Products of Infinite Series . . . . .	33



# Chapter 1

## The Real Numbers

### 1.2 Some Preliminaries

#### Exercise 1.2.1

- (a) Prove that  $\sqrt{3}$  is irrational. Does a similar argument work to show  $\sqrt{6}$  is rational?
- (b) Where does the proof break down if we try to prove  $\sqrt{4}$  is irrational?

#### SOLUTION

- (a) AFSOC that  $\sqrt{3}$  is rational, so  $\exists m, n \in \mathbb{Z}$ , such that

$$\sqrt{3} = \frac{m}{n},$$

where  $\frac{m}{n}$  is in the lowest reduced terms. By squaring both sides, we obtain  $3 = (\frac{m}{n})^2 \implies 3n^2 = m^2$ . Now, we know that  $m^2$  is a multiple of 3 and thus  $m$  must also be a multiple of 3. We can then write  $m = 3k$ , deriving

$$\begin{aligned}(\sqrt{3})^2 &= \left(\frac{3k}{n}\right)^2 \\ 3n^2 &= 9k^2 \\ n^2 &= 3k^2\end{aligned}$$

Similar to above, we can conclude that  $n$  is a multiple of 3. However this is a contradiction since  $m, n$  are both multiples of 3 but we assumed that  $\frac{m}{n}$  was in its lowest reduced term. Thus we conclude that  $\sqrt{3}$  is irrational. The same proof for  $\sqrt{3}$  works for  $\sqrt{6}$  as well.

- (b) We cannot conclude that  $\sqrt{4} = \frac{m}{n}$  imply that  $m$  is a multiple of 4, as we have

$$4n^2 = m^2 \implies 2n = m,$$

preventing us from reaching our contradiction that  $m/n$  is not in its lowest terms.

#### Exercise 1.2.2

Show that there is no rational number  $r$  satisfying  $2^r = 3$ .

#### SOLUTION

If  $r = 0$ , then  $2^r = 1 \neq 3$ . Suppose  $r = p/q$  to get  $2^p = 3^q$ , which is not possible as 2 and 3 share no common factors. Hence  $r$  is not rational.

#### Exercise 1.2.3

Decide which of the following represent true statements about the nature of sets. For any that are false, provide a specific example where the statement in question does not hold.

- (a) If  $A_1 \supseteq A_2 \subseteq A_3 \subseteq A_4 \dots$  are all sets containing an infinite number of elements, then the intersections  $\bigcap_{n=1}^{\infty} A_n$  is infinite as well.
- (b) If  $A_1 \supseteq A_2 \subseteq A_3 \subseteq A_4 \dots$  are all finite, nonempty sets of real numbers, then the intersection  $\bigcap_{n=1}^{\infty} A_n$  is finite and non-empty.
- (c)  $A \cap (B \cup C) = (A \cap B) \cup C$ .
- (d)  $A \cap (B \cap C) = (A \cap B) \cap C$ .
- (e)  $A \cap (B \cup C) = (A \cap B) \cup (A \cap C)$ .

**SOLUTION**

- (a) False. Consider  $A_n = \{n, n+1, n+2, \dots\}$ , then  $\bigcap_{n=1}^{\infty} A_n = \emptyset$ .
- (b) True. Since all  $A_n$  are nonempty,  $\exists n \in \mathbb{N}$  such that  $A_n = \{x\}$  for some real  $x$ . Hence  $\bigcap_{n=1}^{\infty} A_n \subseteq \{x\}$  which is empty. Since  $A_1$  is finite,  $\bigcap_{n=1}^{\infty} A_n \subseteq \{x\} \subset A_1$  is finite.
- (c) False. If  $A = \emptyset$ , then  $\emptyset = C$
- (d) True. Intersection is associative as evident that both LHS and RHS implies the  $x \in A, B, C$
- (e) True. Drawing a Venn Diagram illustrates this.

**Exercise 1.2.4**

Produce an infinite collection of sets  $A_1, A_2, A_3, \dots$  with the property that every  $A_i$  has an infinite number of elements,  $A_i \cap A_j = \emptyset$  for all  $i \neq j$ , and  $\bigcup_{i=1}^{\infty} A_i = \mathbb{N}$ .

**SOLUTION**

Consider arranging the elements of  $\mathbb{N}$  in a square as such.

1	3	6	10	15	...
2	5	9	14	...	
4	8	13	...		
7	12	...			
11	...				
$\vdots$					

By letting  $A_i$  being the set of all natural numbers in the  $i$ -th row, we have satisfied the above conditions above.

**Exercise 1.2.5**

**(De Morgan's Law)** Let  $A$  and  $B$  be subsets of  $\mathbb{R}$ .

- (a) If  $x \in (A \cap B)^c$ , explain why  $x \in A^c \cup B^c$ . This shows that  $(A \cap B)^c \subseteq A^c \cup B^c$ .
- (b) Prove the reverse inclusion  $(A \cap B)^c \supseteq A^c \cup B^c$ , and conclude that  $(A \cap B)^c = A^c \cup B^c$ .
- (c) Show  $(A \cup B)^c = A^c \cap B^c$  by demonstrating inclusion both ways.

**SOLUTION**

- (a) If  $x \in (A \cap B)^c$ , then  $x \notin A \cap B$ , so  $x \notin A$  or  $x \notin B$ , implying  $x \in A^c$  or  $x \in B^c$ , therefore  $x \in A^c \cup B^c$ .
- (b) If  $x \in A^c \cup B^c$ , then  $x \in A^c$  or  $x \in B^c$ , so  $x \notin A$  and  $x \notin B$ , implying  $x \notin A \cap B$ , therefore  $x \in (A \cap B)^c$ . Since  $(A \cap B)^c \subseteq A^c \cup B^c$  and  $(A \cap B)^c \supseteq A^c \cup B^c$ , we can conclude that both sets are equal.
- (c) To show that  $(A \cap B)^c = A^c \cup B^c$ , we need to demonstrate inclusion both ways.
  - (i) If  $x \in (A \cup B)^c$ , then  $x \notin A \cup B$ , so  $x \notin A$  or  $x \notin B$ , implying  $x \in A^c$  or  $x \in B^c$ , therefore  $x \in A^c \cup B^c$ .

- (ii) If  $x \in A^c \cap B^c$ , then  $x \in A^c$  and  $x \in B^c$ , so  $x \notin A$  and  $x \notin B$ , implying  $x \notin A \cup B$ , which is just  $x \in (A \cup B)^c$ .

**Exercise 1.2.6**

- (a) Verify the triangle inequality in the special case where  $a$  and  $b$  have the same sign.  
 (b) Find an efficient proof for all the cases at once by first demonstrating  $(a + b)^2 \leq (|a| + |b|)^2$ .  
 (c) Prove  $|a - b| \leq |a - c| + |c - d| + |d - b|$  for all  $a, b, c$  and  $d$ .  
 (d) Prove  $||a| - |b|| \leq |a - b|$ . (The unremarkable identity  $a = a - b + b$  may be useful.)

**SOLUTION**

- (a) With both  $a$  and  $b$  having the same sign, then  $|a| + |b| = |a + b|$ , which satisfies  $|a| + |b| \geq |a + b|$ .  
 (b) Note that  $(a + b)^2 \leq (|a| + |b|)^2$  reduces to  $ab \leq |a||b|$ , which is true as LHS can be negative while RHS cannot. Since squaring preserves inequality, this implies that  $|a + b| \leq |a| + |b|$ .  
 (c) Notice that  $a - b = (a - c) + (c - d) + (d - b)$ . Hence by triangle inequality,

$$|a - b| = |(a - c) + (c - d) + (d - b)| \leq |a - c| + |c - d| + |d - b|$$

for all  $a, b, c$  and  $d$ .

- (d) Since  $||a| - |b|| = ||b| - |a||$ , WLOG, we can assume that  $|a| \geq |b|$ . Then

$$||a| - |b|| = |a| - |b| = |(a - b) + b| - |b| \leq |a - b| + |b| - |b| = |a - b|$$

**Exercise 1.2.7**

Given a function  $f$  and a subset  $A$  of its domain, let  $f(A)$  represent the range of  $f$  over the set  $A$ ; that is,  $f(A) = \{f(x) : x \in A\}$ .

- (a) Let  $f(x) = x^2$ . If  $A = [0, 2]$  (the closed interval  $\{x \in \mathbb{R} : 0 \leq x \leq 2\}$ ) and  $B = [1, 4]$ , find  $f(A)$  and  $f(B)$ . Does  $f(A \cap B) = f(A) \cap f(B)$  in this case? Does  $f(A \cup B) = f(A) \cup f(B)$ ?  
 (b) Find two sets  $A$  and  $B$  for which  $f(A \cap B) \neq f(A) \cap f(B)$ .  
 (c) Show that, for an arbitrary function  $g : \mathbb{R} \rightarrow \mathbb{R}$ , it is always true that  $g(A \cap B) \subseteq g(A) \cap g(B)$  for all sets  $A, B \subseteq \mathbb{R}$ .  
 (d) Form and prove a conjecture about the relationship between  $g(A \cup B)$  and  $g(A) \cup g(B)$  for an arbitrary function  $g$ .

**SOLUTION**

- (a) For  $f(x) = x^2$ ,  $f(A) = f([0, 2]) = [0, 4]$  and  $f(B) = f([1, 4]) = [1, 16]$ .

$$f(A \cap B) = f([0, 2] \cap [1, 4]) = f([1, 2]) = [1, 4] = [0, 4] \cap [1, 16] = f([1, 2]) \cap f([2, 4]) = f(A) \cap f(B)$$

$$f(A \cup B) = f([0, 2] \cup [1, 4]) = f([0, 4]) = [0, 16] = [0, 4] \cup [1, 16] = f([0, 2]) \cup f([1, 4]) = f(A) \cup f(B)$$

- (b) Consider  $A = [0, 2]$  and  $B = [-2, 0]$ .  $f(A \cap B) = \{0\}$ , but  $f(A) \cap f(B) = [0, 4]$ .  
 (c) Suppose  $y \in g(A \cap B)$ , then  $\exists x \in A \cap B$  such that  $g(x) = y$ . This implies that  $x \in A$  and  $x \in B$ , so  $x \in A \cap B$ , hence  $y \in g(A \cap B)$ . Note that contrary may not always be true as it is possible for  $x_1 \in A \setminus B$  and  $x_2 \in B \setminus A$  such that  $g(x_1) = g(x_2)$ .  
 (d) I conjecture that  $g(A \cup B) = g(A) \cup g(B)$ . To prove this, we have to show inclusion both ways:  
 (i) Let  $y \in g(A \cup B)$ , then  $\exists x \in A \cup B$  such that  $y = g(x)$ . This implies that  $x \in A$  or  $x \in B$ , so  $y \in g(A)$  or  $y \in g(B)$ , hence  $y \in g(A) \cup g(B)$ .  
 (ii) Let  $y \in g(A) \cup g(B)$ , then  $y \in g(A)$  or  $y \in g(B)$ , implying  $x \in A$  or  $x \in B$  such that  $y = g(x)$ . So  $x \in A \cup B$ , hence  $y \in g(A \cup B)$ .

**Exercise 1.2.8**

Here are two important definitions related to a function  $f : A \rightarrow B$ . The function  $f$  is *one-to-one* (1-1) if  $a_1 \neq a_2$  in  $A$  implies that  $f(a_1) \neq f(a_2)$  in  $B$ . The function  $f$  is *onto* if, given any  $b \in B$ , it is possible to find an element  $a \in A$  for which  $f(a) = b$ . Give an example of each or state that the request is impossible:

(a)  $f : \mathbb{N} \rightarrow \mathbb{N}$  that is 1-1 but not onto.

(b)  $f : \mathbb{N} \rightarrow \mathbb{N}$  that is onto but not 1-1.

(c)  $f : \mathbb{N} \rightarrow \mathbb{Z}$  that is 1-1 and onto.

**SOLUTION**

(a) Let  $f(x) = x + 1$ , which is 1-1 but does not have a solution to  $f(x) = 1$ , hence not onto.

(b) Let  $f(x) = 1$  for  $x = 1$  and  $f(x) = x - 1$  for  $x > 1$ , which is onto but not 1-1 as  $f(1) = f(2) = 1$ .

(c) Let  $f(x) = n/2$  when  $n$  is even and  $f(x) = -\frac{x-1}{2}$  when  $n$  is odd.

**Exercise 1.2.9**

Given a function  $f : D \rightarrow \mathbb{R}$  and a subset  $B \subseteq \mathbb{R}$ , let  $f^{-1}(B)$  be the set of all points from the domain  $D$  that get mapped into  $B$ ; that is  $f^{-1}(B) = \{x \in D : f(x) \in B\}$ . This set is called the *preimage* of  $B$ .

(a) Let  $f(x) = x^2$ . If  $A$  is the closed interval  $[0, 4]$  and  $B$  is the closed interval  $[-1, 1]$ , find  $f^{-1}(A)$  and  $f^{-1}(B)$ . Does  $f^{-1}(A \cap B) = f^{-1}(A) \cap f^{-1}(B)$  in this case? Does  $f^{-1}(A \cup B) = f^{-1}(A) \cup f^{-1}(B)$ ?

(b) The good behaviour of preimages demonstrated in (a) is completely general. Show that for an arbitrary function  $g : \mathbb{R} \rightarrow \mathbb{R}$ , it is always true that  $g^{-1}(A \cap B) = g^{-1}(A) \cap g^{-1}(B)$  and  $g^{-1}(A \cup B) = g^{-1}(A) \cup g^{-1}(B)$  for all sets  $A, B \subseteq \mathbb{R}$ .

**SOLUTION**

(a) For  $f(x) = x^2$ ,  $f^{-1}(A) = [-2, 2]$  and  $f^{-1}(B) = [-1, 1]$ .  $f^{-1}(A \cap B) = f^{-1}([0, 1]) = [-1, 1] = f^{-1}(A) \cap f^{-1}(B)$ . Similarly,  $f^{-1}(A \cup B) = f^{-1}([-1, 4]) = [-2, 2] = f^{-1}(A) \cup f^{-1}(B)$ .

(b) To show that  $g^{-1}(A \cap B) = g^{-1}(A) \cap g^{-1}(B)$ , we have to show inclusion both ways:

(i) Let  $x \in g^{-1}(A \cap B)$ , so  $g(x) \in A \cap B$ , which implies  $g(x) \in A$  and  $g(x) \in B$ . This shows that  $x \in g^{-1}(A)$  and  $x \in g^{-1}(B)$ , hence  $x \in g^{-1}(A) \cap g^{-1}(B)$ .

(ii) Let  $x \in g^{-1}(A) \cap g^{-1}(B)$ , so  $x \in g^{-1}(A)$  and  $x \in g^{-1}(B)$ , then  $g(x) \in A$  and  $g(x) \in B$ . This implies that  $g(x) \in A \cap B$ , so  $x \in g^{-1}(A \cap B)$ .

Showing  $g^{-1}(A \cup B) = g^{-1}(A) \cup g^{-1}(B)$  is obvious using Exercise 1.2.7 (d).

**Exercise 1.2.10**

Decide which of the following are true statements. Provide a short justification for those that are valid and a counterexample for those that are not:

(a) Two real numbers satisfy  $a < b$  if and only if  $a < b + \epsilon$  for every  $\epsilon > 0$ .

(b) Two real numbers satisfy  $a < b$  if  $a < b + \epsilon$  for every  $\epsilon > 0$ .

(c) Two real numbers satisfy  $a \leq b$  if and only if  $a < b + \epsilon$  for every  $\epsilon > 0$ .

**SOLUTION**

(a) False. Consider the case where  $a < b + \epsilon$  is true but  $a = b$ .

(b) False. Same reasoning as above.

(c) True. Firstly suppose  $a < b + \epsilon$  for all  $\epsilon > 0$ . We need to show this implies  $a \leq b$ . We either have  $a \leq b$  or  $a > b$ . However,  $a > b$  is not possible as this implies there exists an  $\epsilon$  small enough such that  $a > b + \epsilon$ . Secondly, suppose  $a \leq b$ . It is obvious that  $a < b + \epsilon$  for all  $\epsilon > 0$ .



**Exercise 1.2.11**

Form the logical negation of each claim. One trivial way to do this is to simply add "It is not the case that..." in front of each assertion. To make this more interesting, fashion the negation into a positive statement that avoids using the word "not" altogether. In each case, make an intuitive guess as to whether the claim or its negation is the true statement.

- (a) For all real numbers satisfying  $a < b$ , there exists an  $n \in \mathbb{N}$  such that  $a + 1/n < b$ .
- (b) There exists a real number  $x > 0$  such that  $x < 1/n$  for all  $n \in \mathbb{N}$ .
- (c) Between every two distinct real numbers there is a rational number.

**SOLUTION**

- (a) For all  $n \in \mathbb{N}$ , there exists  $a, b \in \mathbb{R}$  such that  $a + 1/n < b$ . [FALSE]
- (b) For all real number  $x > 0$ , there exists an  $n \in \mathbb{N}$  such that  $x \geq 1/n$ . [TRUE]
- (c) There exists two real numbers  $a < b$  such that if  $r < b$  then  $r < a$  for all  $r \in \mathbb{Q}$ . [FALSE]

**Exercise 1.2.12**

Let  $y_1 = 6$ , and for each  $n \in \mathbb{N}$  define  $y_{n+1} = (2y_n - 6)/3$ .

- (a) Use induction to prove that the sequence satisfies  $y_n > -6$  for all  $n \in \mathbb{N}$ .
- (b) Use another induction argument to show the sequence  $(y_1, y_2, y_3, \dots)$  is decreasing.

**SOLUTION**

- (a) For  $n = 1$ ,  $y_1 = 6 > -6$  (Base Case). Suppose  $y_n > -6$  for some  $n \in \mathbb{N}$ .

$$y_{n+1} = \frac{2y_n - 6}{3} > \frac{2(-6) - 6}{3} = -6$$

Hence, by induction,  $y_n > -6$  for all  $n \in \mathbb{N}$ .

- (b) Suppose  $y_{n+1} < y_n$ . The base case works as  $y_2 = 2 < 6 = y_1$ . Then,

$$\begin{aligned} y_{n+1} < y_n &\implies 2y_{n+1} - 6 < 2y_n - 6 \\ &\implies \frac{2y_{n+1} - 6}{3} < \frac{2y_n - 6}{3} \\ &\implies y_{n+2} < y_{n+1} \end{aligned}$$

Thus,  $y_{n+1} < y_n$  is true for all  $n \in \mathbb{N}$ .

**Exercise 1.2.13**

For this exercise, assume Exercise 1.2.5 has been successfully completed.

- (a) Show how induction can be used to conclude that

$$(A_1 \cup A_2 \cup \dots \cup A_n)^c = A_1^c \cap A_2^c \cap \dots \cap A_n^c$$

for any finite  $n \in \mathbb{N}$ .

- (b) It is tempting to appeal to induction to conclude

$$\left( \bigcup_{i=1}^{\infty} A_i \right)^c = \bigcap_{i=1}^{\infty} A_i^c$$

but induction does not apply here. Induction is used to prove that a particular statement holds for every value of  $n \in \mathbb{N}$ , but this does not imply the validity of the infinite case. To illustrate this point, find an example of a collection of sets  $B_1, B_2, B_3, \dots$  where  $\bigcap_{i=1}^n B_i \neq \emptyset$  is true for every  $n \in \mathbb{N}$ , but  $\bigcap_{i=1}^{\infty} B_i \neq \emptyset$  fails.

- (c) Nevertheless, the infinite version of De Morgan's Law stated in (b) is a valid statement. Provide a proof that does not use induction.

**SOLUTION**

- (a) Using Exercise 1.2.5 as the base case. Suppose  $(A_1 \cup \dots \cup A_n)^c = A_1^c \cap \dots \cap A_n^c$  is true. Using associative law on the  $n + 1$ c case,

$$\begin{aligned} ((A_1 \cup \dots \cup A_n) \cup A_{n+1})^c &= (A_1 \cup \dots \cup A_n)^c \cap A_{n+1}^c \\ &= (A_1^c \cap \dots \cap A_n^c) \cap A_{n+1}^c \\ &= A_1^c \cap \dots \cap A_n^c \cap A_{n+1}^c \end{aligned}$$

- (b) Consider  $B_1 = \{1, 2, \dots\}$ ,  $B_2 = \{2, 3, \dots\}$ ,  $B_n = \{x : x \in \mathbb{N} \cap [n, \infty)\}$ .
- (c) To prove the infinite version of De Morgan's Law, we need to show inclusion both ways.
- (i) Suppose  $x \in (\bigcup_{i=1}^{\infty} A_i)^c$ , then  $x \notin \bigcup_{i=1}^{\infty} A_i$ . This implies that  $x \notin A_i \ \forall i \in \mathbb{N}$ , so  $x \in A_i^c \ \forall i \in \mathbb{N}$ . Hence  $x \in \bigcap_{i=1}^{\infty} A_i^c$ .
- (ii) Suppose  $x \in \bigcap_{i=1}^{\infty} A_i^c$ , then  $x \in A_i^c \ \forall i \in \mathbb{N}$ , so  $x \notin A_i \ \forall i \in \mathbb{N}$ . This implies that  $x \notin \bigcup_{i=1}^{\infty} A_i$ , hence  $x \in (\bigcup_{i=1}^{\infty} A_i)^c$ .

## 1.3 The Axiom of Completeness

**Exercise 1.3.1**

- (a) Write a formal definition in the style of Definition 1.3.2 for the *infimum* or *greatest lower bound* of a set.
- (b) Now, state and prove a version of Lemma 1.3.8 for greatest lower bound.

**SOLUTION**

- (a) A real number  $i$  is the *greatest upper bound* for a set  $A \subseteq \mathbb{R}$  if it meets the following two criteria:
- (i)  $i$  is a lower bound of  $A$ ;
- (ii) if  $b$  is any lower bound for  $A$ , then  $i \geq b$ .
- (b) Assume  $i \in \mathbb{R}$  is a lower bound for a set  $A \subseteq \mathbb{R}$ . Then  $i = \inf A$  if and only if, for every choice of  $\epsilon > 0$ , there  $\exists a \in A$  satisfying  $i + \epsilon > a$ .

PROOF Rephrasing the lemma gives us: Given that  $i$  is a lower bound,  $i$  is the greatest lower bound if and only if any number greater than  $i$  is not a lower bound.

- (i) Suppose  $i = \inf A$  and consider  $i + \epsilon$  for an arbitrarily chosen  $\epsilon > 0$ . Since  $i + \epsilon > i$ , part (ii) of the definition implies that  $i + \epsilon$  is not a lower bound for  $A$ . If this is the case, then there must be some element  $a \in A$  such that  $i + \epsilon > a$ .
- (ii) Conversely, assume  $i$  is a lower bound with the property that for any  $\epsilon > 0$ ,  $i + \epsilon$  is not a lower bound of  $A$ . Note that this implies that if  $b$  is any number more than  $i$ , then  $b$  is not an upperbound. Since we have argued that any larger number than  $i$  cannot be a lower bound, if  $b$  is some other upper bound for  $A$ , then  $i \geq b$ .

**Exercise 1.3.2**

Give an example of each of the following, or state that the request is impossible.

- (a) A set  $B$  with  $\inf B \geq \sup B$ .
- (b) A finite set that contains its infimum but not its supremum.
- (c) A bounded subset of  $\mathbb{Q}$  that contains its supremum but not its infimum.

**SOLUTION**

- (a) Possible. Consider the set  $0$ , where  $\inf\{0\} = \sup\{0\} = 0$ .
- (b) Not possible as all finite sets must contain its supremum and infimum.
- (c) Possible. Consider  $A = \{r \in \mathbb{Q} \mid 1 < r \leq 2\}$ .

**Exercise 1.3.3**

- (a) Let  $A$  be nonempty and bounded below, and define  $B = \{b \in \mathbb{R} : b \text{ is a lower bound for } A\}$ . Show that  $\sup B = \inf A$ .
- (b) Use (a) to explain why there is no need to assert that greatest lower bounds exist as part of the Axiom of Completeness.

**SOLUTION**

- (a) By definition,  $\sup B$  is the greatest lower bound for  $A$ , meaning it equals  $\inf A$ .
- (b) Part (a) proves the greatest lower bound exists using the least upper bound.

**Exercise 1.3.4**

Let  $A_1, A_2, A_3, \dots$  be a collection of nonempty sets, each of which is bounded above.

- (a) Find a formula for  $\sup(A_1 \cup A_2)$ . Extend this to  $\sup(\bigcup_{k=1}^n A_k)$ .
- (b) Consider  $\sup(\bigcup_{k=1}^{\infty} A_k)$ . Does the formula in (a) extend to the infinite case?

**SOLUTION**

- (a)  $\sup(A_1 \cup A_2) = \sup\{\sup A_1, \sup A_2\}$  and  $\sup(\bigcup_{k=1}^n A_k) = \sup\{\sup A_k \mid k = 1, \dots, n\}$
- (b) This formula does not extend to infinity. Consider  $A_k = [k, k+1]$ , where  $\bigcup_{k=1}^{\infty} A_k$  is unbounded.

**Exercise 1.3.5**

As in Example 1.3.7, let  $A \subseteq \mathbb{R}$  be nonempty and bounded above, and let  $c \in \mathbb{R}$ . This time define the set  $cA = \{ca : a \in A\}$ .

- (a) If  $c \geq 0$ , show that  $\sup(cA) = c \sup A$ .
- (b) Postulate a similar type of statement for  $\sup(cA)$  for the case  $c < 0$ .

**SOLUTION**

- (a) The case of  $c = 0$  is trivial as it implies that  $cA = \{0\}$ . Hence  $\sup(cA) = c \sup A = 0$ .  
For  $c > 0$ , we need to show that  $c \sup A$  is the lowest upper bound. Assume  $c > 0$ . Let  $s = c \sup A$ . Suppose  $ca > s$ , then  $a > \sup A$  which is impossible, meaning that  $s$  is an upper bound on  $cA$ . Now suppose  $s'$  is an upper bound on  $cA$  and  $s' < s$ . Then  $s'/c < s/c = \sup A$ , meaning  $s'/c$  cannot bound  $A$ . Hence there  $\exists a \in A$  such that  $s'/c < a$ , meaning  $s' > ca$ , thus  $s'$  cannot be an upper bound on  $cA$ , so  $s = c \sup A$  is the least upper bound.
- (b)  $\sup(cA) = c \inf A$  for  $c < 0$

**Exercise 1.3.6**

Given sets  $A$  and  $B$ , define  $A + B = \{a + b : a \in A \text{ and } b \in B\}$ . Follow these steps to prove that if  $A$  and  $B$  are nonempty and bounded above then  $\sup(A + B) = \sup A + \sup B$ .

- (a) Let  $s = \sup A$  and  $t = \sup B$ . Show  $s + t$  is an upper bound for  $A + B$ .
- (b) Now let  $u$  be an arbitrary upper bound for  $A + B$ , and temporarily fix  $a \in A$ . Show  $t \leq u - a$ .
- (c) Finally, show  $\sup(A + B) = s + t$ .
- (d) Construct another proof of this same fact using Lemma 1.3.8.

**SOLUTION**

- (a) By definition of supremum,  $a \leq s$  and  $b \leq t$ . Adding both equations give  $a + b \leq s + t$ , hence  $s + t$  is an upper bound.
- (b) Since  $a + b \leq u$  implies  $b \leq u - a$ ,  $u - a$  is an upper bound on  $b$ , meaning it is greater or equal to the least upper bound of  $t$ , giving  $t \leq u - a$ .
- (c) From (a), we have shown that  $s + t$  is an upper bound for  $A + B$ , hence it is sufficient to show that  $s + t$  is the least upper bound.
- Let  $u = \sup(A + B)$ , from (b) we have  $t \leq u - a$  and  $s \leq u - b$ . Adding and rearranging gives  $a + b \leq 2u - s - t$ . Since  $2u - s - t$  is an upper bound on  $A + B$ , it must be greater or equal to the least upper bound, giving  $u \leq 2u - s - t$ , implying  $s + t \leq u$ . Since  $u$  is the least upper bound,  $s + t$  must equal  $u$ .
- (d) Showing  $s + t - \epsilon$  is not an upper bound for any  $\epsilon > 0$  proves that it is the least upper bound by Lemma 1.3.8. Rearranging gives  $(s - \epsilon/2) + (t - \epsilon/2)$  we know  $\exists a > (s - \epsilon/2)$  and  $b > (t - \epsilon/2)$ , therefore  $a + b > s + t - \epsilon$ , meaning  $s + t$  cannot be made smaller and thus is the least upper bound.

**Exercise 1.3.7**

Prove that if  $a$  is an upper bound for  $A$ , and if  $a$  is also an element of  $A$ , then it must be that  $a = \sup A$ .

**SOLUTION**

Since it is given that  $a$  is an upper bound for  $A$ , we just have to show that  $a$  is the least upper bound, meaning any number lower than  $a$  would have an  $a' \in A$  such that  $a' > a$ .

Suppose  $a - \epsilon$  is also an upper bound for  $A$  for some  $\epsilon > 0$ . This is not possible has  $a > a'$  and  $a \in A$ . Hence by contradiction,  $a$  is the lowest upper bound, meaning  $a = \sup A$ .

**Exercise 1.3.8**

Compute, without proofs, the suprema and infima (if they exists) of the following sets:

- (a)  $\{m/n : m, n \in \mathbb{N} \text{ with } m < n\}$ .
- (b)  $\{(-1)^m/n : m, n \in \mathbb{N}\}$ .
- (c)  $\{n/(3n + 1) : n \in \mathbb{N}\}$ .
- (d)  $\{m/(m + n) : m, n \in \mathbb{N}\}$ .

**SOLUTION**

- (a)  $\inf = 0$  and  $\sup = 1$
- (b)  $\inf = -1$  and  $\sup = 1$
- (c)  $\inf = 1/4$  and  $\sup = 1/3$
- (d)  $\inf = 0$  and  $\sup = 1$

**Exercise 1.3.9**

- (a) If  $\sup A < \sup B$ , show there exists an element that is an upper bound for  $A$ .
- (b) Give an example to show that this is not always the case if we only assume  $\sup A \leq \sup B$ .

**SOLUTION**

- (a) By Lemma 1.3.8, we know there exists a  $b$  such that  $(\sup B) - \epsilon < b$  for any  $\epsilon > 0$ . We can set  $\epsilon$  to be small enough such that  $\sup B - \sup A < \epsilon$ , implying  $\sup A < \sup B - \epsilon < b$  for some  $b \in B$ , thus  $b$  is an upper bound of  $A$ .
- (b) Consider the sets  $A = (-\infty, 1]$  and  $B = (-\infty, 1)$ . No  $b \in B$  is an upperbound since  $1 \in A$  and  $1 > b$ .

**Exercise 1.3.10**

**(Cut Property)** The *Cut Property* of the real numbers is the following:

If  $A$  and  $B$  are nonempty, disjoint sets with  $A \cup B = \mathbb{R}$  and  $a < b$  for all  $a \in A$  and  $b \in B$ , then there exists a  $c \in \mathbb{R}$  such that  $x \leq c$  whenever  $x \in A$  and  $x \geq c$  whenever  $x \in B$ .

- (a) Use the Axiom of Completeness to prove the Cut Property.
- (b) Show that the implication goes the other way; that is, assume  $\mathbb{R}$  possesses the Cut Property and let  $E$  be a nonempty set that is bounded above. Prove  $\sup E$  exists.
- (c) The punchline of parts (a) and (b) is that the Cut Property could be used in place of the Axiom of Completeness as the fundamental axiom that distinguishes the real numbers from the rational numbers. To drive this point home, give a concrete example showing that the Cut Property is not a valid statement when  $\mathbb{R}$  is replaced by  $\mathbb{Q}$ .

**SOLUTION**

- (a) If  $c = \sup A = \inf B$ , it is obvious that  $a \leq c \leq b$  and we are done. Hence we need to show, by contradiction, that  $\sup A < \inf B$  and  $\sup A > \inf B$  is false.
  - (i) AFSOC  $\sup A < \inf B$ . We can choose  $c = \frac{\sup A + \inf B}{2}$ , which satisfies  $\sup A < c < \inf B$ . Hence it is obvious that  $c \notin A$  and  $c \notin B$ , so  $c \notin A \cup B \neq \mathbb{R}$  which is a contradiction.
  - (ii) AFSOC  $\sup A > \inf B$ . We can find  $a$  such that  $a > b$  by subtracting  $\epsilon > 0$  and using the definition of supremum and infimum similar to Lemma 1.3.8. Thus creating a contradiction.

Since both alternatives are impossible,  $\sup A = \inf B$ .

- (b) If  $E$  is finite or has a maximum element, that is  $\sup E$  and we are done.

Consider the case where  $E$  has not maximum element (for example,  $\{-1/n : n \in \mathbb{N}\}$ ). Let  $B$  be the sets of all upper bounds of  $E$  and let  $A = B^c$ . It can be said  $E \cap B = \emptyset$  otherwise  $E$  has a maximum element. Thus  $E \subseteq A$ .

By the Cut Property, there exists  $c$  such that  $a \leq c \leq b$  for all  $a \in A$  and  $b \in B$ . Since  $c$  is an upper bound on  $A$  and  $E \subseteq A$ ,  $c$  is also an upper bound on  $E$ . And since  $c \leq b$ ,  $c$  is the lowest upper bound. Therefore,  $c = \sup E$ .

- (c) Consider  $A = \{r \in \mathbb{Q} \mid r^2 < 2 \text{ or } r < 0\}$ ,  $B = A^c$  does not satisfy the cut property in  $\mathbb{Q}$  as  $\sqrt{2} \notin \mathbb{Q}$ .

**Exercise 1.3.11**

Decide if the following statements about suprema and infima are true or false. Give a short proof for those that are true. For any that are false, supply an example where the claim in question does not appear to hold.

- (a) If  $A$  and  $B$  are nonempty, bounded and satisfy  $A \subseteq B$ , then  $\sup A \leq \sup B$ .
- (b) If  $\sup A < \inf B$  for sets  $A$  and  $B$ , then there exists a  $c \in \mathbb{R}$  satisfying  $a < c < b$  for all  $a \in A$  and  $b \in B$ .
- (c) If there exists a  $c \in \mathbb{R}$  satisfying  $a < c < b$  for all  $a \in A$  and  $b \in B$ , then  $\sup A < \inf B$ .

**SOLUTION**

- (a) True. We know  $a \leq \sup A$ ,  $a \leq \sup B$  since  $A \subseteq B$ . Since  $\sup A$  is the least upper bound on  $A$ , we have  $\sup A \leq \sup B$ .
- (b) True. Let  $c = \frac{\sup A + \inf B}{2}$ ,  $c > \sup A$  implies  $a < c$  and  $c < \inf B$  implies  $c < b$ , giving  $a < c < b$ .
- (c) False. Consider  $A = (-\infty, 1)$  and  $B = (1, \infty)$ ,  $a < 1 < b$  but  $\sup A \not< \inf B$  as  $\sup A = \inf B = 1$ .

## 1.4 Consequences of Completeness

### Exercise 1.4.1

Recall that  $\mathbb{I}$  stands for the set of irrational numbers.

- (a) Show that if  $a, b \in \mathbb{Q}$ , then  $ab$  and  $a + b$  are elements of  $\mathbb{Q}$  as well.
- (b) Show that if  $a \in \mathbb{Q}$  and  $t \in \mathbb{I}$ , then  $a + t \in \mathbb{I}$  and  $at \in \mathbb{I}$  as long as  $a \neq 0$ .
- (c) Part (a) can be summarized by saying that  $\mathbb{Q}$  is closed under addition and multiplication. Is  $\mathbb{I}$  closed under addition and multiplication? Given two irrational numbers  $s$  and  $t$ , what can we say about  $s + t$  and  $st$ ?

### SOLUTION

- (a) This is trivial. Since  $a, b \in \mathbb{Q}$ , they can be expressed as a fraction of two integers. Let  $a = m/n$  and  $b = x/y$  where  $m, n, x, y \in \mathbb{Z}$ , then  $a + b = \frac{my + xn}{ny}$  and  $ab = \frac{mx}{ny}$ , which are fractions with integer numerators and denominators, hence  $a + b$  and  $ab$  are elements of  $\mathbb{Q}$ .
- (b) AFSOC that  $a + t \in \mathbb{Q}$  and  $at \in \mathbb{Q}$ . Let  $a + t = \alpha$  and  $at = \beta$ , so  $t = \alpha - a$  and  $t = \beta/a$ . Since  $\alpha, \beta, -a, 1/a \in \mathbb{Q}$ , using part (a) gives  $t \in \mathbb{Q}$  which is a contradiction. Hence  $a + t \in \mathbb{I}$  and  $at \in \mathbb{I}$ .
- (c)  $\mathbb{I}$  is not closed under addition or multiplication. Consider  $\sqrt{2} + (-\sqrt{2}) = 0$  and  $\sqrt{2} \cdot \sqrt{2} = 2$ .

### Exercise 1.4.2

Let  $A \subseteq \mathbb{R}$  be nonempty and bounded above, and let  $s \in \mathbb{R}$  have the property that for all  $n \in \mathbb{N}$ ,  $s + \frac{1}{n}$  is an upper bound for  $A$  and  $s - \frac{1}{n}$  is not an upper bound for  $A$ . Show  $s = \sup A$ .

### SOLUTION

We can rephrase Lemma 1.3.8 using the archimedean property.

- (i) AFSOC  $s < \sup A$ , then there exists an  $n$  such that  $s + 1/n < \sup A$ , contradicting  $\sup A$  being the least upper bound.
- (ii) AFSOC  $s > \sup A$ , then there exists an  $n$  such that  $s - 1/n > \sup A$  where  $s - 1/n$  is not an upper bound, contradicting  $\sup A$  being an upper bound.

Hence, we can conclude that  $\sup A = s$ .

### Exercise 1.4.3

Prove that  $\bigcap_{n=1}^{\infty} (0, 1/n) = \emptyset$ . Notice that this demonstrates that the intervals in the Nested Interval Property must be closed for the conclusion of the theorem to hold.

### SOLUTION

AFSOC the  $x \in \bigcap_{n=1}^{\infty} (0, 1/n)$ , so  $0 < x < 1/n$  for all  $n \in \mathbb{N}$  which is impossible by archimedean property.

### Exercise 1.4.4

Let  $a < b$  be real numbers and consider the set  $T = \mathbb{Q} \cap [a, b]$ . Show  $\sup T = b$ .

### SOLUTION

To show that  $\sup T = b$ , it needs to satisfy both conditions of supremum:

- (i) Since  $x \leq b$  for all  $x \in [a, b]$ ,  $y \leq b$  for all  $y \in T$  as  $T \subseteq [a, b]$ .
- (ii) AFSOC  $b'$  is also an upper bound such that  $b' < b$ . Since  $\mathbb{Q}$  is dense in  $\mathbb{R}$ , there exists an  $\alpha \in \mathbb{Q} \cap [b', b] \subseteq T$ . This implies there exists  $t \in T$  satisfying  $b' < t$ , which is a contradiction.

### Exercise 1.4.5

Using Exercise 1.4.1, supply a proof for Corollary 1.4.4 by considering real numbers  $a - \sqrt{2}$  and  $b - \sqrt{2}$ .

### SOLUTION

Recall that **Corollary 1.4.4** states that *Given any two real numbers  $a < b$ , there exists an irrational number  $t$  satisfying  $a < t < b$ .*

Since  $\mathbb{Q}$  is dense in  $\mathbb{R}$ , we can find  $t \in \mathbb{Q}$  that is between any two real numbers  $a - \sqrt{2}$  and  $b - \sqrt{2}$ , with  $a < b$ . Hence,  $a - \sqrt{2} < t < b - \sqrt{2}$ , meaning  $a < t + \sqrt{2} < b$ . By Exercise 1.4.1,  $t + \sqrt{2} \in \mathbb{I}$  and we are done.

**Exercise 1.4.6**

Recall that a set  $B$  is *dense* in  $\mathbb{R}$  if can element  $B$  can be found between any two real numbers  $a < b$ . Which of the following sets are dense in  $\mathbb{R}$ ? Take  $p \in \mathbb{Z}$  and  $q \in \mathbb{N}$  in every case.

- (a) The set of all rational numbers  $p/q$  with  $q \leq 10$ .
- (b) The set of all rational numbers  $p/q$  with  $q$  a power of 2.
- (c) The set of all rational numbers  $p/q$  with  $10|p| \geq q$ .

**SOLUTION**

- (a) Not dense since we cannot make  $|p|/q < 1/10$ .
- (b) Dense.
- (c) Not dense since we cannot make  $|p|/q < 1/10$ .

**Exercise 1.4.7**

Finish the proof of Theorem 1.4.5 by showing that the assumption  $\alpha^2 > 2$  leads to a contradiction of the fact that  $\alpha = \sup T$ .

**SOLUTION**

Recall  $T = \{t \in \mathbb{R} : t^2 < 2\}$  and  $\alpha = \sup T$ . AFSOC  $\alpha^2 > 2$ , we will show that there exists an  $n \in \mathbb{N}$  such that  $(\alpha - 1/n)^2 > 2$ , contradicting the assumption that  $\alpha$  is the least upper bound.

Using  $(\alpha - 1/n)^2 > 2$ , we can find  $n \in \mathbb{N}$  such that  $(\alpha^2 - 1/n) > 2$ .

$$2 < (\alpha - 1/n)^2 = \alpha^2 - \frac{2\alpha}{n} + 1/n^2 < \alpha^2 - \frac{2\alpha - 1}{n}$$

Then

$$2 < \alpha^2 - \frac{2\alpha - 1}{n} \implies n(2 - \alpha^2) < 1 - 2\alpha$$

Since  $2 - \alpha^2 < 0$ , dividing reverses the inequality, giving

$$n > \frac{1 - 2\alpha}{2 - \alpha^2}$$

Hence we can pick  $n \in \mathbb{N}$  such that  $(\alpha^2 - 1/n) > 2$ , so  $\alpha$  is the the least upper bound which is a contradiction. Hence it is not possible for  $\alpha^2 > 2$ .

**Exercise 1.4.8**

Give an example of each or state that the request is impossible. When a request is impossible, provide a compelling argument for why this is the case.

- (a) Two sets  $A$  and  $B$  with  $A \cap B \neq \emptyset$ ,  $\sup A = \sup B$ ,  $\sup A \notin A$  and  $\sup B \notin B$ .
- (b) A sequence of nested open intervals  $J_1 \supseteq J_2 \supseteq J_3 \supseteq \dots$  with  $\bigcap_{n=1}^{\infty} J_n$  nonempty but containing only a finite number of elements.
- (c) A sequence of nested unbounded closed intervals  $L_1 \supseteq L_2 \supseteq L_3 \supseteq \dots$  with  $\bigcap_{n=1}^{\infty} L_n = \emptyset$ . (An unbounded closed interval has the form  $[a, \infty) = \{x \in \mathbb{R} : x \geq a\}$ .)
- (d) A sequence of closed bounded (not necessarily nested) intervals  $I_1, I_2, I_3, \dots$  with the property that  $\bigcap_{n=1}^N I_n \neq \emptyset$  for all  $N \in \mathbb{N}$ , but  $\bigcap_{n=1}^{\infty} I_n = \emptyset$ .

**SOLUTION**

- (a) Consider  $A = (0, 1) \cap \mathbb{Q}$  and  $B = (0, 1) \cap \mathbb{I}$ .  $\sup A = \sup B = 1$  and  $1 \in A, B$ .
- (b) Consider  $J_n = (-1/n, 1/n)$ , then  $\bigcap_{n=1}^{\infty} J_n = \{0\}$  which is nonempty and contains a finite number of elements.
- (c) Consider  $L_n = [n, \infty)$ , then  $\bigcap_{n=1}^{\infty} L_n = \emptyset$ .
- (d) Impossible. Let  $j_n = \bigcap_{k=1}^n I_k$ . Since  $\bigcap_{n=1}^N I_n \neq \emptyset$ , we have  $J_n \neq \emptyset$ . Note that  $J_n$  is the intersection of closed intervals, making it a closed interval. With  $I_{n+1} \cap J_n \subseteq J_n$ , we have  $J_{n+1} \subseteq J_n$ . Since  $J_n$  is a series of nested bounded closed intervals, by Nested Interval Property,  $\bigcap_{n=1}^{\infty} J_n = \bigcap_{n=1}^{\infty} (\bigcap_{k=1}^n I_k) = \bigcap_{n=1}^{\infty} I_n \neq \emptyset$ .

## 1.5 Cardinality

### Exercise 1.5.1

Finish the following proof for Theorem 1.5.7.

Assume  $B$  is a countable set. Thus, there exists  $f : \mathbb{N} \rightarrow B$ , which is 1-1 and onto. Let  $A \subseteq B$  be an infinite subset of  $B$ . We must show that  $A$  is countable.

Let  $n_1 = \min\{n \in \mathbb{N} : f(n) \in A\}$ . As a start to a definition of  $g : \mathbb{N} \rightarrow A$ , set  $g(1) = f(n_1)$ . Show how to inductively continue this process to produce a 1-1 function  $g$  from  $\mathbb{N}$  onto  $A$ .

### SOLUTION

Let  $n_k = \min\{n \in \mathbb{N} : f(n) \in A, n \notin \{n_1, n_2, \dots, n_{k-1}\}\}$  and  $g(k) = f(n_k)$ . Since  $g : \mathbb{N} \rightarrow A$  is 1-1 and onto,  $A$  is countable.

### Exercise 1.5.2

Review the proof of Theorem 1.5.6, part (ii) showing that  $\mathbb{R}$  is uncountable, and then find the flaw in the following erroneous proof that  $\mathbb{Q}$  is uncountable:

Assume, for contradiction, that  $\mathbb{Q}$  is countable. Thus we can write  $\mathbb{Q} = \{r_1, r_2, r_3, \dots\}$  and, as before, construct a nested sequence of closed intervals with  $r_n \notin I_n$ . Our construction implies  $\bigcap_{n=1}^{\infty} I_n = \emptyset$  while NIP implies  $\bigcap_{n=1}^{\infty} I_n \neq \emptyset$ . This contradiction implies  $\mathbb{Q}$  must therefore be uncountable.

### SOLUTION

The Nested Interval Property only applies to  $\mathbb{R}$  and not  $\mathbb{Q}$ .

### Exercise 1.5.3

Use the following proofs for the statements in Theorem 1.5.8.

- (a) First, prove statement (i) for two countable sets,  $A_1$  and  $A_2$ . Example 1.5.3 (ii) may be a useful reference. Some technicalities can be avoided by first replacing  $A_2$  with the set  $B_2 = A_2 \setminus A_1 = \{x \in A_2 : x \notin A_1\}$ . The point of this is that the union  $A_1 \cup B_2$  is equal to  $A_1 \cup A_2$  and the sets  $A_1$  and  $B_2$  are disjoint. (What happens if  $B_2$  is finite?)

Now, explain how that more general statement in (i) follows.

- (b) Explain why induction cannot be used to prove part (ii) of Theorem 1.5.8 from part (i).
- (c) Show how arranging  $\mathbb{N}$  into the two-dimensional array

$$\begin{array}{cccccc} 1 & 3 & 6 & 10 & 15 & \dots \\ 2 & 5 & 9 & 14 & \dots & \\ 4 & 8 & 13 & \dots & & \\ 7 & 12 & \dots & & & \\ \vdots & & & & & \end{array}$$

leads to a proof of Theorem 1.5.8 (ii).

### SOLUTION

- (a) Let  $B = \{b_1, b_2, b_3, \dots\}$  and  $C = \{c_1, c_2, c_3, \dots\}$  be countable, disjoint sets. We can define a function  $g : \mathbb{N} \rightarrow B \cup C$  following a similar method of mapping  $\mathbb{N}$  onto  $\mathbb{Z}$ , listing them as follows:

$$B \cup C = \{b_1, c_1, b_2, c_2, \dots\}$$

Implying that  $B \cup C$  is countable. By letting  $B = A_1$  and  $C = A_2 \setminus A_1$ , we can show that  $A_1 \cup A_2$  is also countable.

By using induction, suppose  $A_1 \cup \dots \cup A_n$  is countable,  $(A_1 \cup \dots \cup A_n) \cup A_{n+1}$  is the unions of two countable sets, which as proven above is also countable.

- (b) Induction only shows something for each  $n \in \mathbb{N}$ , it does not apply in the infinite case. For example,  $\bigcap_{k=1}^n [k, \infty) \neq \emptyset$  is true for all  $n \in \mathbb{N}$ , but the infinite case  $\bigcap_{k=1}^{\infty} [k, \infty) \neq \emptyset$  is false.



- (c) By rearranging  $\mathbb{N}$  as in (c) gives us disjoint sets  $C_n = \{k \in \mathbb{N} : k \text{ is in the } n\text{-th row}\}$ , such that  $\bigcup_{n=1}^{\infty} C_n = \mathbb{N}$ . Let  $B_n$  be disjoint sets, constructed as  $B_1 = A_1$ ,  $B_2 = A_2 \setminus A_1$ ,  $\dots$ ,  $B_n = A_n \setminus (A_1 \cup \dots \cup A_{n-1})$ . If we can find a function  $f : C \rightarrow B$  that is bijective, we can show:

$$f(\mathbb{N}) = f\left(\bigcup_{n=1}^{\infty} C_n\right) = \bigcup_{n=1}^{\infty} f(C_n) = \bigcup_{n=1}^{\infty} B_n = \bigcup_{n=1}^{\infty} A_n$$

Let  $f_n : C_n \rightarrow B_n$  be bijective since  $B_n$  is countable. Define  $f : \mathbb{N} \rightarrow \bigcup_{n=1}^{\infty} B_n$  as

$$f(n) = \begin{cases} f_1(n) & \text{if } n \in C_1 \\ f_2(n) & \text{if } n \in C_2 \\ \vdots & \end{cases}$$

We must now show that  $f$  is bijective:

- (i) Since each  $C_n$  is disjoint and each  $f_n$  is 1-1,  $f(n_1) = f(n_2) \implies n_1 = n_2$ , meaning  $f$  is 1-1.
- (ii) Since any  $b \in \bigcup_{n=1}^{\infty} B_n$  as  $b \in B_n$  for some  $n$ ,  $b = f_n(x)$  has a solution since  $f_n$  is onto. Letting  $x = f_n^{-1}(b)$ , we have  $f(x) = f_n(x) = b$  as  $f_n^{-1}(b) \in C_n$ , meaning  $f$  is onto.

By (i) and (ii),  $f$  is bijective, so  $\bigcup_{n=1}^{\infty} B_n$  is countable, implying that  $\bigcup_{n=1}^{\infty} A_n$  is also countable, completing the proof.

#### Exercise 1.5.4

- (a) Show  $(a, b) \sim \mathbb{R}$  for any interval  $(a, b)$ .
- (b) Show that an unbounded interval like  $(a, \infty) = \{x : x > a\}$  has the same cardinality as  $\mathbb{R}$  as well.
- (c) Using open intervals makes it more convenient to produce the required 1-1, onto functions, but it is not really necessary. Show that  $[0, 1) \sim (0, 1)$  by exhibiting a 1-1 onto function between the two sets.

#### SOLUTION

- (a) Example 1.5.4 provides the function  $f(x) = \frac{x}{x^2-1}$  which takes the interval  $(-1, 1)$  onto  $\mathbb{R}$  in a 1-1 fashion. If we shift  $f$  to the interval of  $(a, b)$ , we get:

$$g(x) = f\left(\frac{2x-1}{b-a} - a\right)$$

which maps  $(a, b)$  onto  $\mathbb{R}$  in a 1-1 fashion.

- (b) To show that  $(a, \infty) \sim \mathbb{R}$ , we need to find a function  $h$  that maps  $\mathbb{R}$  onto  $(a, \infty)$ . Consider the function  $h(x) = e^x + a$ , which maps  $\mathbb{R}$  onto  $(a, \infty)$  in a 1-1 fashion. Hence we are done.
- (c) Define  $f : [0, 1) \rightarrow (0, 1)$  as

$$f(x) = \begin{cases} 1/2 & \text{if } x = 0 \\ 1/4 & \text{if } x = 1/2 \\ 1/8 & \text{if } x = 1/4 \\ \vdots & \\ x & \text{otherwise} \end{cases}$$

To show that both  $[0, 1)$  and  $(0, 1)$  have the both cardinality, we need to prove that  $f$  is 1-1 and onto.

We start by showing that  $y = f(x)$  has exactly one solution for all  $y \in (0, 1)$ .

If  $y = 1/2^n$ , then the only solution is  $y = f(1/2^{n-1})$  (or  $x = 0$  in the special case  $n = 1$ ).

Otherwise, the only solution is  $y = f(y)$ .

#### Exercise 1.5.5

- (a) Why is  $A \sim A$  for every set  $A$ ?

- (b) Given sets  $A$  and  $B$ , explain why  $A \sim B$  is equivalent to asserting  $B \sim A$ .
- (c) For three sets  $A$ ,  $B$ , and  $C$ , show that  $A \sim B$  and  $B \sim C$  implies  $A \sim C$ . These three properties are what is meant by saying that  $\sim$  is an *equivalence relation*.

**SOLUTION**

- (a) The identity function  $f(x) = x$  is trivially bijective.
- (b) If  $f : A \rightarrow B$  is bijective, then the inverse  $f^{-1} : B \rightarrow A$  is also bijective.
- (c) If  $f : A \rightarrow B$  and  $g : B \rightarrow C$ , then  $g \circ f : A \rightarrow C$  is a bijection, thus  $A \sim C$ .

**Exercise 1.5.6**

- (a) Give an example of a countable collection of disjoint open intervals.
- (b) Give an example of an uncountable collection of disjoint open intervals, or argue that no such collection exists.

**SOLUTION**

- (a) Consider  $I_1 = (1, 2)$ ,  $I_2 = (2, 3)$ ,  $\dots$ ,  $I_n = (n, n + 1)$ .
- (b) No such collection exists. Let  $A$  denote this set.

For any nonempty interval  $I_n$ , since  $\mathbb{Q}$  is dense in  $\mathbb{R}$ , we can find an  $r \in \mathbb{Q}$  such that  $r \in I_n$ . Assigning each  $I \in A$  a rational number  $r \in I$  proves that  $I \subseteq \mathbb{Q}$ , thus  $I$  is countable.

**Exercise 1.5.7**

Consider the open interval  $(0, 1)$ , and let  $S$  be the set of points in the open unit square; that is,  $S = \{(x, y) : 0 < x, y < 1\}$ .

- (a) Find a 1-1 function that maps  $(0, 1)$  into, but not necessarily onto,  $S$ . (This is easy)
- (b) Use the fact that every real number has a decimal expansion to produce a 1-1 function that maps  $S$  into  $(0, 1)$ . Discuss whether the formulated function is onto. (Keep in mind that any terminating decimal expansion such as .235 represents the same real number as .234999...)

The Schröder-Bernstein Theorem discussed in Exercise 1.5.11 can now be applied to conclude that  $(0, 1) \sim S$ .

**SOLUTION**

- (a) Consider  $f(x) = (\frac{1}{4}x, \frac{1}{3})$ .
- (b) Let  $g : S \rightarrow (0, 1)$  be a function that interweaves decimals in the representation without trailing nines, padding with zeros if necessary.  $g(0.12, 0.34) = 0.1324$ ,  $g(0.1\bar{9}, 0.2) = g(0.2, 0.2) = 0.22$ ,  $g(0.4, 0.89) = 0.4809$ ,  $g(0.6, 0.\bar{7}) = 0.670\bar{7}$ , etc.

To prove that  $g$  is 1-1, we need show that there does not exist any solutions for  $g(x_1, y_1) = g(x_2, y_2)$ . Every real number can be written with two representations, one with trailing 9's and one without. However,  $g(x, y) = 0.d_1d_2d_3\dots\bar{9}$  is impossible as it would imply both  $x$  and  $y$  has trailing 9's, which contradicts the definition of  $g$ . Therefore,  $g(s)$  is unique and so  $g$  is 1-1.

$g$  is not onto since  $g(x, y) = 0.1$  has no solutions.

**Exercise 1.5.8**

Let  $B$  be a set of positive real numbers with the property that adding together any finite subset of elements from  $B$  will always give a sum of 2 or less. Show  $B$  must be finite or countable.

**SOLUTION**

It is obvious that  $B \cap [a, 2)$  must be finite for some  $a \in (0, 2)$ , otherwise, we can choose  $\lceil \frac{2}{a} \rceil$  number of elements from  $B \cap [a, 2)$  would result in a sum of more than 2. Since  $B$  is the countable union of finite sets  $\bigcup_{n=1}^{\infty} B \cap [\frac{1}{n}, 2)$ ,  $B$  must be countable or finite.

**Exercise 1.5.9**

A real number  $x \in \mathbb{R}$  is called *algebraic* if there exist integers  $a_0, a_1, a_2, \dots, a_n \in \mathbb{Z}$ , not all zero, such that

$$a_n x^n + a_{n-1} x^{n-1} + \dots + a_1 x + a_0 = 0.$$

Said another way, a real number is algebraic if it is the root of a polynomial with integer coefficients. Real numbers that are not algebraic are called *transcendental* numbers. Reread the last paragraph of Section 1.1. The final question posed here is closely related to the question of whether or not transcendental numbers exist.

- (a) Show that  $\sqrt{2}$ ,  $\sqrt[3]{2}$  and  $\sqrt{3} + \sqrt{2}$  are algebraic.
- (b) Fix  $n \in \mathbb{N}$  and let  $A_n$  be the algebraic numbers obtained as roots of polynomials with integer coefficients that have degree  $n$ . Using the fact that every polynomial has a finite number of roots, show that  $A_n$  is countable.
- (c) Now, argue that the set of all algebraic numbers is countable. What may we conclude about the set of transcendental numbers?

**SOLUTION**

- (a) The first two are trivial as they are roots of  $x^2 - 2 = 0$  and  $x^3 - 2 = 0$ . For  $\sqrt{3} + \sqrt{2}$ , we can start working backwards:

$$\begin{aligned} x = \sqrt{3} + \sqrt{2} &\implies x^2 = 5 + 2\sqrt{6} \\ &\implies (x^2 - 5)^2 - 24 = 0, \end{aligned}$$

thus giving a polynomial, hence  $\sqrt{2} + \sqrt{3}$  is algebraic.

- (b) We need to show that  $A_n \sim \mathbb{Z}^n \sim \mathbb{N}^n \sim \mathbb{N}$  and hence countable.
  - (i)  $A_n \sim \mathbb{Z}^n$  since choosing  $n$  integer coefficients for a polynomial of degree  $n$  is equivalent to an ordered list of  $n$  integers.
  - (ii)  $\mathbb{Z}^n \sim \mathbb{N}^n$  since  $f : \mathbb{N}^n \rightarrow \mathbb{Z}^n$  is just the piecewise application of  $g : \mathbb{N} \rightarrow \mathbb{Z}$ .
  - (iii)  $\mathbb{N}^n \sim \mathbb{N}$  since it is the intersection of finite sets  $\bigcup_{k=1}^{\infty} \{(x_1, \dots, x_n) : x_1 + \dots + x_n = k\}$ .
- (c) The set of algebraic numbers is countable as it is the union of countable sets  $\bigcup_{n=1}^{\infty} A_n$ . The set of transcendental numbers is uncountable, otherwise, it implies that the set of real numbers is countable, which is not possible.

**Exercise 1.5.10**

- (a) Let  $C \subseteq [0, 1]$  be uncountable. Show that there exists  $a \in (0, 1)$  such that  $C \cap [a, 1]$  is countable.
- (b) Now let  $A$  be the set of all  $a \in (0, 1)$  such that  $C \cap [a, 1]$  is uncountable, and set  $\alpha = \sup A$ . Is  $C \cap [\alpha, 1]$  an uncountable set?
- (c) Does the statement in (a) remain true if "uncountable" is replaced by "infinite"?

**SOLUTION**

- (a) AFSOC  $a$  does not exist, so  $C \cap [a, 1]$  is countable for all  $a \in (0, 1)$ . Therefore  $C = \bigcup_{n=1}^{\infty} C \cap [1/n, 1]$  is also countable, which is a contradiction. Hence, there exist a  $a \in (0, 1)$  such that  $C \cap [a, 1]$  is uncountable.
- (b) Since  $\alpha = \sup A$ , for any  $\epsilon > 0$ ,  $C \cap [\alpha + \epsilon, 1]$  is countable. Therefore,  $C \cap [\alpha, 1] = \alpha \cup \bigcup_{n=1}^{\infty} C \cap [1/n, 1]$  is countable.
- (c) No. Consider  $C = \{1/n : n \in \mathbb{N}\}$ . It has  $C \cap [\alpha, 1]$  finite for every  $\alpha$ , but  $C \cap [0, 1]$  is infinite.

**Exercise 1.5.11**

**(Schröder-Bernstein Theorem)** Assume there exists a 1-1 function  $f : X \rightarrow Y$  and another 1-1 function  $g : Y \rightarrow X$ . Follow the steps to show that there exists a 1-1, onto function  $h : X \rightarrow Y$  and hence  $X \sim Y$ .

The strategy is to partition  $X$  and  $Y$  into components

$$X = A \cup A' \quad \text{and} \quad Y = B \cup B'$$

with  $A \cap A' = \emptyset$  and  $B \cap B' = \emptyset$ , in such a way that  $f$  maps  $A$  onto  $B$ , and  $g$  maps  $B'$  and  $A'$ .

- (a) Explain how achieving this would lead to a proof that  $X \sim Y$ .
- (b) Set  $A_1 = X \setminus g(Y) = \{x \in X : x \notin g(Y)\}$  (what happens if  $A_1 = \emptyset$ ?) and inductively define a sequence of sets by letting  $A_{n+1} = g(f(A_n))$ . Show that  $\{A_n : n \in \mathbb{N}\}$  is a pairwise disjoint collection of subsets of  $X$ , while  $\{f(A_n) : n \in \mathbb{N}\}$  is a similar collection in  $Y$ .
- (c) Let  $A = \bigcup_{n=1}^{\infty} A_n$  and  $B = \bigcup_{n=1}^{\infty} f(A_n)$ . Show that  $f$  maps  $A$  onto  $B$ .
- (d) Let  $A' = X \setminus A$  and  $B' = Y \setminus B$ . Show  $g$  maps  $B'$  onto  $A'$ .

**SOLUTION**

- (a)  $f : A \rightarrow B$  and  $g : B' \rightarrow A'$  are bijective, therefore we can define

$$h(x) = \begin{cases} f(x) & \text{if } x \in A \\ g^{-1}(x) & \text{if } x \in A' \end{cases}$$

which is bijective.

- (b) If  $A_1 = \emptyset$ , we are done as it implies  $g(Y) = X$  so  $X \sim Y$

Since  $A_n \subseteq X$ , then  $f(A_n) \subseteq Y$ , hence  $A_n \cap A_1 = \emptyset$  for all  $n \in \mathbb{N}$  as  $A_1 = X \setminus g(Y)$ .

Note that  $g \circ f$  is bijective implies that  $g(f(A \cap B)) = g(f(A)) \cap g(f(B))$ . We will show that by inclusion both ways. Let  $h$  be an arbitrary bijective function.

- (i) Let  $y \in h(A \cap B)$ , so there exists  $x \in A \cap B$  such that  $h(x) = y$ , then  $x \in A$  and  $x \in B$ . This implies that  $h(x) \in h(A)$  and  $h(x) \in h(B)$ , hence  $y = h(x) \in h(A) \cap h(B)$ .
- (ii) Let  $y \in h(A) \cap h(B)$ , so  $y \in h(A)$  and  $y \in h(B)$ , then there exist  $x \in A$  and  $x \in B$  such that  $h(x) = y$ . This implies  $x \in A \cap B$ , hence  $y \in h(A \cap B)$ .

Now to show pairwise disjointment for all subsets, consider  $i, j \in \mathbb{N}$ , with  $i < j$ .

$$A_j \cap A_i = g(f(A_j \cap A_i)) = g(f(A_j)) \cap g(f(A_i)) = A_{j-1} \cap A_{i-1} = \cdots = A_{j-i+1} \cap A_1 = \emptyset$$

Similarly,  $\{f(A_n) : n \in \mathbb{N}\}$  is also pairwise disjoint as

$$A_j \cap A_i = \emptyset \implies f(A_j) \cap f(A_i) = f(A_j \cap A_i) = f(\emptyset) = \emptyset$$

- (c)  $f(A) = B$  as  $f(\bigcup_{n=1}^{\infty} A_n) = \bigcup_{n=1}^{\infty} f(A_n) = B$ . ( $B$  is defined to be the range of  $f$ .)
- (d) To show that  $g(B') = A'$ , we will show inclusion both ways:
  - (i)  $g(B') \subseteq A'$ . AFSOC  $g(b') \in A$ , then since  $A_1 \cap g(Y) = \emptyset$ ,  $g(b') \notin A$ , implying  $g(b') \in \bigcup_{n=2}^{\infty} A_n = g(B)$ , meaning  $\exists b \in B$  such that  $g(b) = g(b')$ , contradicting the fact that  $g$  is 1-1.
  - (ii)  $g(B') \supseteq A'$ . AFSOC  $\exists a' \in A'$ , with  $a' \notin g(B')$ . Since  $A' \subseteq g(Y)$ , we have  $a' \in g(B)$ , since  $a' \notin g(B')$ , and  $g(B) \subseteq A$  contradicting  $a' \in A$ .

## 1.6 Cantor's Theorem

### Exercise 1.6.1

Show that  $(0, 1)$  is uncountable if and only if  $\mathbb{R}$  is uncountable.

#### SOLUTION

In Exercise 1.5.4 (a), we have found a bijection  $f : (0, 1) \rightarrow \mathbb{R}$ . Suppose  $g : (0, 1) \rightarrow \mathbb{N}$  is some map, we must show  $g$  is bijective if and only if  $(g \circ f) : \mathbb{R} \rightarrow \mathbb{N}$  is bijective. This is true as if  $g$  is bijective, then  $(g \circ f)$  is bijective as it is the composition of bijective functions. Similarly, if  $(g \circ f)$  is bijective, then  $(g \circ f) \circ f^{-1} = g$  is bijective.

This proof works as  $(0, 1)$  is uncountable, implies that  $g$  is not bijective, then  $(g \circ f) : \mathbb{R} \rightarrow \mathbb{N}$  is not bijective, implying that  $\mathbb{R}$  is not countable and hence uncountable.

### Exercise 1.6.2

- (a) Explain why the real number  $x = .b_1b_2b_3b_4\dots$  cannot be  $f(1)$ .
- (b) Now, explain why  $x \neq f(2)$ , and in general why  $x \neq f(n)$  for any  $n \in \mathbb{N}$ .
- (c) Point out the contradiction that arises from these observations and conclude that  $(0, 1)$  is uncountable.

#### SOLUTION

- (a) Since  $b_1 \neq a_{11}$ , the first digit of  $f(1)$  differ from  $x$ , hence cannot be equal.
- (b) Since  $b_n \neq a_{nn}$ , the  $n$ -th digit of  $f(n)$  differ from  $x$ , hence cannot be equal.
- (c) Since  $x$  is not in the list, it is a contradiction.

### Exercise 1.6.3

Supply rebuttals to the following complaints about the proof of Theorem 1.6.1.

- (a) Every rational number has a decimal expansion, so we could apply this same argument to show that the set of rational numbers between 0 and 1 is uncountable. However, because we know that any subset of  $\mathbb{Q}$  must be countable, the proof of Theorem 1.6.1 must be flawed.
- (b) Some numbers have *two* different decimal representations. Specifically, any decimal expansion that terminates can also be written with repeating 9's. For instance,  $1/2$  can be written as  $.5$  or as  $.4999\dots$ . Doesn't this cause some problems?

#### SOLUTION

- (a) False, since the constructed number has an infinite number of decimals, it is irrational.
- (b) No, since changing the  $n$ -th digit would still result in a different number.

### Exercise 1.6.4

Let  $S$  be the set consisting of all sequences of 0's and 1's. Observe that  $S$  is not a particular sequence, but rather a large set whose elements are sequences; namely,

$$S = \{(a_1, a_2, a_3, \dots) : a_n = 0 \text{ or } 1\}.$$

As an example, the sequence  $(1, 0, 1, 0, 1, 0, 1, 0, \dots)$  is an element of  $S$ , as is the sequence  $(1, 1, 1, 1, 1, 1, \dots)$ .

Give a rigorous argument showing  $S$  is uncountable.

#### SOLUTION

Similar to the proof in Exercise 1.6.1, suppose there exists a function  $f : \mathbb{N} \rightarrow S$  that is 1-1 and onto. For each  $n \in \mathbb{N}$ ,  $f(n)$  is an element in  $S$ , represented as:

$$f(n) = (a_{n1}, a_{n2}, a_{n3}, \dots)$$

where  $a_{nm} = 0$  or  $1$  for each  $m, n \in \mathbb{N}$ . We construct a  $s \in S$ , with  $s = (b_1, b_2, b_3, \dots)$ , where  $b_n = 0$  if  $a_{nn} = 1$  and  $b_n = 1$  otherwise. Since the  $n$ -th digit of sequence  $s$  differs from  $f(n)$  for all  $n \in \mathbb{N}$ ,  $s \notin S$ , which is a contradiction. Hence  $S$  is not countable, therefore uncountable.

Alternatively, we can define  $g : S \rightarrow \bar{S}$  where  $g(a_1, a_2, a_3, \dots) = .a_1a_2a_3\dots$  which is trivially bijective, and  $h : \mathbb{R}_2 \rightarrow \mathbb{R}$  which converts a number in base 2 to base 10 and is clearly bijective. Hence  $h \circ g : S \rightarrow \mathbb{R}$  is bijective and hence shows that  $S \sim \mathbb{R}$  so  $S$  is uncountable.

**Exercise 1.6.5**

- (a) Let  $A = \{a, b, c\}$ . List the eight elements of  $P(A)$ . (Do not forget that  $\emptyset$  is considered to be a subset of every set.)
- (b) If  $A$  is finite with  $n$  elements, show that  $P(A)$  has  $2^n$  elements.

**SOLUTION**

- (a)  $P(A) = \{\emptyset, \{a\}, \{b\}, \{c\}, \{a, b\}, \{a, c\}, \{b, c\}, \{a, b, c\}\}$
- (b) There are  $n$  elements, with 2 possible states: included or excluded, implying that there are  $2^n$  elements.

**Exercise 1.6.6**

- (a) Using the particular set  $A = \{a, b, c\}$ , exhibit two different 1-1 mappings from  $A$  into  $P(A)$ .
- (b) Letting  $C = \{1, 2, 3, 4\}$ , produce an example of a 1-1 map  $g : C \rightarrow P(C)$ .
- (c) Explain *why*, in parts (a) and (b), it is impossible to construct mappings that are *onto*.

**SOLUTION**

- (a) Consider  $f(x) = \{x\}$  and  $f(x) = \{x\}$  for  $x \neq b$ ,  $\{x, c\}$  for  $x = b$ .
- (b) Let  $g(x) = \{x\}$ .
- (c)  $|P(C)| > |C|$ .

**Exercise 1.6.7**

Return to the particular functions constructed in Exercise 1.6.6 and construct the subset  $B$  that results using the preceding rule. In each case, note that  $B$  is not in the range of the function used.

**SOLUTION**

- (i) For  $A = \{a, b, c\}$ ,  $B_A = \{b\}$
- (ii) For  $C = \{1, 2, 3, 4\}$ ,  $B_C = \emptyset$

**Exercise 1.6.8**

- (a) First, show that the case  $a' \in B$  leads to a contradiction.
- (b) Now, finish the argument by showing that the case  $a' \notin B$  is equally unacceptable.

**SOLUTION**

- (a) AFSOC  $a' \in B$ . This means  $a' \notin f(a')$ , then  $a' \notin B$  which is a contradiction.
- (b) AFSOC  $a' \notin B$ . This means  $a' \in f(a')$ , then  $a' \in B$ , which is a contradiction.

**Exercise 1.6.9**

Using the various tools and techniques developed in the last two sections (including the exercises from Section 1.5), give a compelling argument showing that  $P(\mathbb{N}) \sim \mathbb{R}$ .

**SOLUTION**

We can make a function  $f : P(\mathbb{N}) \rightarrow S$ , with  $S$  following the definition given in Exercise 1.6.4, being a set consisting of all sequences of 0's and 1's.

$$f(P_N) = (a_1, a_2, a_3, \dots) \text{ with } a_n = 0 \text{ if } a_n \notin P_N \text{ and } 1 \text{ otherwise}$$

$f$  is clearly onto as every element of  $S$  can be mapped to by its definition. To show that it is injective, AFSOC that there exists  $X, Y \in P(\mathbb{N})$ , where  $X \neq Y$ , such that  $f(X) = f(Y)$ . There will exist  $n \in \mathbb{N}$  where  $n \in X$  but  $n \notin Y$ . By definition, the  $n$ -th element of  $f(X)$  is 1, but for  $f(Y)$  is 0, which is a contradiction.

Using Exercise 1.6.4,  $S \sim \mathbb{R}$ , hence  $P(\mathbb{N}) \sim \mathbb{R}$ .

**Exercise 1.6.10**

As a final exercise, answer each of the following by establishing a 1-1 correspondence with a set of known cardinality.

- (a) Is the set of all functions from  $\{0, 1\}$  to  $\mathbb{N}$  countable or uncountable?
- (b) Is the set of all functions from  $\mathbb{N}$  to  $\{0, 1\}$  countable or uncountable?
- (c) Given a set  $B$ , a subset  $\mathcal{A}$  of  $P(B)$  is called an *antichain* if no element of  $\mathcal{A}$  is a subset of any other element of  $\mathcal{A}$ . Does  $P(\mathbb{N})$  contain an uncountable antichain?

**SOLUTION**

- (a) Both elements of  $\{0, 1\}$  can be mapped to an element in  $\mathbb{N}$ , hence we can find a bijective function to show that this set is the same as  $\mathbb{N}^2$  hence countable.
- (b) As shown in Exercise 1.6.9, this set maps to  $S \sim \mathbb{R}$  which is uncountable.
- (c) Let  $\mathcal{A}$  be an antichain of  $P(\mathbb{N})$  and let  $\mathcal{A}_l$  be the sets in  $\mathcal{A}$  of size  $l$ . For finite  $l$ ,  $\mathcal{A}_l$  is countable since  $\mathcal{A}_l \subseteq \mathbb{N}^l$  is countable. Hence the countable union  $\bigcup_{l=0}^{\infty} \mathcal{A}_l = \mathcal{A}$  is countable.

If  $l$  is infinite, only countably many elements of  $\mathcal{A}$  can be finite, while there will be uncountably many that must be infinite, hence forming an uncountable antichain  $\mathcal{A}$ .





## Chapter 2

# Sequences and Series

## 2.2 The Limit of a Sequence

### Exercise 2.2.1

What happens if we reverse the order of the quantifiers in Definition 2.2.3?

*Definition:* A sequence  $(x_n)$  *verconges* to  $x$  if *there exists* an  $\epsilon > 0$  such that *for all*  $N \in \mathbb{N}$  it is true that  $n \geq N$  implies  $|x_n - x| < \epsilon$ .

Give an example of a vercongent sequence. Is there an example of a vercongent dequence that is divergent? Can a sequence verconge to two different values? What exactly is being described in this strange definition?

### SOLUTION

- (i) Consider the sequence  $x_n = \sin x$ .  $(x_n)$  verconges to 0 as we can choose  $\epsilon = 1$  such that  $|x_n - 0| < \epsilon$  for any value of  $n \in \mathbb{N}$ .
- (ii) There is not vercongent sequence that diverges. The proof is obvious and left as an exercise for the reader.
- (iii) The sequence in (i) verconges to  $x \in \mathbb{R}$  by choosing  $\epsilon = x + 2$ .
- (iv) The definition of verconges described a bounded sequence.

### Exercise 2.2.2

Verify, using the definition of convergence of a sequence, that the following sequences converge to the proposed limit.

(a)  $\lim_{n \rightarrow \infty} \frac{2n+1}{5n+4} = \frac{2}{5}$ .

(b)  $\lim_{n \rightarrow \infty} \frac{2n^2}{n^3+3} = 0$ .

(c)  $\lim_{n \rightarrow \infty} \frac{\sin(n^2)}{\sqrt[3]{n}}$ .

### SOLUTION

- (a) For  $\epsilon > 0$ , we can choose  $N = 1/\epsilon$ , such that for all  $n > N$ ,

$$\left| \frac{2n+1}{5n+4} - \frac{2}{5} \right| = \left| \frac{-3}{5(5n+4)} \right| = \frac{3}{5(5n+4)} < \frac{1}{n} < \epsilon \implies n > \frac{1}{\epsilon}$$

- (b) For  $\epsilon > 0$ , we can choose  $N = 2/\epsilon$ , such that for all  $n > N$ ,

$$\left| \frac{2n^2}{n^3+3} - 0 \right| = \frac{2n^2}{n^3+3} < \frac{2n^2}{n^3} = \frac{2}{n} < \epsilon \implies n > \frac{2}{\epsilon}$$

- (c) For  $\epsilon > 0$ , we can choose  $N = 1/\epsilon^3$ , such that for all  $n > N$ ,

$$\left| \frac{\sin(n^2)}{\sqrt[3]{n}} - 0 \right| = \frac{\sin(n^2)}{n^{1/3}} \leq \frac{1}{n^{1/3}} \implies n > \frac{1}{\epsilon^3}$$

### Exercise 2.2.3

Describe what we would have to demonstrate in order to disprove each of the following statements.

- (a) At every college in the United States, there is a student who is at least seven feet tall.
- (b) For all colleges in the United States, there exists a professor who gives every student a grade of either A or B.
- (c) There exists a college in the United States where every student is at least six feet tall.

### SOLUTION

- (a) Find a college in the United States that has every student under seven feet.
- (b) Find a college in the United States that have every professor giving at least one grade lower than a B.
- (c) Find a student in each college in the United States that is under six feet

### Exercise 2.2.4

Give an example of each or state that the request is impossible. For any that are impossible, give a compelling argument for why that is the case.

- (a) A sequence with an infinite number of ones that does not converge to one.
- (b) A sequence with an infinite number of ones that converges to a limit not equal to one.
- (c) A divergent sequence such that for every  $n \in \mathbb{N}$  it is possible to find  $n$  consecutive ones somewhere in the sequence.

### SOLUTION

- (a) Consider  $x_n = (-1)^n$ .
- (b) Impossible. Suppose  $\lim x_n = a \neq 1$ . Since there are infinite ones, we cannot find  $N > 0$  for  $\epsilon < |1 - a|$ .
- (c) Consider the sequence  $(1, 2, 1, 1, 3, 1, 1, 1, \dots)$ .

### Exercise 2.2.5

Let  $\llbracket x \rrbracket$  be the greatest integer less than or equal to  $x$ . For example,  $\llbracket \pi \rrbracket = 3$  and  $\llbracket 3 \rrbracket = 3$ . For each sequence, find  $\lim a_n$  and verify it with the definition of convergence.

- (a)  $a_n = \llbracket 5/n \rrbracket$ .
- (b)  $a_n = \llbracket (12 + 4n)/3n \rrbracket$ .

Reflecting on these examples, comment on the statement following the Definition 2.2.3 that "the smaller the  $\epsilon$ -neighborhood, the larger  $N$  may have to be."

### SOLUTION

- (a) For all  $n > 5$ , we have  $\llbracket 5/n \rrbracket = 0$ , hence  $\lim a_n = 0$ .
- (b) The inside converges to  $4/3$  from above, so  $\lim a_n = 1$ .  
Some sequences eventually reach their limit, meaning that  $N$  no longer has to increase.

### Exercise 2.2.6

Prove Theorem 2.2.7. To get started assume  $(a_n) \rightarrow a$  and also that  $(a_n) \rightarrow b$ . Now argue  $a = b$ .

**SOLUTION**

**Theorem 2.2.7 (Uniqueness of Limits).** The limit of a sequence, when it exists, must be unique.

AFSOC that  $\lim a_n = a$  and  $\lim a_n = b$ , such that  $a \neq b$ . Then for  $\epsilon > 0$ , choosing  $n > N_a$  implies  $|a_n - a| < \epsilon/2$  and choosing  $n > N_b$  implies  $|a_n - b| < \epsilon/2$ . Let  $N = \max(N_a, N_b)$ , hence  $|a_n - a| < \epsilon/2$  and  $|a_n - b| < \epsilon/2$ . By triangle inequality,

$$\epsilon = \epsilon/2 + \epsilon/2 > |a_n - a| + |a_n - b| = |a - a_n| + |a_n - b| \leq |a - b|$$

showing that  $a = \lim a = b$ , which is a contradiction.

**Exercise 2.2.7**

Here are two useful definitions:

- (i) A sequence  $(a_n)$  is *eventually* in a set  $A \subseteq \mathbb{R}$  if there exists an  $N \in \mathbb{N}$  such that  $a_n \in A$  for all  $n \geq N$ .
- (ii) A sequence  $(a_n)$  is *frequently* in a set  $A \subseteq \mathbb{R}$  if, for every  $N \in \mathbb{N}$ , there exists an  $n \geq N$  such that  $a_n \in A$ .
  - (a) Is the sequence  $(-1)^n$  eventually or frequently in the set  $\{1\}$ ?
  - (b) Which definition is stronger? Does frequently imply eventually or does eventually imply frequently?
  - (c) Give an alternate rephrasing of Definition 2.2.3B using either frequently or eventually. Which is the term we want?
  - (d) Suppose an infinite number of terms of a sequence  $(x_n)$  are equal to 2. Is  $(x_n)$  necessarily eventually in the interval  $(1.9, 2.1)$ ? Is it frequently in  $(1.9, 2.1)$ ?

**SOLUTION**

- (a)  $(-1)^n$  is frequently but not eventually in  $\{1\}$ .
- (b) Eventually implies frequently, but the converse is not true.
- (c)  $(x_n) \rightarrow x$  if and only if  $x_n$  is eventually in any  $\epsilon$ -neighborhood around  $x$ .
- (d)  $(x_n)$  is frequently in  $(1.9, 2.1)$  but not always eventually (considering  $x_n = 2(-1)^n$ ).

**Exercise 2.2.8**

For some additional practice with nested quantifiers, consider the following invented definition:

Let's call a sequence  $(x_n)$  *zero-heavy* if there exists  $M \in \mathbb{N}$  such that for all  $N \in \mathbb{N}$  there exists  $n$  satisfying  $N \leq n \leq N + M$  where  $x_n = 0$ .

- (a) Is the sequence  $(0, 1, 0, 1, 0, 1, \dots)$  zero heavy?
- (b) If a sequence is zero-heavy does it necessarily contain an infinite number of zeros? If not, provide a counterexample.
- (c) If a sequence contains an infinite number of zeros, is it necessarily zero-heavy? If not, provide a counterexample.
- (d) From the logical negation of the above definition. That is, complete the sentence: A sequence is *not zero-heavy* if  $\dots$

**SOLUTION**

- (a) The sequence is zero-heavy, choosing  $M = 1$ .
- (b) Yes. If there were a finite number of zeros, with the last zero at  $x_K$ , then choosing  $N > K$  would lead to a contradiction.
- (c) No, consider  $(0, 1, 0, 1, 1, 0, 1, 1, 1, 0, 1, 1, 1, 0, \dots)$  where the gap between the 0's grows to infinity.
- (d) A sequence is not zero-heavy if for all  $M \in \mathbb{N}$ , there exists some  $N \in \mathbb{N}$  such that for all  $n \in \mathbb{N}$ ,  $N \leq n \leq N + M$ ,  $x_n \neq 0$ .

## 2.3 The Algebraic and Order Limit Theorem

### Exercise 2.3.1

Let  $x_n \geq 0$  for all  $n \in \mathbb{N}$ .

- (a) If  $(x_n) \rightarrow 0$ , show that  $(\sqrt{x_n}) \rightarrow 0$ .
- (b) If  $(x_n) \rightarrow x$ , show that  $(\sqrt{x_n}) \rightarrow \sqrt{x}$ .

### SOLUTION

- (a) For  $\epsilon^2 > 0$ , there exists  $N > 0$ , such that for all  $n > N$ ,  $x_n < \epsilon^2$ , implying  $\sqrt{x_n} < \epsilon$ .
- (b) We want to show that  $|\sqrt{x_n} - \sqrt{x}| < \epsilon$ . Multiplying by  $(\sqrt{x_n} + \sqrt{x})$  gives us  $|x_n - x| < (\sqrt{x_n} + \sqrt{x})\epsilon$ . Since  $(x_n)$  converges, it is bounded, giving  $|x_n| \leq M$ , then  $\sqrt{x_n} \leq \sqrt{M}$ .

$$|x_n - x| < (\sqrt{x_n} + \sqrt{x})\epsilon \leq (M + \sqrt{x})\epsilon$$

Setting  $|x_n - x| < (M + \sqrt{x})\epsilon$  for  $n > N$ , we get

$$|\sqrt{x_n} - \sqrt{x}| = \frac{|x_n - x|}{\sqrt{x_n} + \sqrt{x}} \leq \frac{|x_n - x|}{M + \sqrt{x}} = \epsilon$$

thus completing the proof.

### Exercise 2.3.2

Using only Definition 2.2.3, prove that if  $(x_n) \rightarrow 2$ , then

- (a)  $(\frac{2x_n-1}{3}) \rightarrow 1$ ;
- (b)  $(1/x_n) \rightarrow 1/2$ ;

(For this exercise the Algebraic Limit Theorem is off-limits, so to speak.)

### SOLUTION

- (a) We have  $|\frac{2x_n-1}{3} - 1| = |\frac{2}{3}x_n - \frac{4}{3}| = \frac{2}{3}|x_n - 2|$ . Setting  $|x_n - 2| < \frac{3}{2}\epsilon$  for all  $n > N$ , we get  $|\frac{2x_n-1}{3} - 1| < \epsilon$ .
- (b) We set  $N$  such that  $|x_n - 2| < \epsilon$ . Since  $x_n$  is at least 1, we can bound  $|1/x_n| \leq 1$ , giving

$$|1/x_n - 1/2| = \frac{|2 - x_n|}{|2x_n|} \leq \frac{|x_n - 2|}{2} \leq \frac{\epsilon}{2} < \epsilon.$$

### Exercise 2.3.3

**(Squeeze Theorem)** Show that if  $x_n \leq y_n \leq z_n$  for all  $n \in \mathbb{N}$ , and if  $\lim x_n = \lim z_n = l$ , then  $\lim y_n = l$  as well.

### SOLUTION

Let  $\epsilon > 0$ , set  $N > 0$  so that  $|x_n - l| < \epsilon/3$  and  $|z_n - l| < \epsilon/3$ . By triangle inequality, we get

$$|z_n - x_n| \leq |z_n - l| + |l - x_n| < 2\epsilon/3$$

Since  $z_n \geq y_n \geq x_n$ ,  $|y_n - x_n| = y_n - x_n \leq z_n - x_n$ . So by triangle inequality,

$$|y_n - l| \leq |y_n - x_n| + |x_n - l| \leq |z_n - x_n| + |x_n - l| < 2\epsilon/3 + \epsilon/3 = \epsilon$$

### Exercise 2.3.4

Let  $(a_n) \rightarrow 0$ , and use the Algebraic Limit Theorem to compute each of the following limits (assuming the fractions are always defined):

- (a)  $\lim \left( \frac{1+2a_n}{1+3a_n-4a_n^2} \right)$
- (b)  $\lim \left( \frac{(a_n+2)^2-4}{a_n} \right)$

$$(c) \lim \left( \frac{\frac{2}{a_n} + 3}{\frac{1}{a_n} + 5} \right).$$

**SOLUTION**

(a) By ALT,

$$\begin{aligned} \lim \left( \frac{1 + 2a_n}{1 + 3a_n - 4a_n^2} \right) &= \frac{\lim(1 + 2a_n)}{\lim(1 + 3a_n - 4a_n^2)} \\ &= \frac{1 + 2 \lim a_n}{1 + 3 \lim a_n - 4(\lim a_n)^2} \\ &= 1 \end{aligned}$$

(b) By ALT,

$$\begin{aligned} \lim \left( \frac{(a_n + 2)^2 - 4}{a_n} \right) &= \lim \left( \frac{a_n^2 + 4a_n}{a_n} \right) \\ &= \lim(a_n + 4) = 4 + \lim a_n = 4 \end{aligned}$$

(c) By ALT,

$$\begin{aligned} \lim \left( \frac{\frac{2}{a_n} + 3}{\frac{1}{a_n} + 5} \right) &= \lim \left( \frac{2 + 3a_n}{1 + 5a_n} \right) \\ &= \frac{\lim(2 + 3a_n)}{\lim(1 + 5a_n)} \\ &= \frac{2 + 3 \lim a_n}{1 + 5 \lim a_n} = 2 \end{aligned}$$

**Exercise 2.3.5**

Let  $(x_n)$  and  $(y_n)$  be given, and define  $(z_n)$  to be the "shuffled" sequence  $(x_1, y_1, x_2, y_2, x_3, y_3, \dots, x_n, y_n, \dots)$ . Prove that  $(z_n)$  is convergent if and only if  $(x_n)$  and  $(y_n)$  are both convergent with  $\lim x_n = \lim y_n$ .

**SOLUTION**

Obviously, if  $\lim x_n = \lim y_n = l$ , then for  $\epsilon > 0$ , we can set  $N > 0$  such that  $|x_n - l| < \epsilon$  and  $|y_n - l| < \epsilon$  for  $n > N$ , thus for all  $n > 2N$ ,  $|z_n - l| < \epsilon$  will be true.

Conversely, if  $(z_n) \rightarrow l$ , for  $\epsilon > 0$ , we can set  $2N > 0$  such that for all  $n > 2N$ ,  $|z_n - l| < \epsilon$ , implying that for  $n > N$ ,  $|x_n - l| < \epsilon$  and  $|y_n - l| < \epsilon$  will also be true.

**Exercise 2.3.6**

Consider the sequence given by  $b_n = n - \sqrt{n^2 + 2n}$ . Taking  $(1/n) \rightarrow 0$  as given, and using both the Algebraic Limit Theorem and the result in Exercise 2.3.1, show  $\lim b_n$  exists and find the value of the limit.

**SOLUTION**

Let  $(b_n) \rightarrow b$ . To remove the radical from the term, consider:

$$(n - \sqrt{n^2 + 2n})(n + \sqrt{n^2 + 2n}) = n^2 - (n^2 + 2n) = -2n$$

By ALT,

$$b = \lim \left( \frac{-2n}{n + \sqrt{n^2 + 2n}} \right) = \lim \frac{-2}{1 + \sqrt{1 + 2/n}} = \frac{-2}{1 + \sqrt{1 + 2 \lim(1/n)}} = \frac{-2}{1 + \sqrt{1 + 0}} = -1$$

thus showing that the limit exists and  $(b_n) \rightarrow -1$ .

**Exercise 2.3.7**

Give an example of each of the following, or state that such a request is impossible by referencing the proper theorem(s):

- (a) sequences  $(x_n)$  and  $(y_n)$ , which both diverge but whose sum  $(x_n + y_n)$  converges;
- (b) sequences  $(x_n)$  and  $(y_n)$ , where  $(x_n)$  converges,  $(y_n)$  diverges, and  $(x_n + y_n)$  converges;
- (c) a convergent sequence  $(b_n)$  with  $b_n \neq 0$  for all  $n$  such that  $(1/b_n)$  diverges;
- (d) an unbounded sequence  $(a_n)$  and a convergent sequence  $(b_n)$  with  $(a_n - b_n)$  bounded;
- (e) two sequences  $(a_n)$  and  $(b_n)$ , where  $(a_nb_n)$  and  $(a_n)$  converge but  $(b_n)$  does not.

**SOLUTION**

- (a) Consider  $x_n = n$  and  $y_n = -n$  but  $x_n + y_n = 0$  for all  $n \in \mathbb{N}$ .
- (b) Impossible. By ALT,  $\lim y_n = \lim(x_n + y_n) - \lim x_n$  must converge, which is a contradiction.
- (c) Consider  $b_n = 1/n$ , then  $1/b_n = n$ , so  $(1/b_n) \rightarrow \infty$ .
- (d) Impossible. Let  $|b_n| \leq M$  and  $|a_n - b_n| \leq N$ . By triangle inequality,

$$|a_n| \leq |a_n - b_n| + |b_n| \leq M + N$$

, hence is also bounded, which is a contradiction.

- (e) Consider  $a_n = 1/n$  and  $b_n = n$ , then  $a_nb_n = 1$ .

**Exercise 2.3.8**

Let  $(x_n) \rightarrow x$  and let  $p(x)$  be a polynomial.

- (a) Show  $p(x_n) \rightarrow p(x)$ .
- (b) Find an example of a function  $f(x)$  and a convergent sequence  $(x_n) \rightarrow x$  where the sequence  $f(x_n)$  converges, but not to  $f(x)$ .

**SOLUTION**

- (a) Applying the Algebraic Limit Theorem repeatedly results in  $(x_n^d) \rightarrow x^d$ , meaning

$$\lim p(x_n) = \lim \sum_{i=0}^d k_i x_n^i = \sum_{i=0}^d k_i (\lim x_n)^i = p(\lim x_n) = p(x)$$

- (b) Consider the sequence  $(x_n)$ , where  $x_n = 1/n$ , and let  $f$  be defined as

$$f(x) = \begin{cases} 1 & \text{if } x \neq 0 \\ 0 & \text{otherwise} \end{cases}$$

where  $(x_n) \rightarrow x$  but  $f(x_n) \rightarrow 1 \neq f(x) = 0$ .

**Exercise 2.3.9**

- (a) Let  $(a_n)$  be a bounded (not necessarily convergent) sequence, and assume  $\lim b_n = 0$ . Show that  $\lim(a_nb_n) = 0$ . Why are we not allowed to use the Algebraic Limit Theorem to prove this?
- (b) Can we conclude anything about the convergence of  $(a_nb_n)$  if we assume that  $(b_n)$  converges to some nonzero limit  $b$ ?
- (c) Use (a) to prove Theorem 2.3.3, part (iii), for the case when  $a = 0$ .

**SOLUTION**

- (a) Since  $(a_n)$  is bounded, let  $|a_n| \leq M_a$ . Consider

$$|a_nb_n| = |a_n||b_n| \leq M|b_n|$$

For  $\epsilon > 0$ , we can set  $N > 0$  such that  $|b_n| < \epsilon/M$ , giving us  $|a_nb_n| < \epsilon$ , hence  $\lim(a_nb_n) = 0$ .

ALT does not hold as  $(a_n)$  does not converge.

- (b) If  $(a_n)$  converges, then  $(a_nb_n)$  converges by ALT. Otherwise,  $(a_nb_n)$  does not converge.
- (c) Shown in (a),  $\lim(a_nb_n) = 0 = ab$  for  $b = 0$ , proving part (iii) of the ALT.

**Exercise 2.3.10**

Consider the following list of conjectures. Provide a short proof for those that are true and a counterexample for any that are false.

- (a) If  $\lim(a_n - b_n) = 0$ , then  $\lim a_n = \lim b_n$ .
- (b) If  $(b_n) \rightarrow b$ , then  $|b_n| \rightarrow |b|$ .
- (c) If  $(a_n) \rightarrow a$  and  $(b_n - a_n) \rightarrow 0$ , then  $(b_n) \rightarrow a$ .
- (d) If  $(a_n) \rightarrow 0$  and  $|b_n - b| \leq a_n$  for all  $n \in \mathbb{N}$ , then  $(b_n) \rightarrow b$ .

**SOLUTION**

- (a) False. Consider  $a_n = n$  and  $b_n = -n$ , then  $\lim(a_n - b_n) = 0$  but  $\infty = \lim a_n \neq \lim b_n = -\infty$ .
- (b) True. Since  $|b_n - b| < \epsilon$ , then without loss of generality, we assume  $|b_n| > |b|$ , so

$$||b_n| - |b|| = |b_n| - |b| = |b_n - b + b| - |b| \leq |b_n - b| + |b| - |b| = |b_n - b| < \epsilon$$

- (c) True. By ALT,  $\lim b_n = \lim a_n + \lim(b_n - a_n) = a + 0 = a$ .
- (d) True. Note that  $0 \leq |b_n - b| \leq a_n$ . Since  $(a_n) \rightarrow 0$ , by Squeeze Theorem in Exercise 2.3.3,  $\lim |b_n - b| = 0$ , hence  $\lim b_n = b$ .

**Exercise 2.3.11**

(Cesaro Means).

- (a) Show that if  $(x_n)$  is a convergent sequence, then the sequence given by the averages

$$y_n = \frac{x_1 + x_2 + \cdots + x_n}{n}$$

also converges to the same limit.

- (b) Give an example to show that it is possible for the sequence  $(y_n)$  of averages to converge even if  $(x_n)$  does not.

**SOLUTION**

- (a) Let  $D = \sup\{|x_n - x| : n \in \mathbb{N}\}$  and  $0 < \epsilon < D$ , we have

$$|y_n - x| = \left| \frac{x_1 + x_2 + \cdots + x_n}{n} - \frac{xn}{n} \right| \leq \left| \frac{|x_1 - x| + |x_2 - x| + \cdots + |x_n - x|}{n} \right| \leq D$$

Set  $N_1 > 0$  such that  $|x_n - x| < \epsilon/2$  for  $n > N_1$ , giving

$$|y_n - x| = \left| \frac{|x_1 - x| + \cdots + |x_{N_1} - x| + \cdots + |x_n - x|}{n} \right| \leq \left| \frac{N_1 D + (n - N_1)\epsilon/2}{n} \right|$$

Set  $N_2 > 0$  such that for all  $n > N_2$ ,

$$0 < \frac{N_1(D - \epsilon/2)}{n} < \epsilon/2$$

Therefore

$$|y_n - x| \leq \left| \frac{N_1(D - \epsilon/2)}{n} + \epsilon/2 \right| \leq \left| \frac{N_1(D - \epsilon/2)}{n} \right| + \epsilon/2 < \epsilon$$

Setting  $N = \max\{N_1, N_2\}$  completes the proof.

- (b)  $x_n = (-1)^n$  diverges but  $(y_n) \rightarrow 0$ .

### Exercise 2.3.12

A typical task in analysis is to decipher whether a property possessed by every term in a convergent sequence is necessarily inherited by the limit. Assume  $(a_n) \rightarrow a$ , and determine the validity of each claim. Try to produce a counterexample for any that are false.

- (a) If every  $a_n$  is an upper bound for a set  $B$ , then  $a$  is also an upper bound for  $B$ .
- (b) If every  $a_n$  is in the complement of the interval  $(0, 1)$ , then  $a$  is also in the complement of  $(0, 1)$ .
- (c) If every  $a_n$  is rational, then  $a$  is rational.

### SOLUTION

- (a) True. For all  $n \in \mathbb{N}$ ,  $a_n \geq \sup B$ , by Order Limit Theorem,  $a = \lim a_n \geq \sup B$ .
- (b) True. If  $a \in (0, 1)$ , then there would exist an  $\epsilon$ -neighborhood inside  $(0, 1)$  that  $a_n$  would have to fall in, contradicting the fact that  $a \notin (0, 1)$ .
- (c) False. Consider the sequence  $(1, 1.4, 1.41, 1.414, \dots) \rightarrow \sqrt{2}$ .

### Exercise 2.3.13

**(Iterated Limits).** Given a doubly indexed array  $a_{mn}$  where  $m, n \in \mathbb{N}$ , what should  $\lim_{m, n \rightarrow \infty} a_{mn}$  represent?

- (a) Let  $a_{mn} = m/(m+n)$  and compute the *iterated* limits

$$\lim_{n \rightarrow \infty} \left( \lim_{m \rightarrow \infty} a_{mn} \right) \quad \text{and} \quad \lim_{m \rightarrow \infty} \left( \lim_{n \rightarrow \infty} a_{mn} \right).$$

Define  $\lim_{m, n \rightarrow \infty} a_{mn} = a$  to mean that all  $\epsilon > 0$  there exists an  $N \in \mathbb{N}$  such that if both  $m, n \geq N$ , then  $|a_{mn} - a| < \epsilon$ .

- (b) Let  $a_{mn} = 1/(m+n)$ . Does  $\lim_{m, n \rightarrow \infty} a_{mn}$  exist in this case? Do the two iterated limits exist? How do these three values compare? Answer these same questions for  $a_{mn} = mn/(m^2 + n^2)$ .
- (c) Produce an example where  $\lim_{m, n \rightarrow \infty} a_{mn}$  exists but neither iterated limit can be computed.
- (d) Assume  $\lim_{m, n \rightarrow \infty} a_{mn} = a$ , and assume that for each fixed  $m \in \mathbb{N}$ ,  $\lim_{n \rightarrow \infty} (a_{mn}) \rightarrow b_m$ . Show  $\lim_{m \rightarrow \infty} b_m = a$ .
- (e) Prove that if  $\lim_{m, n \rightarrow \infty} a_{mn}$  exists and the iterated limits both exist, then all three limits must be equal.

### SOLUTION

- (a)

$$\lim_{n \rightarrow \infty} \left( \lim_{m \rightarrow \infty} a_{mn} \right) = 1 \quad \text{and} \quad \lim_{m \rightarrow \infty} \left( \lim_{n \rightarrow \infty} a_{mn} \right) = 0$$

- (b) For  $a_{mn} = a/(m+n)$ , both iterated limits exist and equal to  $\lim_{m, n \rightarrow \infty} a_{mn} = 0$ .

For  $a_{mn} = mn/(m^2 + n^2)$ , both iterated limits equal to zero. However,  $\lim_{m, n \rightarrow \infty} a_{mn}$  does not exist.

- (c) Consider  $b_{mn} = 1/\min\{m, n\}$ . WLOG, we can fix  $m$  and as  $n$  approaches infinity,  $n > m$  causing the limit to be  $1/m$ . To ensure that the limit does not exist, we can introduce the term  $(-1)^{n+m}$ , giving

$$a_{mn} = \frac{(-1)^{m+n}}{\min\{m, n\}}$$

This works as the inner limits of the iterated limits are not defined since it is oscillating between  $1/m$  and  $-1/m$ . However, if both  $m, n \geq N$  for some  $N > 0$ , the limit equals zero as  $0 < \lim |a_{mn}| \leq \lim |1/(m+n)| \leq \lim |1/2N| = 0$ .



- (d) For  $\epsilon > 0$ , we can choose  $N > 0$  such that for all  $m, n > N_0$ ,  $|a_{mn} - a| < \epsilon/2$ . For each fixed  $m \in \mathbb{N}$ , we can choose  $N_m > 0$  such that for  $n > N_m$ ,  $|a_{mn} - b_m| < \epsilon/2$ . By triangle inequality,

$$|b_m - a| = |b_m - a_{mn} + a_{mn} - a| \leq |b_m - a_{mn}| + |a_{mn} - a| \leq \epsilon/2 + \epsilon/2 = \epsilon$$

Hence we can choose  $N = \max\{N_0, N_m\}$  and we are done.

- (e) Let  $b_m = \lim_{n \rightarrow \infty} (a_{mn})$ ,  $c_n = \lim_{m \rightarrow \infty} (a_{mn})$ , and  $a = \lim_{m, n \rightarrow \infty} (a_{mn})$ . In (d), we have shown that  $(b_m) \rightarrow a$ . A similar argument shows  $(c_n) \rightarrow a$ . Thus all three limits are equal to  $a$ .

## 2.4 The Monotone Convergence Theorem and a First Look at Infinite Series

### Exercise 2.4.1

- (a) Prove that the sequence defined by  $x_1 = 3$  and

$$x_{n+1} = \frac{1}{a - x_n}$$

converges.

- (b) Now that we know  $\lim x_n$  exists, explain why  $\lim x_{n+1}$  must also exist and equal the same value.  
 (c) Take the limit of each side of the recursive equation in part (a) to explicitly compute  $\lim x_n$ .

### SOLUTION

- (a) Calculating the first few terms, we get  $(3, 1, 1/3, \dots)$ . I conjecture that  $(x_n)$  is decreasing.

Suppose  $x_{n+1} < x_n$ , then

$$4 - x_n > 4 - x_{n+1} \implies \frac{1}{4 - x_n} < \frac{1}{4 - x_{n+1}} \implies x_{n+2} < x_{n+1}$$

Thus,  $(x_n)$  is decreasing. Since  $x_n < 4$  for all  $n \in \mathbb{N}$ ,  $4 - x_n > 0$  and hence  $x_{n+1} > 0$ . Since  $(x_n)$  is decreasing and bounded below,  $(x_n)$  converges.

- (b) By definition of limits, changing the index of the sequence does not change its limit.  
 (c) Since  $x = \lim x_n = \lim x_{n+1}$ ,

$$x_{n+1} = \frac{1}{4 - x_n} \implies x = \frac{1}{4 - x} \implies x^2 - 4x + 1 = 0 \implies x = 2 \pm \sqrt{3}$$

Since  $x_n < 3$ , then  $x < 3$ , hence  $x \neq 2 + \sqrt{3}$  and  $x = 2 - \sqrt{3}$ .

### Exercise 2.4.2

- (a) Consider the recursively defined sequence  $y_1 = 1$ ,

$$y_{n+1} = 3 - y_n,$$

and set  $y = \lim y_n$ . Because  $(y_n)$  and  $(y_{n+1})$  have the same limit, taking the limit across the recursive equation gives  $y = 3 - y$ . Solving for  $y$ , we conclude  $\lim y_n = 3/2$ .

What is wrong with this argument?

- (b) This time set  $y_1 = 1$  and  $y_{n+1} = 3 - \frac{1}{y_n}$ . Can the strategy in (a) be applied to compute the limit of this sequence?

### SOLUTION

- (a) The argument only works if the series converges, which does not in this case.

- (b) Yes it can as  $0 < y_n < 3$  and  $(y_n)$  is increasing (Proving this is left as an exercise for the reader).

**Exercise 2.4.3**

- (a) Show that

$$\sqrt{2}, \sqrt{2 + \sqrt{2}}, \sqrt{2 + \sqrt{2 + \sqrt{2}}}, \dots$$

converges and find the limit.

- (b) Does the sequence

$$\sqrt{2}, \sqrt{2\sqrt{2}}, \sqrt{2\sqrt{2\sqrt{2}}}, \dots$$

converge? If so, find the limit.

**SOLUTION**

- (a) Let  $x_1 = \sqrt{2}$  and  $x_{n+1} = \sqrt{2 + x_n}$ , clearly  $x_2 > x_1$ . Suppose  $x_{n+1} > x_n$ , then

$$2 + x_{n+1} > 2 + x_n \implies \sqrt{2 + x_{n+1}} > \sqrt{2 + x_n} \implies x_{n+2} > x_{n+1}$$

showing that  $x_n$  is monotonically increasing.

Next we show that  $x_n \leq 2$ , clearing  $x_1 = \sqrt{2} \leq 2$ . Suppose  $x_n \leq 2$ , then

$$2 + x_n \leq 4 \implies \sqrt{2 + x_n} \leq 2 \implies x_{n+1} \leq 2$$

Since  $(x_n)$  is monotone and bounded, the monotone convergence theorem tells us that  $(x_n) \rightarrow x$ . Equating the limits on both sides gives

$$x = \sqrt{2 + x} \implies x^2 - x - 2 = 0 \implies x = \frac{1}{2} \pm \frac{3}{2}$$

Since  $x > 0$ , we have  $x = 2$ .

- (b) Let  $x_1 = \sqrt{2}$  and  $x_{n+1} = \sqrt{2x_n}$ , clearly  $x_2 > x_1$ . Suppose  $x_{n+1} > x_n$ , then

$$2x_{n+1} > 2x_n \implies \sqrt{2x_{n+1}} > \sqrt{2x_n} \implies x_{n+2} > x_{n+1}$$

showing that  $(x_n)$  is monotonically increase. Next we show that  $x_n \leq 2$ , which is clearly true for  $x_1$ . Suppose  $x_n \leq 2$ , then

$$2x_n \leq 4 \implies x_{n+1} = \sqrt{2x_n} \leq 2$$

Hence,  $x_n$  is bounded. By monotone convergence theorem, the sequence converges, hence  $(x_n) \rightarrow x$ . Taking limits on both sides,

$$x = \sqrt{2x} \implies x^2 - 2x = 0 \implies x = 1 \pm 1$$

Since  $x > 0$ , then  $x = 2$ .

**Exercise 2.4.4**

- (a) In Section 1.4, we used the Axiom of Completeness (AoC) to prove the Archimedean Property of  $\mathbb{R}$  (Theorem 1.4.2). Show that the Monotone Convergence Theorem can also be used to prove the Archimedean Property without making any use of AoC.
- (b) Use the Monotone Convergence Theorem to supply a proof for the Nested Interval Property (Theorem 1.4.1) that doesn't make use of AoC

These two results suggest that we could have used the Monotone Convergence Theorem in place of AoC as our starting axiom for building a proper theory of the real numbers.

**SOLUTION**

- (a) MCT tells us  $(1/n)$  converges, obviously it must converge to zero, giving us  $|1/n - 0| = 1/n < \epsilon$  for any  $\epsilon > 0$ , which is the Archimedean Property.

- (b) We have  $I_n = [a_n, b_n]$  with  $a_n \leq b_n$  since  $I_n \neq \emptyset$ . Since  $I_{n+1} \subseteq I_n$ , we must have  $b_{n+1} \leq b_n$  and  $a_{n+1} \geq a_n$ , the MCT tells us that  $(a_n) \rightarrow a$  and  $(b_n) \rightarrow b$ . By Order Limit Theorem, we have  $a \leq b$ , hence  $a \in I_n$  for all  $n \in \mathbb{N}$ , meaning  $a \in \bigcap_{n=1}^{\infty} I_n \neq \emptyset$ .

**Exercise 2.4.5**

**(Calculating Square Roots)** Let  $x_1 = 2$ , and define

$$x_{n+1} = \frac{1}{2} \left( x_n + \frac{2}{x_n} \right).$$

- (a) Show that  $x_n^2$  is always greater than or equal to 2, and then use this to prove that  $x_n - x_{n+1} \geq 0$ . Conclude that  $\lim x_n = \sqrt{2}$ .
- (b) Modify the sequence  $(x_n)$  so that it converges to  $\sqrt{c}$ .

**SOLUTION**

- (a) Clearing  $x_1 \geq 2$ . For any  $n$ ,

$$\begin{aligned} x_{n+1}^2 &= \left[ \frac{1}{2} \left( x_n + \frac{2}{x_n} \right) \right]^2 \\ &= \frac{1}{4} \left( x_n^2 + 4 + \frac{4}{x_n^2} \right) \\ &= \frac{1}{4} \left( x_n^2 - 4 + \frac{4}{x_n^2} \right) + 2 \\ &= \frac{1}{4} \left( x_n + \frac{2}{x_n} \right) + 2 \geq 2 \end{aligned}$$

Now we show  $x_n - x_{n+1} \geq 0$ .

$$\begin{aligned} x_n - x_{n+1} &= x_n - \frac{1}{2} \left( x_n + \frac{2}{x_n} \right) \\ &= \frac{1}{2} x_n - \frac{1}{x_n} \geq 0 \because x_n \geq 0 \end{aligned}$$

MCT tells us that  $(x_n) \rightarrow x$ . By taking limits on both sides,

$$x = \frac{1}{2} \left( x + \frac{2}{x} \right) \implies \frac{1}{2} x^2 = 1 \implies x = \pm \sqrt{2}$$

Since  $x_n \geq 0$ ,  $x = \sqrt{2}$ .

- (b)

$$x_{n+1} = \frac{1}{2} \left( x_n + \frac{c}{x_n} \right)$$

Proof of this is similar to (a) and left as an exercise for the reader.

**Exercise 2.4.6**

**(Arithmetic-Geometric Mean)**

- (a) Explain why  $\sqrt{xy} \leq (x+y)/2$  for any two positive real numbers  $x$  and  $y$ . (The geometric mean is always less than the arithmetic mean.)
- (b) Now let  $0 \leq x_1 \leq y_1$  and define

$$x_{n+1} = \sqrt{x_n y_n} \quad \text{and} \quad y_{n+1} = \frac{x_n + y_n}{2}$$

Show  $\lim x_n$  and  $\lim y_n$  both exist and are equal.

**SOLUTION**

(a) We have

$$\sqrt{xy} \leq \frac{x+y}{2} \iff 4xy \leq x^2 + 2xy + y^2 \iff 0 \leq (x-y)^2$$

(b) As shown above, if  $\lim x_n = x$  and  $\lim y_n = y$ , then  $x_n = y_n$  is the only fixed point, hence  $x = y$ . Thus we only need to show both sequences converge.

The inequality  $0 \leq x_n \leq y_n$  is always true. It is obvious both terms will always be positive. Suppose  $x_n \leq y_n$ , then by (a),  $x_{n+1} \leq y_{n+1}$  true.

To continue,  $x_n \leq y_n$  implies  $(x_n + y_n)/2 = y_{n+1} \leq y_n$ . Similarly,  $\sqrt{x_n y_n} = x_{n+1} \geq x_n$ . This means both sequences are monotone and bounded by each other, hence Monotone Convergence Theorem tells us both sequences converge.

**Exercise 2.4.7**

**(Limit Superior)** Let  $(a_n)$  be a bounded sequence.

(a) Prove that the sequence defined by  $y_n = \sup\{a_k : k \geq n\}$  converges.

(b) The *limit superior* of  $(a_n)$ , or  $\limsup a_n$ , is defined by

$$\limsup a_n = \lim y_n,$$

where  $y_n$  is the sequence from part (a) of this exercise. Provide a reasonable definition for  $\liminf a_n$  and briefly explain why it always exists for any bounded sequence.

(c) Prove that  $\liminf a_n \leq \limsup a_n$  for every bounded sequence, and give an example of a sequence for which the inequality is strict.

(d) Show that  $\liminf a_n = \limsup a_n$  if and only if  $\lim a_n$  exists. In this case, all three share the same value.

**SOLUTION**

(a)  $(y_n)$  is decreasing, hence MCT tells us it converges.

(b) The *limit inferior* of  $(a_n)$ , or  $\liminf a_n$ , is defined by  $\liminf a_n = \lim x_n$ , where  $x_n = \inf\{a_k : k \geq n\}$ . Since  $(x_n)$  is increasing, MCT tells us it converges, hence the limit inferior exists.

(c) Obviously,  $\inf\{a_k : k \geq n\} \leq \sup\{a_k : k \geq n\}$ , so Order Limit Theorem implies  $\liminf a_n \leq \limsup a_n$ . The equality is strict when the series does not converge, for example  $a_n = (-1)^n$ .

(d)  $(\Rightarrow)$  If  $\liminf a_n = \limsup a_n$ , then Squeeze Theorem implies that  $a_n$  converges to the same value, since  $\inf\{a_{k \geq n}\} \leq a_n \leq \sup\{a_{k \geq n}\}$ .

$(\Leftarrow)$  Assume that  $\lim a_n = a$ . For  $\epsilon > 0$ , we can find  $N > 0$  such that for all  $k > N$ ,  $|a_k - a| < \epsilon$ . This means

$$a - \epsilon < x_N \leq x_k \leq y_k \leq y_N < a + \epsilon$$

implying that both  $x_n$  and  $y_n$  converge to  $a$ .

**Exercise 2.4.8**

For each series, find an explicit formula for the sequence of partial sums and determine if the series converges.

$$(a) \sum_{n=1}^{\infty} \frac{1}{2^n} \quad (b) \sum_{n=1}^{\infty} \frac{1}{n(n+1)} \quad (c) \sum_{n=1}^{\infty} \log\left(\frac{n+1}{n}\right)$$

(In (c),  $\log(x)$  refers to the natural logarithm function from calculus.)

**SOLUTION**

(a) This is a geometric series, allowing us to use the usual trick to determine  $s_n$ .

$$\begin{aligned}s_n &= 1 + \frac{1}{2} + \frac{1}{4} + \cdots + \frac{1}{2^n} \\ \frac{1}{2}s_n &= \frac{1}{2} + \frac{1}{4} + \cdots + \frac{1}{2^{n+1}} \\ \frac{1}{2}s_n - s_n &= \frac{1}{2^{n+1}} - 1 \implies s_n = \frac{2^{-n-1} - 1}{2^{-1} - 1}\end{aligned}$$

As  $n$  tends to infinity,

$$\sum_{n=1}^{\infty} \frac{1}{2^n} = -1 + \sum_{n=1}^{\infty} \frac{1}{2^n} = -1 + \lim_{n \rightarrow \infty} \frac{2^{-n-1} - 1}{2^{-1} - 1} = 1$$

(b) We can use partial fractions to get

$$\frac{1}{n(n+1)} = \frac{1}{n} - \frac{1}{n+1}$$

which gives a telescoping sum

$$s_n = 1 - \frac{1}{n+1}$$

Therefore,

$$\sum_{n=1}^{\infty} \frac{1}{n(n+1)} = \lim_{n \rightarrow \infty} \left(1 - \frac{1}{n+1}\right) = 1$$

(c) Similar to (b), we can use telescoping sum since

$$\log\left(\frac{n+1}{n}\right) = \log(n+1) - \log(n)$$

Hence, we get

$$\sum_{n=1}^{\infty} \log\left(\frac{n+1}{n}\right) = \log(n+1)$$

which does not converge.

**2.5 Subsequences and the Bolzano-Weierstrass Theorem****2.6 The Cauchy Criterion****2.7 Properties of Infinite Series****2.8 Double Summations and Products of Infinite Series**