

Table 2. Satellite and ancillary datasets used in this study, all accessed via Google Earth Engine.

Dataset	GEE Collection ID	Temporal Range	Native Resolution	Role
Sentinel-1 GRD (IW, VH+VV, descending)	COPERNICUS/S1_GRD	2017–2024	10 m	Primary SAR backscatter; classifier features; phenology retrieval
Sentinel-2 L2A (harmonised)	COPERNICUS/S2_SR_HARMONIZED	2017–2024	10 m	Optical indices (NDVI, NDWI, LSWI, CIRE); cloud-free phenology support
JRC GSW Monthly History v1.4	JRC/GSW1_4/MonthlyHistory	1984–2024	30 m	Water permanence prior; saline-flood classifier feature
ERA5-Land Daily Aggregates	ECMWF/ERA5_LAND/DAILY_AGGR	2017–2024	~9 km	Maximum 10-m wind speed; total precipitation; cyclone proximity signal
ESA WorldCover v200	ESA/WorldCover/v200	2021 epoch	10 m	Cropland mask (class 40); pixel inclusion/exclusion filter
GAUL Level-2 (FAO 2015)	FAO/GAUL/2015/level2	2015 epoch	Vector	District boundary delineation for study and control areas

Source: see manuscript §2.2–2.3 for full provenance. Identical to the inline Table appearing in the main manuscript file.