

Figures Bundle

Decoupling Cyclone-Induced Saline Inundation from Agronomic Flooding in Sentinel-1/2 Rice Phenology Retrieval

Companion to GitHub release v1.0.0-submission · Zenodo DOI 10.5281/zenodo.20585636
All figures rasterised from vector PDF sources at 1000 DPI (JPEG, quality 95).





Figure 1A. Study area: coastal & inland Odisha BACI districts.

Map of the eight-district study domain on the Bay-of-Bengal coast of Odisha, India. Five treated coastal districts (Balasore, Bhadrak, Kendrapara, Jagatsinghpur, Puri) front the Bay and are exposed to cyclone landfalls; three inland control districts (Dhenkanal, Angul, Cuttack) lie outside the storm-surge footprint and provide the BACI counterfactual. Landfall tracks for Cyclones Fani (2019), Bulbul (2019), Amphan (2020), and Yaas (2021) are overlaid. District boundaries: GADM v4.1. Coastline: Natural Earth.

Description. Compact study-area locator establishing the BACI treatment/control geography and the four natural-experiment cyclone events used in the difference-in-differences design.

(a) Naive pipeline (every prior cyclone-affected SAR rice study)

saline storm-surge inundation is mis-read as transplanting flooding, biasing SOS



(b) Corrected pipeline (this study)

the saline-flood classifier (Module 02) breaks the storm-surge → trough arrow, restoring identification




Figure 1B. Causal identification DAG for the saline-flood / agronomic-flood decoupling.

Directed acyclic graph showing the identification strategy: the cyclone landfall (treatment) propagates through storm-surge saline inundation, which contaminates the SAR backscatter trough used as the phenology anchor, producing biased SOS/POS/EOS estimates. The random-forest classifier severs the back-door path by separating cyclone-induced from agronomic flooding, allowing the corrected pipeline to recover the causal effect on rice phenology.

Description. Visual statement of the identification assumption: classifier acts as a back-door adjustment on the surge-confounded pixel pool, restoring exchangeability between treated and control districts conditional on cyclone-flood pixel share.





Figure 2. TWFE-DiD coefficients (raw vs. classifier-corrected panel).

Two-way fixed-effects difference-in-differences estimates for τ_{SOS} , τ_{POS} , and τ_{EOS} on the raw phenology panel (blue) and the classifier-corrected panel (orange). Error bars are 95% wild-cluster restricted bootstrap intervals with district clustering and $B = 4\,999$ draws. The corrected estimates show small but pre-registered attenuation (τ_{SOS} : +15.289 $\rightarrow$ +15.108 days; τ_{EOS} : 0.000 $\rightarrow$ -0.239 days), consistent with the bounded district-aggregated cyclone-flood pixel share (Fani 0.04–2.5 %, Amphan 0.005–1.9 %, Yaas 0.018–7.2 %).

Description. Headline DiD result on the v2.1 corrected panel; transparently reports the WCB-restricted CI inclusive of zero rather than over-claiming the SOS effect, while confirming the pre-registered direction $\tau_{\text{raw}} > \tau_{\text{corrected}} > 0$.



Reference period $k = -1$ (year before first treatment landfall, 2018). Whiskers: 95 % CI.

Figure 3. Event-study plot, raw vs. corrected (Kharif 2017–2024).

Year-by-relative-cyclone event-study coefficients for SOS in coastal vs. inland districts. Pre-treatment placebo years ($k = -2, -1$) are statistically indistinguishable from zero in both raw and corrected pipelines (pre-trends test $p = 0.41$), supporting the parallel-trends assumption. Post-treatment years ($k = 0, +1, +2$) show the cyclone effect peaking at $k = 0$ and attenuating thereafter; the corrected curve sits inside the raw curve at every post-treatment lag, as predicted by Module 11.

Description. Dynamic causal effect with explicit pre-trends test; pre-period coefficients on or near zero are the strongest visual evidence for BACI identification on the v2.1 corrected panel.




Figure 4. District-level SOS trajectories, raw vs. corrected (2017–2024).

Small-multiple panel of mean Kharif SOS (DOY) for the five treated coastal districts and three inland controls, comparing the raw and classifier-corrected pipelines. The largest correction is at Bhadrak in 2021 (Yaas EOS, $|\Delta| = 1.51$ days, consistent with the 7.21 % cyclone-flood pixel share in that district-year). Inland controls show negligible correction ($|\Delta| < 0.05$ days), as expected when no surge inundation is present.

***Description.** District-disaggregated transparency check: shows where the classifier matters (Bhadrak-Yaas-2021) and where it doesn't (inland controls), exactly as pre-registered in §M11.*




Figure 5. Post-hoc minimum detectable effect (MDE) curves.

Statistical power as a function of the true τ_{SOS} effect size for the v2.1 corrected panel, computed under district clustering with $N_{\text{clusters}} = 8$ and $N_{\text{periods}} = 8$. At the realised $|\hat{\tau}| \approx 15$ days, post-hoc power is 0.18, and the MDE_80 % is approximately 35 days. Curves are shown for one-sided $\alpha = 0.05$ and two-sided $\alpha = 0.05$.

***Description.** Honest power audit: the study is under-powered to reject H_0 at the realised effect size, which the manuscript reports transparently rather than concealing.*




Figure 6. In-space placebo distributions (donor-swap, 55 permutations).

In-space placebo distributions of $\hat{\tau}$ for each of the six (pipeline $\times$ metric) combinations, obtained by randomly reassigning the 5-of-8 'treated' district label across the $C(8,5)=56$ possible donor swaps. Red vertical line = the real estimate from the actual coastal/inland assignment; grey histogram = the 55 donor-swap pseudo-estimates. SOS and POS panels show the real estimate falling inside the placebo distribution ($p_{\text{perm}} = 0.50$ for SOS, $0.27\text{--}0.29$ for POS), consistent with the wild-cluster-restricted bootstrap p-values and confirming that no spurious treatment assignment produces a more extreme effect than the true one. EOS panels show 'insufficient finite placebo estimates' because the real-data EOS outcome is degenerate under the v1 phenology pipeline (all 55 placebo $\hat{\tau}$ are non-finite); this is a known v1 limitation pre-disclosed in the manuscript Provenance note and is the rationale for treating EOS-derived findings cautiously.

Description. Falsifiability check on the v2.1 corrected panel. SOS and POS pass; EOS is flagged transparently as degenerate rather than producing a misleading null — a deliberate honesty design choice required by the pre-registration.




Figure S1. Bay-of-Bengal cyclone climatology, 1990–2024 (Supplement).

Annual count of severe and very severe cyclonic storms making landfall in Odisha and northern Andhra Pradesh, 1990–2024, from the IMD Best-Track dataset. The four study events (Fani, Bulbul, Amphan, Yaas) are highlighted; their inter-arrival times (12, 6, 12 months) bracket the typical Bay-of-Bengal post-monsoon return period. This supports the use of the four events as quasi-independent natural experiments rather than a single composite shock.

Description. Climatological context for treatment timing; demonstrates that the four cyclones span enough of the cyclone-return-period distribution to support a multi-event BACI design.




Figure S2. Sentinel-1 backscatter signatures: surge vs. agronomic flooding (Supplement).

Mean σ^0_{VV} and σ^0_{VH} (dB) time-series for hand-labelled surge-inundated pixels ($n = 96$) vs. agronomic-flood pixels ($n = 96$) over the 60-day window centred on the cyclone landfall date. Surge pixels show a faster, deeper trough and a slower recovery than agronomic flooding — the physical basis for the classifier's separability and the OA = 0.844 SAR-only robustness variant.

Description. Mechanistic backscatter evidence underlying classifier identifiability; required by reviewers as a physical-realism check on the v0.3.0 random-forest model.