

# Elementary Number Theory

KTW

October 3, 2023

---

*Contents*

<b>Contents</b>	<b>2</b>
<b>1 Preliminaries</b>	<b>3</b>
1.1 Mathematical Induction . . . . .	3
1.2 The Binomial Theorem . . . . .	9

Preliminaries

## 1.1 Mathematical Induction

**Well-Ordering Principle.** *Every nonempty set  $S$  of nonnegative integers contains a least element; that is, there is some integer  $a$  in  $S$  such that  $a \leq b$  for all  $b$ 's belonging to  $S$ .*

$$\forall S \subseteq \mathbb{Z}_{\geq 0}, \exists a \in S : \forall b \in S, a \leq b$$

**Theorem 1.1** (Archimedean Property). *If  $a$  and  $b$  are any positive integers, then there exists a positive integer  $n$  such that  $na \geq b$ .*

$$\forall a, b \in \mathbb{Z}_{>0}, \exists n \in \mathbb{Z}_{>0} : na \geq b$$

*Proof.* For reductio ad absurdum, suppose that the statement of the theorem is false. Then:

$$\begin{aligned} & \neg(\forall a, b \in \mathbb{Z}_{>0}, \exists n \in \mathbb{Z}_{>0} : na \geq b) \\ \Leftrightarrow & \exists a, b \in \mathbb{Z}_{>0} : \forall n \in \mathbb{Z}_{>0}, na < b \\ \Leftrightarrow & \exists a, b \in \mathbb{Z}_{>0} : \forall n \in \mathbb{Z}_{>0}, 0 < b - na \\ \Leftrightarrow & S = \{b - na \mid n \in \mathbb{Z}_{>0}\} = \mathbb{Z}_{>0} \\ \Rightarrow & \exists b - ma \in S : \forall b - na \in S, b - ma \leq b - na \quad [\text{by the Well-Ordering Principle}] \\ \Rightarrow & \exists b - ma, b - (m+1)a \in S : \forall b - na \in S, b - (m+1)a < b - ma \leq b - na \longrightarrow \perp \\ \Rightarrow & \forall a, b \in \mathbb{Z}_{>0}, \exists n \in \mathbb{Z}_{>0} : na \geq b \end{aligned}$$

□

**Theorem 1.2** (First Principle of Finite Induction). *Let  $S$  be a set of positive integers with the following properties:*

- (a)  $\exists 1 \in S$
- (b)  $\forall k \in S, \exists k+1 \in S$

*Then  $S$  is the set of all positive integers.*

*Proof.* Let  $S$  be a set of positive integers such that:

$$\exists 1 \in S \ \& \ \forall k \in S, \exists k+1 \in S$$

For reductio ad absurdum, let  $T$  be a nonempty set of all positive integers not in  $S$ , that is:

$$T = \{t \in \mathbb{Z}_{>0} \mid t \notin S\}$$

By the Well-Ordering Principle,  $T$  has a least element  $a$ , and:

$$\begin{aligned} 1 \in S & \Rightarrow 1 < a \in T \\ \Leftrightarrow & 0 < a-1 \notin T \\ \Leftrightarrow & a-1 \in S \\ \Leftrightarrow & a \in S \longrightarrow \perp \\ \Rightarrow & T = \emptyset \\ \Leftrightarrow & S = \mathbb{Z}_{>0} \end{aligned}$$

□

**Theorem 1.3** (Second Principle of Finite Induction). *Let  $S$  be a set of positive integers with the following properties:*

$$(a) \exists 1 \in S$$

$$(b) \exists 1, 2, \dots, k \in S \Rightarrow \exists k+1 \in S$$

*Then  $S$  is the set of all positive integers.*

*Proof.* Let  $S$  be a set of positive integers following properties above. For reductio ad absurdum, let  $T$  be a nonempty set of all positive integers not in  $S$ . By the Well-Ordering Principle,  $T$  has a least element  $a$ , and:

$$\begin{aligned} 1 \in S &\Rightarrow 1 < a \in T \\ &\Leftrightarrow 0 < 1, \dots, a-1 \notin T \\ &\Leftrightarrow 1, \dots, a-1 \in S \\ &\Leftrightarrow a \in S \longrightarrow \perp \\ &\Rightarrow T = \emptyset \\ &\Leftrightarrow S = \mathbb{Z}_{>0} \end{aligned}$$

□

**Example 1.4** (Lucas sequence).

$$1, 3, 4, 7, 11, 18, 29, 47, 76, \dots$$

Sequence above may be defined inductively by

$$\begin{cases} a_1 = 1 \\ a_2 = 3 \\ a_n = a_{n-1} + a_{n-2} \quad \text{for all } n \geq 3 \end{cases}$$

We contend that the inequality

$$a_n < \left(\frac{7}{4}\right)^n$$

holds for every positive integer  $n$ . First of all, for  $n = 1$  and  $2$ , we have

$$a_1 = 1 < \left(\frac{7}{4}\right)^1 \quad \& \quad a_2 = 3 < \left(\frac{7}{4}\right)^2 = 3\frac{1}{16}$$

and this provides a basis for the induction. For the induction step, choose an integer  $k \geq 3$  and assume that the inequality is valid for  $n = 1, 2, \dots, k-1$ . Then, in particular:

$$a_{k-1} < \left(\frac{7}{4}\right)^{k-1} \quad \& \quad a_{k-2} < \left(\frac{7}{4}\right)^{k-2}$$

By the way in which the sequence is formed, it follows that:

$$\begin{aligned} a_k = a_{k-1} + a_{k-2} &< \left(\frac{7}{4}\right)^{k-1} + \left(\frac{7}{4}\right)^{k-2} \\ &= \left(\frac{7}{4}\right)^{k-2} \left(\frac{7}{4} + 1\right) \\ &= \left(\frac{7}{4}\right)^{k-2} \left(\frac{11}{4}\right) \\ &< \left(\frac{7}{4}\right)^{k-2} \left(\frac{7}{4}\right)^2 = \left(\frac{7}{4}\right)^k \end{aligned}$$

Because the inequality is true for  $n = k$  whenever it is true for the integers  $1, 2, \dots, k-1$ , we conclude by the second induction principle that  $a_n < (7/4)^n$  for all  $n \geq 1$ .

**Problems 1.1**

1. Establish the formulas below by mathematical induction:

- (a)  $1 + 2 + 3 + \cdots + n = \frac{n(n+1)}{2}$  for all  $n \geq 1$ .
- (b)  $1 + 3 + 5 + \cdots + (2n-1) = n^2$  for all  $n \geq 1$ .
- (c)  $1 \cdot 2 + 2 \cdot 3 + 3 \cdot 4 + \cdots + n(n+1) = \frac{n(n+1)(n+2)}{3}$  for all  $n \geq 1$ .
- (d)  $1^2 + 3^2 + 5^2 + \cdots + (2n-1)^2 = \frac{n(2n-1)(2n+1)}{3}$  for all  $n \geq 1$ .
- (e)  $1^3 + 2^3 + 3^3 + \cdots + n^3 = \left[ \frac{n(n+1)}{2} \right]^2$  for all  $n \geq 1$ .

#1(a). Let  $S$  be a set such that:

$$S = \left\{ n \in \mathbb{N} : 1 + 2 + 3 + \cdots + n = \frac{n(n+1)}{2} \right\}$$

$1 \in S$  because for  $n = 1$ :

$$1 = \frac{1(1+1)}{2}$$

Let any  $k \in \mathbb{N}$  be the member of  $S$ , that is:

$$1 + 2 + 3 + \cdots + k = \frac{k(k+1)}{2}$$

Then;

$$\begin{aligned} 1 + 2 + 3 + \cdots + k + (k+1) &= \frac{k(k+1)}{2} + (k+1) \\ &= \frac{k(k+1)}{2} + \frac{2(k+1)}{2} \\ &= \frac{(k+1)(k+2)}{2} \end{aligned}$$

therefore by the First Principle of Finite Induction,  $S = \mathbb{N}$ . ■

#1(b). Let  $S$  be a set such that:

$$S = \left\{ n \in \mathbb{N} : 1 + 3 + 5 + \cdots + (2n-1) = n^2 \right\}$$

$1 \in S$  because for  $n = 1$ :

$$(2 \cdot 1 - 1) = 1 = 1^2$$

Let any  $k \in \mathbb{N}$  be the member of  $S$ , that is:

$$1 + 3 + 5 + \cdots + (2k-1) = k^2$$

Then;

$$\begin{aligned} 1 + 3 + \cdots + (2k-1) + [2(k+1)-1] &= k^2 + [2(k+1)-1] \\ &= k^2 + 2k + 1 \\ &= (k+1)^2 \end{aligned}$$

therefore by the First Principle of Finite Induction,  $S = \mathbb{N}$ . ■

#1(c). Let  $S$  be a set such that:

$$S = \left\{ n \in \mathbb{N} : 1 \cdot 2 + 2 \cdot 3 + \cdots + n(n+1) = \frac{n(n+1)(n+2)}{3} \right\}$$

$1 \in S$  because for  $n = 1$ :

$$1(1+1) = 2 = \frac{1(1+1)(1+2)}{3}$$

Let any  $k \in \mathbb{N}$  be the member of  $S$ , that is:

$$1 \cdot 2 + 2 \cdot 3 + \cdots + k(k+1) = \frac{k(k+1)(k+2)}{3}$$

Then;

$$\begin{aligned} 1 \cdot 2 + 2 \cdot 3 + \cdots + k(k+1) + (k+1)(k+2) &= \frac{k(k+1)(k+2)}{3} + (k+1)(k+2) \\ &= \frac{k(k+1)(k+2)}{3} + \frac{3(k+1)(k+2)}{3} \\ &= \frac{(k+1)(k+2)(k+3)}{3} \end{aligned}$$

therefore by the First Principle of Finite Induction,  $S = \mathbb{N}$ .

**#1(d).** Let  $S$  be a set such that:

$$S = \left\{ n \in \mathbb{N} : 1^2 + 3^2 + 5^2 + \cdots + (2n-1)^2 = \frac{n(2n-1)(2n+1)}{3} \right\}$$

$1 \in S$  because for  $n = 1$ :

$$(2 \cdot 1 - 1)^2 = 1 = \frac{3}{3}$$

Let any  $k \in \mathbb{N}$  be the member of  $S$ , that is:

$$1^2 + 3^2 + \cdots + (2k-1)^2 = \frac{k(2k-1)(2k+1)}{3}$$

Then;

$$\begin{aligned} 1^2 + 3^2 + \cdots + (2k-1)^2 + (2k+1)^2 &= \frac{k(2k-1)(2k+1)}{3} + (2k+1)^2 \\ &= \frac{k(2k-1)(2k+1)}{3} + \frac{3(2k+1)^2}{3} \\ &= \frac{(2k+1)[k(2k-1) + 3(2k+1)]}{3} \\ &= \frac{(2k+1)[2k^2 + 5k + 3]}{3} \\ &= \frac{(2k+1)(2k+3)(k+1)}{3} \\ &= \frac{(k+1)(2k+1)(2k+3)}{3} \end{aligned}$$

therefore by the First Principle of Finite Induction,  $S = \mathbb{N}$ . ■

**#1(e).** Let  $S$  be a set such that:

$$S = \left\{ n \in \mathbb{N} : 1^3 + 2^3 + \cdots + n^3 = \left[ \frac{n(n+1)}{2} \right]^2 \right\}$$

$1 \in S$  for  $n = 1$  becuase:

$$1^3 = 1 = 1^2$$

Let any  $k \in \mathbb{N}$  be the member of  $S$ , that is:

$$1^3 + 2^3 + \cdots + k^3 = \left[ \frac{k(k+1)}{2} \right]^2$$

Then;

$$\begin{aligned} 1^3 + 2^3 + \cdots + k^3 + (k+1)^3 &= \left[ \frac{k(k+1)}{2} \right]^2 + (k+1)^3 \\ &= \frac{k^2}{4}(k+1)(k+1) + (k+1)(k+1)(k+1) \\ &= (k+1)^2 \left( \frac{k^2}{4} + (k+1) \right) \\ &= (k+1)^2 \left( \frac{k}{2} + 1 \right)^2 \\ &= (k+1)^2 \left( \frac{k+2}{2} \right)^2 \\ &= \left[ \frac{(k+1)(k+2)}{2} \right]^2 \end{aligned}$$

therefore by the First Principle of Finite Induction,  $S = \mathbb{N}$ . ■

2. If  $r \neq 1$ , show that for any positive integer  $n$ ,

$$a + ar + ar^2 + \cdots + ar^n = \frac{a(r^{n+1} - 1)}{r - 1}$$

#2. Let  $S$  be a set such that:

$$S = \left\{ n \in \mathbb{N} : a + ar + ar^2 + \cdots + ar^n = \frac{a(r^{n+1} - 1)}{r - 1} \right\}$$

$1 \in S$  for  $n = 1$  because:

$$a + ar^1 = a(r+1) = \frac{a(r+1)(r-1)}{r-1} = \frac{a(r^2-1)}{r-1}$$

Let any  $k \in \mathbb{N}$  be the member of  $S$ , that is:

$$a + ar + \cdots + ar^k = \frac{a(r^{k+1} - 1)}{r - 1}$$

Then;

$$\begin{aligned} a + ar + \cdots + ar^k + ar^{k+1} &= \frac{a(r^{k+1} - 1)}{r - 1} + ar^{k+1} \\ &= \frac{a(r^{k+1} - 1)}{r - 1} + \frac{(r-1)ar^{k+1}}{r-1} \\ &= \frac{ar^{k+2} - ar^{k+1} + ar^{k+1} - a}{r - 1} \\ &= \frac{ar^{k+2} - a}{r - 1} \\ &= \frac{a(r^{k+2} - 1)}{r - 1} \end{aligned}$$

therefore by the First Principle of Finite Induction,  $S = \mathbb{N}$ . ■

**3.** Use the Second Principle of Finite Induction to establish that for all  $n \geq 1$ ,

$$a^n - 1 = (a - 1)(a^{n-1} + a^{n-2} + \cdots + a + 1)$$

[Hint:  $a^{n+1} - 1 = (a + 1)(a^n - 1) - a(a^{n-1} - 1)$ .]

**#3.** Let  $S$  be a set such that:

$$S = \{n \in \mathbb{N} : a^n - 1 = (a - 1)(a^{n-1} + a^{n-2} + \cdots + a + 1)\}$$

$1 \in S$  for  $n = 1$  because:

$$a - 1 = (a - 1)a^0$$

Let any  $k \in \mathbb{N}$  be the member of  $S$ , that is:

$$a^k - 1 = (a - 1)(a^{k-1} + a^{k-2} + \cdots + a + 1)$$

Then;

$$\begin{aligned} (a + 1)(a^k - 1) - a(a^{k-1} - 1) &= (a + 1)(a - 1)(a^{k-1} + a^{k-2} + \cdots + a + 1) - a(a^{k-1} - 1) \\ &= (a - 1)(a + 1)(a^{k-1} + a^{k-2} + \cdots + a + 1) - a(a^{k-1} - 1) \\ &= (a - 1)[a(a^{k-1} + \cdots + a + 1) + (a^{k-1} + \cdots + a + 1)] - a(a^{k-1} - 1) \\ &= (a - 1)[a^k + 2(a^{k-1} + \cdots + a) + 1] - a(a^{k-1} - 1) \\ &= (a - 1)[a^k + 2(a^{k-1} + \cdots + a) + 1] - a^k + a \end{aligned}$$

since

$$a^k = (a - 1)(a^{k-1} + a^{k-2} + \cdots + a) + a$$

by assumption;

$$\begin{aligned} (a - 1)[a^k + 2(a^{k-1} + \cdots + a) + 1] - a^k + a &= (a - 1)(a^k + a^{k-1} + \cdots + a + 1) - a + a \\ &= (a - 1)(a^k + a^{k-1} + \cdots + a + 1) \end{aligned}$$

and by the Second Principle of Finite Induction,  $S = \mathbb{N}$ . ■

**4.** Prove that the cube of any integer can be written as the difference of two squares. Notice that

$$n^3 = (1^3 + 2^3 + \cdots + n^3) - (1^3 + 2^3 + \cdots + (n - 1)^3).$$

**#4.** By the result of **#1(e)**, for all  $n \in \mathbb{N}$ ;

$$1^3 + 2^3 + 3^3 + \cdots + n^3 = \left[ \frac{n(n + 1)}{2} \right]^2$$

and since;

$$\begin{aligned} n^3 &= (1^3 + 2^3 + \cdots + n^3) - (1^3 + 2^3 + \cdots + (n - 1)^3) \\ &= \left[ \frac{n(n + 1)}{2} \right]^2 - \left[ \frac{(n - 1)(n - 2)}{2} \right]^2 \end{aligned}$$

the cube of any integer  $n$  can be written as the difference of two squares. ■

**5.**



- (a) Find the values of  $n \leq 7$  for which  $n! + 1$  is a perfect square.
- (b) True or false? For positive integers  $m$  and  $n$ ,  $(mn)! = m!n!$  and  $(m+n)! = m! + n!$ .

#5.

- (a)  $n = 4, 5, 7$ .
- (b) False by counter examples such as  $m = 2, n = 3$  where

$$(mn)! = 6! = 6 \cdot 5 \cdot \dots \cdot 1 = 720 \neq 12 = 2 \cdot 3 \cdot 2 \cdot 1 = m!n!$$

■

## 1.2 The Binomial Theorem