# Table of Contents

1. Composition and Structure of the Atmosphere

2. Constituents — Major Gases

3. Variable Gases, Greenhouse Gases & Dry/Saturated Air

4. Carbon Dioxide ($CO_2$) & Ozone ($O_3$)

5. Vertical Distribution of Air Mass

6. Thermal Structure of the Atmosphere

7. Atmospheric Layers in Detail

8. International Standard Atmosphere (ISA)

9. Jet Standard Atmosphere (JSA)

10. Quick Revision Summary

11. Practice Questions & Answers (Q1–Q45)

12. Master Reference Tables

GHOST AVIATOR

GHOST AVIATOR


Join live classes for guided practice and doubt clearing.

# 1. Composition and Structure of the Atmosphere

**📘 Definition — Atmosphere**

The atmosphere of the earth is an **envelope of homogeneous mixture of gases**, called **Air**. It surrounds the earth and is attached to it due to gravitation. It moves with the earth at the same speed and direction.

- Total thickness: approximately `480 km`
- `80%` of the atmosphere is within `16 km`
- No exact upper boundary — gradually merges with outer space




Composition of dry air near the surface

Join live classes for guided practice and doubt clearing.

### Characteristics of the Atmosphere

| Property | Detail |
|---|---|
| Weight | Has weight and hence exerts pressure |
| Compressibility | Compressible and expandable |
| Shape | Occupies space, has no definite shape |
| Mobility | Mobile — allows transfer of heat and moisture |
| Conductivity | **Poor conductor of heat and electricity** |

## 2. Constituents — Major Gases

| Gas | Symbol | % by Volume |
|---|---|---|
| Nitrogen | $N_2$ | `78.09%` |
| Oxygen | $O_2$ | `20.95%` |
| Argon | Ar | `0.93%` |
| Carbon Dioxide | $CO_2$ | `0.035%` |

### Trace Gases

Neon, Helium, Methane, Krypton, Xenon, Nitrous Oxide, Iodine, $H_2$, $O_3$ ( `0.000007%` ), Ammonia, Carbon Monoxide, $CO_2$, SO, Nitrogen Dioxide. The atmosphere also contains water vapour (WV) and solid particles.

> ☆ **$N_2$ : $O_2$ Ratio — Exam Favourite**
>
> **By Volume:** $N_2$ : $O_2$ = `4 : 1`
>
> **By Weight:** $N_2$ : $O_2$ = `3 : 1`
>
> Mnemonic: *"Volume 4, Weight 3 — N before O"*

> ⚠ **Supplementary Oxygen Required**
>
> Due to rapid reduction of gases with height, **supplementary oxygen is needed above** `10,000 ft` .

Join live classes for guided practice and doubt clearing.

### Homosphere vs Heterosphere

| Layer | Height | Characteristic |
|---|---|---|
| **Homosphere** | Up to `~80 km` | Nearly uniform composition; well-mixed |
| **Heterosphere** | Above `80 km` | Gases separate by molecular weight; composition varies |

## 3. Variable Gases, Greenhouse Gases & Dry/Saturated Air

### Variable Gases

The gases like Water Vapour (WV), Carbon Mono Oxide, Sulphur di Oxide, Nitrogen di Oxide, and Methane **vary in amount from place to place**, being concentrated more in industrial areas, cities and water bodies, than in open areas. Though very small in quantity, these gases are very significant for weather and life.

### Greenhouse Gases

> 📘 **Greenhouse Effect**
>
> Water vapour, $CO_2$, $O_3$, Methane and some other gases are **transparent to solar (Short Wave) radiation** but **partially absorb terrestrial (Long Wave) radiation**, and re-radiate it. This keeps the earth warmer than it would have otherwise been.
>
> Greenhouse gases: **$H_2O$, $CO_2$, $O_3$, $CH_4$ (Methane)**

> ⚠️ **Global Warming Warning**
>
> Excessive amount of greenhouse gases is causing **Global Warming, which is a serious threat to our life and needs to be limited.**

### Dry Air vs Saturated Air

Water in the atmosphere can exist in three states:

- **Solid:** Snow, Ice, Hail
- **Liquid:** Drizzle, Rain, Shower
- **Gas:** Water Vapour (WV)

Join live classes for guided practice and doubt clearing.

| Concept | Detail |
|---|---|
| WV concentration | Rapidly **decreases with height** |
| Warmer air | Can hold **greater amount** of WV |
| Tropics ( `23½°N to 23½°S` ) | Air can hold up to `4%` WV by volume |
| Above `30,000 ft` | Very low WV due to very low temperatures |
| Saturated air | The water-vapour pressure equals the saturation vapour pressure at that temperature: `100% RH`. The required amount of vapour changes with temperature. |
| Unsaturated air | RH < 100%. Unsaturated air may still contain water vapour; “dry air” in composition and gas-law calculations means air without water vapour. |

## 4. Carbon Dioxide ($CO_2$) & Ozone ($O_3$)

### Carbon Dioxide ($CO_2$)

- Produced by burning of fuel, wood etc.
- Concentration substantial in **industrial areas**; negligible in Polar regions and higher altitudes
- Large amount of $CO_2$ is dissolved in the oceans
- Plants absorb $CO_2$, use its carbon as food and release $O_2$ into atmosphere

### Ozone ($O_3$)

**Ozone Properties**

- Forms in the **upper atmosphere**
- Absorbs **Ultra Violet (UV) radiation** from the Sun → raises temperature
- The ozone distribution depends on chemical formation, destruction and transport by atmospheric circulation.
- Maximum $O_3$ concentration: `20–25 km`
- $O_3$ found between `10–50 km`

Join live classes for guided practice and doubt clearing.

✅ **Why Ozone Matters for Aviation & Life**

**Ozone protects us from the ill effects (like skin cancer, sunburn and cataracts) of UV radiation.**

### Ozone Hole

⚠️ **Ozone Depletion**

In recent times the $O_3$ layer has **thinned** and large holes have developed mainly over the Polar and other latitudes due to:

- Atmospheric pollution
- Excessive use of **halocarbon refrigerants, solvents, chlorofluoro carbons**
- On reaching Stratosphere they release halogen atoms through **photodissociation** → breaks down $O_3$ into $O_2$

### Solid Particles (Aerosols)

Particles such as salt from evaporating sea water, dust from arid regions, industrial particles:

- Obscure solar radiation and affect visibility and temperature
- Act as **condensation nuclei** → promote condensation of WV prematurely
- In larger cities/industrial areas: frequent **fog and smog** (mixture of fog and smoke) due to high concentration of particles

## 5. Vertical Distribution of Air Mass

| Height | Fraction of Atmosphere Contained Below |
|---|---|
| `6 km` | `1/2` (50%) |
| `10 km` | `3/4` (75%) |
| `35 km` | `99%` |

☆ **Memory Aid**

**"6–10–35 = Half–3/4–All"**


Half the atmosphere below 6 km; ¾ below 10 km; 99% below 35 km.

Join live classes for guided practice and doubt clearing.

# 6. Thermal Structure of the Atmosphere

### 📘 How the Atmosphere is Heated

Solar radiation heats the earth and the earth in turn heats up the atmosphere by **conduction, convection, radiation** and release of latent heat by condensation of water vapour. Hence **temperature falls with height**, up to a certain height. **The atmosphere is thus heated from below and not from above.**




Temperature through the layers of the atmosphere


## Heat Flow from Earth's Surface

| Type of Heat | Mechanism | Percentage |
|---|---|---|
| **Sensible Heat** | Conduction, Convection, Radiation | 23% |
| **Latent Heat** | Evaporation, Condensation, Sublimation | 77% |

Join live classes for guided practice and doubt clearing.


> ☆ **Exam Tip**
>
> Latent heat transfer is **dominant** at 77%. Sensible heat = 23%. Mnemonic: *"Latent is Larger (77)"*

## Thermal Structure — Visual Diagram





Join live classes for guided practice and doubt clearing.

## 7. Atmospheric Layers in Detail




Tropopause height is highest over the tropics, lower in winter and higher in summer




The atmosphere in layers, with the temperature profile beside it

Join live classes for guided practice and doubt clearing.

## A. Troposphere

### 📘 Troposphere

- Lowest portion of the atmosphere
- Extends from surface to `16–18 km` at equator and `8–10 km` at poles
- Higher height at equator due to **rising hot air convection currents**
- Temperature falls at `~6.5°C/km` — called the **Lapse Rate**
- All kinds of weather occur here; turbulent mixing
- Contains `75%` of atmospheric mass and `99%` of water vapour and aerosols

| Sub-Layer | Height Range |
|---|---|
| Lower Troposphere | Surface to `2.1 km` |
| Mid Troposphere | `2.1 km` to `7.6 km` |
| Upper Troposphere | `7.6 km` to Tropopause |

## B. Tropopause

### 📘 Tropopause

- Top of the troposphere; marks boundary between Troposphere and Stratosphere
- At tropopause: lapse rate reduces to `1–2°C/km` or temperature stops falling
- Controlled by: Surface Temperature, Latitude, Season, Land-Sea distribution, Synoptic Situation
- Temperature at equator: `-70° to -75°C`
- Temperature at poles: `-40° to -45°C`
- At lower height in **Winters** than Summers
- **Warmer the surface → Higher the tropopause**

### ☆ Tropopause Breaks/Folds

There are two regions where tropopause **abruptly changes height**, called **breaks/folds**:

- At about `40°` Lat and `60°` Lat
- Break at `40°` Lat is more prominent
- **Jet streams occur at these breaks**

Join live classes for guided practice and doubt clearing.

| Type | Pressure Level | Latitude Range | Height (India) | Temp (India) | Associated Jet |
|---|---|---|---|---|---|
| **Polar Tropopause** | `~300 hPa` | Lat. 45°–60° polewards | – | – | Polar Front Jet (southern end) |
| **Middle Tropopause** | `~200 hPa` | Between Polar & Tropical | ~11.5 km (winters ~23°N) | –45°C | – |
| **Tropical Tropopause** | `~100 hPa` | Lat. 35°–40° | ~11.5 km (winters ~23°N) | –45°C | Subtropical Jet (STJ) at southern end |

✅ **Importance of Tropopause for Pilots**

- Maximum height clouds can reach
- Jet streams and maximum wind speeds occur **just below it**
- **Clear Air Turbulence (CAT)** occurs just below it

## C. Stratosphere

🟦 **Stratosphere (Tropopause to ~50 km)**

- Initially (8–10 km above tropopause) **isothermal**
- Temperature then increases slowly and towards the top sharply
- **Inversion produced by absorption of UV radiation from sun by ozone**
- Ozone layer found at height of `~25 km`
- Very **dry and stable** region — low humidity, no weather
- Above stormy weather; steady, strong, horizontal winds
- **Nacreous Clouds** (Mother of Pearl Clouds) sometimes seen in higher latitudes in winters

**Stratopause:** Upper boundary of the Stratosphere.

Join live classes for guided practice and doubt clearing.

## D. Mesosphere

📘 **Mesosphere (Stratopause to ~80 km)**

- Extends `50–80 km` from Stratopause
- Lack of solar radiation absorption + weak vertical mixing → temperatures **fall with height**
- **Noctilucent Clouds** seen occasionally in Polar regions

**Mesopause:** Top of Mesosphere. Temperatures stop falling. Coldest area of atmosphere. Temperature at 80 km = `-100°C` ( `173.15 K` ). Meteors burn up in this layer.

## E. Thermosphere

GHOST AVIATOR

📘 **Thermosphere / Ionosphere (80 km+)**

- From Mesopause to end of atmosphere
- Temperatures **increase with height**
- Above `60 km` also called **Ionosphere**
- Important for **radio wave propagation**
- Serious interference in radio propagation during solar disturbances

## F. Exosphere

📘 **Exosphere (outermost layer)**

- Outermost layer extending up to `10,000 km` from top of Thermosphere
- Molecules, ozone and atoms escape into space
- Air is very thin — few particles; very active with very high temperature but far apart → very little heat present in this layer

GHOST AVIATOR

### Atmospheric Layers — Process Flowchart

```
flowchart TD
A["🌍 Earth's Surface\n(0 km)"] --> B["TROPOSPHERE\n0–18 km (equator), 8–10 km (poles)\nTemp ↓ at 6.5°C/km\nAll weather here | 75% mass | 99% WV"]
B --> C{"TROPOPAUSE\nTemp stops falling\nJet streams & CAT below it"}
C --> D["STRATOSPHERE\n~18–50 km\nOzone at 25 km absorbs UV\nTemp ↑ (inversion)\nStable, dry, no weather"]
D --> E{"STRATOPAUSE"}
E --> F["MESOSPHERE\n~50–80 km\nTemp ↓ with height\nNoctilucent clouds (polar)\nMeteors burn here"]
F --> G{"MESOPAUSE\n−100°C / 173 K\nColdest point"}
G --> H["THERMOSPHERE / IONOSPHERE\n80 km+\nTemp ↑ with height\nRadio wave propagation"]
H --> I["EXOSPHERE\nTo 10,000 km\nMolecules escape to space"]
style A fill:#90caf9
style B fill:#64b5f6
style C fill:#fff9c4
style D fill:#a5d6a7
style E fill:#fff9c4
style F fill:#ffcc80
style G fill:#fff9c4
style H fill:#ef9a9a
style I fill:#ce93d8
```

## 8. International Standard Atmosphere (ISA)

> **📘 What is ISA?**
>
> A standard average atmosphere specified for various purposes like design and testing of aircraft, evaluation of aircraft performance, calibration of altimeters. The most widely used is the **ICAO ISA (International Standard Atmosphere)**.

GHOST AVIATOR

## ISA Specifications

| Parameter | Value |
|---|---|
| Air Condition | `Dry` |
| Temperature at MSL | `15°C (288.15 K)` |
| Pressure at MSL | `1013.25 hPa` |
| Density at MSL | `1225 g/m³` |
| Acceleration due to gravity | `980.665 cm/s²` |
| Lapse Rate (up to 11 km / 36,090 ft) | `6.5°C/km (1.98°C/1000 ft ≈ 2°C/1000 ft)` |
| Temperature 11–20 km (up to 65,617 ft) | `-56.5°C` (constant / isothermal) |
| Temperature 20–30 km | Rises at `1°C/km` with temp `-44.5°C` at 32 km (104,987 ft) |

### ISA Deviation Formula

**ISA Deviation = Actual Temperature – ISA Temperature**

**Worked Example 1:**
Actual temp at 2 km = 05°C
ISA temp at 2 km = 15 – (6.5 × 2) = 15 – 13 = 2°C
ISA Deviation = 05 – 02 = **+03°C**

**Worked Example 2:**
Pressure at MSL = 1002.25 hPa
ISA Pressure at MSL = 1013.25 hPa
ISA Deviation = 1002.25 – 1013.25 = **-11 hPa**

**Worked Example 3:**
Actual temp at 19 km = -60°C
ISA temp at 19 km = -56.5°C (constant 11–20 km)
ISA Deviation = -60 – (-56.5) = **-3.5°C** → answer -03.5°C

GHOST AVIATOR

> ⭐ **Key ISA Numbers to Remember**
>
> **15-25-56.5-36090**
>
> MSL Temp = 15°C | MSL Pressure = 1013.25 hPa | Isothermal at –56.5°C | Isothermal starts at 36,090 ft (11 km)

## 9. Jet Standard Atmosphere (JSA)

> 📘 **Why JSA?**
>
> ISA values are satisfactory for aircraft operating below `30,000 ft`, but NOT above. For space flights and very high altitude flying and engine manufacturing, additional specific values are required. JSA was introduced to meet these requirements.

| Parameter | ISA | JSA |
|---|---|---|
| MSL Temperature | `15°C (288.15 K)` | `15°C (288.15 K)` |
| Lapse Rate | `6.5°C/km (1.98°C/1000 ft)` | `2°C/1000 ft` |
| Tropopause | Yes (at 11 km) | **No Tropopause** |

GHOST AVIATOR

Join live classes for guided practice and doubt clearing.

## 🚀 Quick Revision Summary — Chapter 1: Atmosphere

| Key Fact | Value |
|---|---|
| Total atmosphere thickness | ~480 km |
| 80% atmosphere within | 16 km |
| $N_2$ : $O_2$ by Volume | 4:1 \| By Weight: 3:1 |
| Supplementary $O_2$ above | 10,000 ft |
| Homosphere up to | ~80 km |
| Max WV (tropics) | 4% by volume |
| Saturated Air RH | 100% |
| Ozone max concentration | 20–25 km |
| Ozone range | 10–50 km |
| Sensible heat | 23% |
| Latent heat | 77% |
| Troposphere height (equator) | 16–18 km |
| Troposphere height (poles) | 8–10 km |
| Lapse rate | ~6.5°C/km |
| WV + aerosols in troposphere | 99% |
| Atmospheric mass in troposphere | 75% |
| Tropopause temp (equator) | –70° to –75°C |
| Tropopause temp (poles) | –40° to –45°C |
| Stratosphere to | ~50 km |
| Ozone layer height | ~25 km |
| Mesosphere range | 50–80 km |
| Mesopause temperature | –100°C (173.15 K) |

GHOST AVIATOR

Join live classes for guided practice and doubt clearing.

| | |
|---|---|
| Ionosphere above | 60 km |
| Exosphere up to | 10,000 km |
| ISA MSL Temp | 15°C (288.15 K) |
| ISA MSL Pressure | 1013.25 hPa |
| ISA MSL Density | 1225 g/m³ |
| ISA Lapse Rate | 6.5°C/km up to 11 km |
| ISA Isothermal zone | –56.5°C, 11–20 km |
| JSA Lapse Rate | 2°C/1000 ft (no tropopause) |

GHOST AVIATOR

GHOST AVIATOR

Join live classes for guided practice and doubt clearing.

## Practice Questions & Answers

Q1

**Lowest layer of atmosphere is …… (a) Troposphere (b) Tropopause (c) Stratosphere**

✅ CORRECT ANSWER: (A) TROPOSPHERE

The Troposphere is the lowest layer, starting at earth's surface and extending to the tropopause. All weather occurs here.

DISTRACTOR ANALYSIS

(b) Tropopause is a boundary, not a layer. (c) Stratosphere is above the Troposphere.

📌 *INSTRUCTOR'S NOTE*

*Remember the order: **T-S-M-T-E** = Troposphere, Stratosphere, Mesosphere, Thermosphere, Exosphere from bottom to top.*

Q2

**Height of Tropopause at equator is …… (a) 8–10 km (b) 16–18 km (c) 14 km**

✅ CORRECT ANSWER: (B) 16–18 KM

At the equator, rising hot air due to convection pushes the tropopause higher — up to 16–18 km.

DISTRACTOR ANALYSIS

(a) 8–10 km is the tropopause height at the poles. (c) 14 km is intermediate but incorrect.

📌 *INSTRUCTOR'S NOTE*

*Equator = Hot = Higher tropopause. Poles = Cold = Lower tropopause. "Hotter higher, colder lower."*

Q3

**Height of Tropopause at Poles is …… (a) 12 km (b) 13 km (c) 08 km**

✅ CORRECT ANSWER: (C) 08 KM

At the poles, the atmosphere is colder; therefore convective activity is minimal and the tropopause is lower, around 8–10 km.

DISTRACTOR ANALYSIS

Join live classes for guided practice and doubt clearing.

(a) and (b) are too high for polar regions.

📌 *INSTRUCTOR'S NOTE*

*Polar tropopause ~300 hPa; Tropical tropopause ~100 hPa (lower pressure = higher altitude).*

Q4

**Higher the surface temperature …… would be the tropopause. (a) Higher (b) Lower (c) Same**

✅ CORRECT ANSWER: (A) HIGHER

Warmer surfaces generate more convective activity, pushing the tropopause to a greater height.

DISTRACTOR ANALYSIS

(b) Lower — incorrect; this is the effect at colder surfaces/poles. (c) Same — temperature definitely affects tropopause height.

GHOST AVIATOR

📌 *INSTRUCTOR'S NOTE*

*"Warmer surface = Higher tropopause" — think of hot air balloons rising higher when it's warmer.*

Q5

**Height of tropopause …… (a) Is constant (b) Varies with altitude (c) Varies with Latitude**

✅ CORRECT ANSWER: (C) VARIES WITH LATITUDE

Tropopause height depends on latitude (equator ~16–18 km; poles ~8 km) and also on season, surface temperature, synoptic situation.

DISTRACTOR ANALYSIS

(a) Is constant — definitely not; it varies significantly. (b) Varies with altitude — altitude is not the controlling factor.

GHOST AVIATOR

📌 *INSTRUCTOR'S NOTE*

*Tropopause height is controlled by: Surface Temp, Latitude, Season, Land-Sea distribution, Synoptic situation.*

Q6

**Above 8 km the lower temperatures are over …… (a) Equator (b) Mid Latitudes (c) Poles**

✅ Answer: (a) Equator

Join live classes for guided practice and doubt clearing.

Above the low polar tropopause (about 8 km) the polar air is already in the stratosphere, where temperature stops falling and begins to rise. Over the Equator the troposphere is still 16–18 km deep and keeps cooling, reaching about −75 to −80 °C at its top, against −40 to −50 °C at the Poles. So above 8 km it is colder over the Equator.

❌ (b) Mid-latitudes lie between the two extremes. (c) Poles: above 8 km the polar air is the warmer of the two.

💡 *At the surface the Equator is warmest; at tropopause level the Equator is coldest. The picture turns over at about 8 km.*

Q7

**Atmosphere is heated by …… (a) Solar Radiation (b) Heat from earth surface (c) From above**

GHOST AVIATOR

✅ CORRECT ANSWER: (B) HEAT FROM EARTH SURFACE

The atmosphere is heated from below by the earth's surface (via conduction, convection, radiation and latent heat). The atmosphere is a poor absorber of direct solar radiation.

DISTRACTOR ANALYSIS

(a) Solar radiation heats the EARTH, not directly the lower atmosphere. (c) From above is incorrect — atmosphere is heated from below.

📌 *INSTRUCTOR'S NOTE*

*Remember: **Temperature falls with height** in troposphere BECAUSE heating is from below.*

Q8

**The temperature at 80 km is …… (a) 173 K (b) 100 K (c) –137°C**

GHOST AVIATOR

✅ CORRECT ANSWER: (A) 173 K (= –100°C)

The text explicitly states: "The temperature at 80 km is about –100°C (173.15 K)." 173 K ≈ –100°C. This is the Mesopause, the coldest point in the atmosphere.

DISTRACTOR ANALYSIS

(b) 100 K = –173°C, far too cold. (c) –137°C = 136 K, not matching the textbook value.

📌 *INSTRUCTOR'S NOTE*

*Mesopause = –100°C = 173.15 K. Mnemonic: "80 km → 173 K → Meteors burn".*

Q9

Join live classes for guided practice and doubt clearing.

**$CO_2$ and $H_2O$ are also called …… (a) Green House Gases (b) Rare Earth Gases (c) [other]**

✅ **CORRECT ANSWER: (A) GREEN HOUSE GASES**

$CO_2$ and $H_2O$ (water vapour) along with $O_3$ and Methane are greenhouse gases — they are transparent to shortwave solar radiation but absorb longwave terrestrial radiation.

**DISTRACTOR ANALYSIS**

(b) Rare Earth Gases (Noble gases like Argon, Neon, Helium) are a different category entirely.

📌 ***INSTRUCTOR'S NOTE***

*Greenhouse gases: **W-C-O-M** = Water vapour, $CO_2$, $O_3$, Methane.*

**Q10**

**Troposphere is generally …… (a) Stable (b) Unstable (c) Neutral**

✅ **CORRECT ANSWER: (B) UNSTABLE**

The Troposphere has a lapse rate of ~6.5°C/km. With turbulent mixing, convective activity and all weather occurring here, it is generally unstable.

**DISTRACTOR ANALYSIS**

(a) Stable — that's the Stratosphere. (c) Neutral — not the defining characteristic.

📌 ***INSTRUCTOR'S NOTE***

***Troposphere = Unstable; Stratosphere = Stable**. Easy contrast for exams.*

**Q11**

**Stratosphere is …… (a) Unstable (b) Neutral (c) Stable**

✅ **CORRECT ANSWER: (C) STABLE**

The Stratosphere has a temperature inversion (temperature increases with height due to ozone absorbing UV), making it very stable with no convective weather.

**DISTRACTOR ANALYSIS**

(a) Unstable — that's the Troposphere. (b) Neutral — not the dominant characteristic.

📌 ***INSTRUCTOR'S NOTE***

*Stratosphere = "Stratified" = Layered = Stable. Also: no weather, steady winds.*

Join live classes for guided practice and doubt clearing.

Q12

**Tropopause is discontinuous at about …… (a) 30°lat (b) 40°lat (c) 80°lat**

✅ CORRECT ANSWER: (B) 40°LAT

The breaks/folds in the tropopause occur at ~40° and ~60° latitude, with the break at 40°lat being more prominent. Jet streams occur at these breaks.

DISTRACTOR ANALYSIS

(a) 30°lat — not a significant break. (c) 80°lat — too poleward, not where the prominent break is.

📌 *INSTRUCTOR'S NOTE*

*"Break at 40, Fold at 60." Jet streams at these breaks — important for upper-level wind planning.*

GHOST AVIATOR

Q13

**Most of atmospheric mass is contained in …… (a) Troposphere (b) Stratosphere (c) Heterosphere**

✅ CORRECT ANSWER: (A) TROPOSPHERE

The Troposphere contains 75% of the atmospheric mass and 99% of water vapour and aerosols.

DISTRACTOR ANALYSIS

(b) Stratosphere is relatively thin in mass. (c) Heterosphere is the outer part with very low density.

📌 *INSTRUCTOR'S NOTE*

*Troposphere = **75% mass, 99% WV & aerosols**. Two key numbers.*

Q14

**Stratosphere extends from Tropopause to …… (a) 50 km (b) 60 km (c) 40 km**

GHOST AVIATOR

✅ CORRECT ANSWER: (A) 50 KM

The Stratosphere extends up to ~50 km from the earth's surface (approximately 30–32 km above the tropopause at equator).

DISTRACTOR ANALYSIS

(b) 60 km is too high — enters Mesosphere region. (c) 40 km is not the correct upper boundary.

📌 *INSTRUCTOR'S NOTE*

*Layer tops: Tropo ~18, Strato ~50, Meso ~80. Easy: **18–50–80**.*

Q15

**The middle atmosphere layer with temperature inversion and stability is …… (a) Troposphere (b) Tropopause (c) Stratosphere**

✅ CORRECT ANSWER: (C) STRATOSPHERE

The Stratosphere has a temperature inversion (UV absorption by ozone warms upper layers) making it very stable. It lies between the Tropopause and Stratopause.

DISTRACTOR ANALYSIS

(a) Troposphere is unstable with a normal lapse rate. (b) Tropopause is a boundary, not a layer.

📌 *INSTRUCTOR'S NOTE*

*Stratosphere = Stable + Inversion = Ozone = UV absorption. All linked.*

Q16

**Mother of Pearl clouds occur in …… (a) Mesosphere (b) Thermosphere (c) Stratosphere**

✅ CORRECT ANSWER: (C) STRATOSPHERE

Nacreous Clouds (Mother of Pearl Clouds) are sometimes seen in the Stratosphere in higher latitudes in winters.

DISTRACTOR ANALYSIS

(a) Mesosphere — Noctilucent clouds occur here. (b) Thermosphere — no clouds form here.

📌 *INSTRUCTOR'S NOTE*

*Nacreous = Stratosphere | Noctilucent = Mesosphere. "Na-Stra, Noc-Meso".*

Q17

**The temperature in ISA at 17 km is …… (a) –56.5°C (b) –65.5°C (c) –35.5°C**

✅ CORRECT ANSWER: (A) –56.5°C

In ISA, from 11 km to 20 km (65,617 ft), temperature is assumed constant at **–56.5°C**. 17 km falls within this isothermal zone.

DISTRACTOR ANALYSIS

(b) –65.5°C and (c) –35.5°C are both incorrect — no lapse rate change occurs in this zone.

📌 *INSTRUCTOR'S NOTE*

Join live classes for guided practice and doubt clearing.

*ISA isothermal zone: 11 km to 20 km = –56.5°C throughout. Any question with height 11–20 km → answer is –56.5°C.*

Q18

**By weight, approximate ratio of $O_2$ to $N_2$ in the atmosphere is …… (a) 1:3 (b) 1:4 (c) 1:5**

✅ CORRECT ANSWER: (A) 1:3 (I.E., $O_2$ : $N_2$ = 1:3 BY WEIGHT)

$N_2$ : $O_2$ by weight = 3:1, so $O_2$ : $N_2$ by weight = 1:3.

DISTRACTOR ANALYSIS

(b) 1:4 — that is the VOLUME ratio ($O_2$:$N_2$ = 1:4). Don't confuse weight and volume ratios.

📌 *INSTRUCTOR'S NOTE*

***Volume: 4:1* | *Weight: 3:1*** *($N_2$:$O_2$). Volume ratio is larger because $N_2$ is less dense per unit volume.*

Q19

**By volume, the approximate ratio of $O_2$ to $N_2$ in the atmosphere is …… (a) 1:3 (b) 1:4 (c) 1:5**

✅ CORRECT ANSWER: (B) 1:4 ($O_2$ : $N_2$ = 1:4 BY VOLUME)

$N_2$ : $O_2$ by volume = 4:1, therefore $O_2$ : $N_2$ = 1:4. $N_2$ = 78.09%, $O_2$ = 20.95% — ratio is almost 4:1.

DISTRACTOR ANALYSIS

(a) 1:3 is the weight ratio. (c) 1:5 is incorrect.

📌 *INSTRUCTOR'S NOTE*

*Volume ratio is 4:1 (larger). Weight ratio is 3:1 (smaller). "Volume is Vast (4)."*

Q20

**By volume, the proportion of $CO_2$ in the atmosphere is …… (a) 3% (b) 0.3% (c) 0.03%**

✅ CORRECT ANSWER: (C) 0.03% (0.035%)

$CO_2$ constitutes approximately 0.035% by volume in the atmosphere.

DISTRACTOR ANALYSIS

(a) 3% — vastly overstated. (b) 0.3% — 10 times too high.

📌 *INSTRUCTOR'S NOTE*

Join live classes for guided practice and doubt clearing.

*$CO_2$ = 0.035% ≈ 0.03% — tiny amount but significant for greenhouse effect.*

Q21

**In ISA, the mean sea level temperature is …… (a) 15°C (b) 10°C (c) 25°C**

✅ **CORRECT ANSWER: (A) 15°C**

ISA MSL temperature = 15°C (288.15 K). This is a fundamental ISA value.

**DISTRACTOR ANALYSIS**

(b) 10°C — incorrect. (c) 25°C — too warm, not ISA standard.

📌 ***INSTRUCTOR'S NOTE***

*ISA MSL: Temp = **15°C**, Pressure = **1013.25 hPa**, Density = **1225 g/m³**. Memorise all three together.*

GHOST AVIATOR

Q22

**Maximum concentration of ozone is at a height of …… (a) 10–15 km (b) 20–25 km (c) 30–35 km**

✅ **CORRECT ANSWER: (B) 20–25 KM**

Ozone is found between 10–50 km with maximum concentration at 20–25 km in the lower Stratosphere.

**DISTRACTOR ANALYSIS**

(a) 10–15 km — lower end of ozone range, not maximum. (c) 30–35 km — past the maximum.

📌 ***INSTRUCTOR'S NOTE***

*Ozone range: 10–50 km | Max: 20–25 km | Ozone layer: ~25 km. "20-25 is Ozone's favourite zone."*

GHOST AVIATOR

Q23

**Additional oxygen is needed while flying above …… (a) 5000 ft (b) 7000 ft (c) 10000 ft**

✅ **CORRECT ANSWER: (C) 10000 FT**

Due to rapid reduction of gases with height, supplementary oxygen is needed above **10,000 ft**.

**DISTRACTOR ANALYSIS**

(a) 5000 ft and (b) 7000 ft — below the threshold requiring supplemental oxygen.

📌 ***INSTRUCTOR'S NOTE***

*10,000 ft = critical altitude for supplemental $O_2$. Also relevant for cabin pressurization regulations.*

Join live classes for guided practice and doubt clearing.

Q24

**$CO_2$ and $H_2O$ keep the atmosphere …… (a) Warm (b) Cold (c) Have no effect**

✅ CORRECT ANSWER: (A) WARM

$CO_2$ and $H_2O$ are greenhouse gases that absorb terrestrial longwave radiation and re-radiate it, keeping the earth warmer than it would otherwise be.

DISTRACTOR ANALYSIS

(b) Cold — incorrect, they trap heat. (c) Have no effect — incorrect, they are the primary drivers of the greenhouse effect.

📌 *INSTRUCTOR'S NOTE*

*Think of greenhouse gases as a "blanket" around the earth, trapping warmth.*

GHOST AVIATOR

Q25

**Noctilucent clouds occur in …… (a) Thermosphere (b) Mesosphere (c) Stratosphere**

✅ CORRECT ANSWER: (B) MESOSPHERE

Noctilucent Clouds are seen on rare occasions in the upper Mesosphere, in Polar regions.

DISTRACTOR ANALYSIS

(a) Thermosphere — no clouds form there. (c) Stratosphere — Nacreous/Mother of Pearl clouds occur here, not Noctilucent.

📌 *INSTRUCTOR'S NOTE*

***Noctilucent = Mesosphere (Polar) | Nacreous = Stratosphere (higher latitudes, winters)**. Know both!*

GHOST AVIATOR

Q26

**Temperature at 2 km is 05°C. What is ISA deviation? Hint: (Actual – ISA) (a) –05°C (b) –02°C (c) 03°C**

✅ CORRECT ANSWER: (C) 03°C (I.E. +03°C)

ISA temp at 2 km = 15 – (6.5 × 2) = 15 – 13 = 2°C. ISA Deviation = 5 – 2 = **+3°C**.

DISTRACTOR ANALYSIS

(a) –5°C wrong sign and magnitude. (b) –2°C — wrong direction entirely.

📌 *INSTRUCTOR'S NOTE*

Join live classes for guided practice and doubt clearing.

*ISA Deviation = Actual – ISA. Positive deviation = warmer than ISA. Always compute ISA temp first using lapse rate.*

Q27

**Pressure at MSL is 1002.25 hPa. Find the ISA deviation. Hint: (Actual – ISA) (a) –11 hPa (b) 10 hPa (c) 12 hPa**

✅ **CORRECT ANSWER: (A) –11 HPA**

ISA MSL Pressure = 1013.25 hPa. Deviation = 1002.25 – 1013.25 = –11 hPa. Negative means pressure is lower than ISA standard.

**DISTRACTOR ANALYSIS**

(b) 10 hPa and (c) 12 hPa — wrong sign and magnitude.

GHOST AVIATOR

📌 ***INSTRUCTOR'S NOTE***

*Same formula applies for pressure deviation as temperature deviation. ISA MSL Pressure = 1013.25 hPa.*

Q28

**In actual atmosphere temp. at 19 km is –60°C. Find the ISA deviation. Hint: (Actual – ISA) (a) –4.5°C (b) –05.5°C (c) –03.5°C**

✅ **CORRECT ANSWER: (C) –03.5°C**

ISA temp at 19 km = –56.5°C (isothermal zone 11–20 km). Deviation = –60 – (–56.5) = –60 + 56.5 = **–3.5°C**.

**DISTRACTOR ANALYSIS**

(a) –4.5°C and (b) –5.5°C are arithmetic errors — make sure signs are handled correctly.

GHOST AVIATOR

📌 ***INSTRUCTOR'S NOTE***

*For any altitude between 11–20 km: ISA temp is always –56.5°C. Don't try to apply lapse rate in this zone.*

Q29

**Nacreous clouds occur in …… (a) Thermosphere (b) Mesosphere (c) Upper Stratosphere**

✅ **CORRECT ANSWER: (C) UPPER STRATOSPHERE (STRATOSPHERE)**

Nacreous Clouds (Mother of Pearl Clouds) are sometimes seen in the Stratosphere in higher latitudes in winters.

**DISTRACTOR ANALYSIS**

Join live classes for guided practice and doubt clearing.

(a) Thermosphere — no clouds. (b) Mesosphere — Noctilucent clouds.

📌 *INSTRUCTOR'S NOTE*

*Nacreous clouds are iridescent and beautiful — found in the stratosphere over polar regions in winter.*

Q30

**The atmosphere up to 80 km has a nearly similar composition and is called the Homosphere. Its uniform composition is due to …… (a) Pressure (b) Gravitation of earth (c) Mixing due to turbulence**

✅ CORRECT ANSWER: (C) MIXING DUE TO TURBULENCE

The atmosphere is generally well mixed up to ~80 km due to turbulent mixing, maintaining a nearly uniform composition.

GHOST AVIATOR

DISTRACTOR ANALYSIS

(a) Pressure alone doesn't cause uniform mixing. (b) Gravitation causes vertical stratification, not uniform mixing.

📌 *INSTRUCTOR'S NOTE*

*Homosphere = uniform mixing by turbulence. Heterosphere = no mixing = separation by molecular weight.*

Q31

**Half of the atmospheric air mass is contained …… below (a) 20,000 ft (b) 15,000 ft (c) 10,000 ft**

✅ Answer: (a) 20,000 ft

Half of the mass of the atmosphere lies below about 5.5–6 km, roughly 18,000–20,000 ft. Of the three options, 20,000 ft is the closest.

GHOST AVIATOR

❌ (b) 15,000 ft (about 4.6 km) and (c) 10,000 ft (about 3 km) are too low: well over half of the air is still above those levels.

💡 *Half the air is below about FL180, which is why the 500 hPa level sits near FL180.*

Q32

**In jet standard atmosphere the Lapse Rate is …… (a) 2°C/1000 ft (b) 2°C/km (c) 5°C/km**

✅ Answer: (a) 2°C/1000 ft

Join live classes for guided practice and doubt clearing.

The Jet Standard Atmosphere assumes a constant lapse rate of 2 °C per 1000 ft and has no tropopause.

❌ (b) 2 °C/km is far too small a lapse rate. (c) 5 °C/km is not a standard rate.

💡 *JSA: 2 °C/1000 ft, no tropopause. ISA: 6.5 °C/km (about 2 °C/1000 ft) up to 11 km, then isothermal.*

Q33

**The rate of fall of temperatures with height, called …… (a) Isothermal rate (b) Inversion Rate (c) Lapse Rate**

✅ CORRECT ANSWER: (C) LAPSE RATE

Lapse rate is the rate at which temperature decreases with increasing altitude, approximately 6.5°C/km in the ISA troposphere.

GHOST AVIATOR

DISTRACTOR ANALYSIS

(a) Isothermal rate — temperature does NOT change with height (isothermal). (b) Inversion Rate — temperature INCREASES with height (inverse of normal).

📌 *INSTRUCTOR'S NOTE*

*Lapse Rate = normal fall of temperature with height. Inversion = temperature rise with height. Isothermal = constant temperature.*

Q34

**In actual atmosphere the lapse rate could …… (a) Assume any value (b) Fall up to 8 km. (c) Rise up to 50 km**

✅ CORRECT ANSWER: (A) ASSUME ANY VALUE

GHOST AVIATOR

In the actual (real) atmosphere — unlike ISA — the lapse rate can assume any value; it can be positive, zero (isothermal), or negative (inversion) depending on atmospheric conditions.

DISTRACTOR ANALYSIS

(b) and (c) describe fixed behaviours. The actual atmosphere is highly variable.

📌 *INSTRUCTOR'S NOTE*

*ISA lapse rate = fixed at 6.5°C/km. ACTUAL lapse rate = variable, can be anything. This distinction is key for stability analysis.*

Q35

Join live classes for guided practice and doubt clearing.

**Tropical Tropopause extends from the equator to Lat. 35°–40°. Over India it is at …… (a) 20–21 km (b) 14–15 km (c) 16–16.5 km**

✅ Answer: (c) 16–16.5 km

The tropical tropopause lies near the 100 hPa level, roughly 16–16.5 km, and extends from the equator to about 35°–40° latitude. It is the tropopause over most of India in the summer.

❌ (a) 20–21 km is higher than any tropopause. (b) 14–15 km is lower than the ~100 hPa tropical level.

💡 *Tropical tropopause: about 100 hPa, 16 km. Polar tropopause: about 300 hPa, 8–10 km.*

Q36

**Lapse rate in the troposphere is produced by …… and in the stratosphere by …… (a) Evaporation; condensation (b) Condensation; solar radiation (c) Terrestrial radiation; solar radiation (d) Solar radiation; convection**

GHOST AVIATOR

✅ CORRECT ANSWER: (C) TERRESTRIAL RADIATION; SOLAR RADIATION

In the Troposphere: cooling with height is due to terrestrial radiation heating from below. In Stratosphere: temperature inversion is due to solar radiation (UV) being absorbed by ozone — causing warming with height.

DISTRACTOR ANALYSIS

Other combinations mix up the mechanisms. Condensation releases latent heat but is not the primary cause of the lapse rate.

📌 *INSTRUCTOR'S NOTE*

*Troposphere heated from Earth (below) → cools upward. Stratosphere heated by ozone absorbing UV (from Sun above) → warms upward.*

GHOST AVIATOR

Q37

**Most of the water vapour in the atmosphere is confined up to …… (a) stratosphere (b) 30,000 ft (c) Mid troposphere (d) lower troposphere**

✅ Answer: (b) 30,000 ft

Water vapour is held in the lowest part of the troposphere and falls off quickly with height. Above about 30,000 ft the amount is negligible, so that is the limit up to which it is confined.

❌ (a) The stratosphere holds almost no water vapour. (c) and (d): most of the vapour is in the lower

Join live classes for guided practice and doubt clearing.

troposphere, but the level asked for is the upper limit of its presence.

💡 *Water vapour: troposphere only, mostly in the lowest few km, negligible above 30,000 ft.*

Q38

**Negative lapse rate of temperature is …… (a) Isothermal rate (b) Temperature rise with lowering height (c) Temperature rise with height (d) Temperature fall with height**

✅ CORRECT ANSWER: (C) TEMPERATURE RISE WITH HEIGHT

A "negative lapse rate" means temperature increases with altitude — this is called an **inversion**. This occurs in the Stratosphere and in inversions within the Troposphere.

DISTRACTOR ANALYSIS

(a) Isothermal = zero lapse rate (no temperature change). (d) Temperature fall with height = positive lapse rate (normal). (b) is phrased incorrectly.

📌 *INSTRUCTOR'S NOTE*

*Positive lapse rate = temp falls with height (normal). Zero lapse rate = isothermal. Negative lapse rate = inversion (temp rises with height).*

Q39

**In ICAO ISA the atmosphere is assumed to be isothermal …… (a) in stratosphere (b) 11 to 16 km (c) 11 to 20 km (d) 11 to 32 km**

✅ CORRECT ANSWER: (C) 11 TO 20 KM

In ICAO ISA, temperature is assumed constant at –56.5°C from 11 km (36,090 ft) to 20 km (65,617 ft). This is the isothermal (stratospheric) zone.

DISTRACTOR ANALYSIS

(b) 11–16 km — too narrow. (d) 11–32 km — too wide; from 20 km temperature starts rising again. (a) "in stratosphere" is vague and incorrect if taken to mean the whole stratosphere.

📌 *INSTRUCTOR'S NOTE*

*ISA Isothermal: 11–20 km at –56.5°C. After 20 km temp rises. After 32 km another zone begins.*

Q40

Join live classes for guided practice and doubt clearing.

**One of the characteristics of our atmosphere is …… (a) Poor conductor of heat and electricity (b) Equator is warmer than poles above 10 km (c) Lapse rate is positive in stratosphere (d) Density is constant above 8 km**

✅ CORRECT ANSWER: (A) POOR CONDUCTOR OF HEAT AND ELECTRICITY

The atmosphere is a poor conductor of heat and electricity — one of its key characteristics listed at the beginning of the chapter.

DISTRACTOR ANALYSIS

(b) Above 10 km, poles are actually warmer than equator at the tropopause (poles –40 to –45°C vs equator –70 to –75°C). (c) Lapse rate in stratosphere is negative (inversion). (d) Density is NOT constant above 8 km.

📌 *INSTRUCTOR'S NOTE*

*Atmosphere = poor conductor. Counterintuitively, tropopause at POLES is warmer (–40°C) than EQUATOR (–75°C) — despite surface being opposite!*

GHOST AVIATOR

Q41

**Heat transfer in the atmosphere is maximum due to …… (a) convection (b) radiation (c) sensible heat (d) latent heat**

✅ CORRECT ANSWER: (D) LATENT HEAT

Latent Heat (Evaporation, Condensation, Sublimation) accounts for 77% of heat transfer from earth to atmosphere — much more than sensible heat at 23%.

DISTRACTOR ANALYSIS

(a) Convection is a mechanism within sensible heat, not the maximum overall. (b) Radiation contributes but is part of the 23%. (c) Sensible heat is only 23%.

GHOST AVIATOR

📌 *INSTRUCTOR'S NOTE*

*Latent heat = 77% (dominant!). Sensible heat = 23%. "Latent is Larger at 77%."*

Q42

**The knowledge of the height of tropopause is important for a pilot because …… (a) weather is mainly confined up to this level (b) clouds rarely reach up to this height due to jetstream (c) stratosphere starts at this height where all solar radiation is absorbed**

✅ CORRECT ANSWER: (A) WEATHER IS MAINLY CONFINED UP TO THIS LEVEL

Join live classes for guided practice and doubt clearing.

The tropopause is the maximum height clouds can reach. All weather, turbulence, cloud formations are below it. Knowing its height helps pilots plan for weather-free cruising altitudes.

**DISTRACTOR ANALYSIS**

(b) Clouds do grow to just below tropopause — not because of jet streams. (c) Solar radiation is absorbed by ozone in the stratosphere, not at the tropopause itself.

**📌 *INSTRUCTOR'S NOTE***

*Tropopause = weather ceiling. Above it = stable, clear, steady stratospheric winds. CAT and Jet streams just BELOW it.*

**Q43**

**In ISA atmosphere the tropopause occurs at a height of …… (a) 8–10 km (b) 11 km (c) 16–18 km**

GHOST AVIATOR

**✅ CORRECT ANSWER: (B) 11 KM**

In ICAO ISA, the tropopause is fixed at **11 km (36,090 ft)**, where the lapse rate stops and isothermal zone begins (–56.5°C).

**DISTRACTOR ANALYSIS**

(a) 8–10 km = actual polar tropopause. (c) 16–18 km = actual equatorial tropopause. ISA is a standard — fixed at 11 km.

**📌 *INSTRUCTOR'S NOTE***

*ISA tropopause = **11 km = 36,090 ft**. This is a fixed constant in ISA, unlike the actual variable tropopause.*

**Q44**

**Most of the transfer of heat in the atmosphere is due to …… (a) conduction (b) convection (c) Both A and B (d) latent heat**

GHOST AVIATOR

**✅ CORRECT ANSWER: (D) LATENT HEAT**

As stated in the chapter: Latent Heat transfer = 77% (dominant). Sensible heat (including conduction, convection, radiation) = 23%.

**DISTRACTOR ANALYSIS**

(a) Conduction — atmosphere is a poor conductor. (b) Convection — important but part of 23%. (c) Both — only 23% combined.

**📌 *INSTRUCTOR'S NOTE***

*This question is almost identical to Q41. Reinforce: Latent heat dominates atmospheric heat transfer.*

Join live classes for guided practice and doubt clearing.

# Master Reference Tables

## 1. All Numerical Values — Chapter 1

| Parameter | Value | Source |
|---|---|---|
| Atmosphere total thickness | ~480 km | Composition |
| 80% atmosphere below | 16 km | Composition |
| $N_2$ by volume | 78.09% | Constituents |
| $O_2$ by volume | 20.95% | Constituents |
| Argon by volume | 0.93% | Constituents |
| $CO_2$ by volume | 0.035% | Constituents |
| $O_3$ trace concentration | 0.000007% | Constituents |
| $N_2$:$O_2$ by Volume | 4:1 | Ratios |
| $N_2$:$O_2$ by Weight | 3:1 | Ratios |
| Supplemental $O_2$ required above | 10,000 ft | Physiology |
| Homosphere up to | ~80 km | Structure |
| Max WV (tropics) | 4% by volume | WV |
| WV minimal above | 30,000 ft | WV |
| Tropics latitude band | 23½°N to 23½°S | WV |
| Ozone range | 10–50 km | Ozone |
| Ozone max concentration | 20–25 km | Ozone |
| Ozone layer (Stratosphere) | ~25 km | Ozone |
| Airmass below 6 km | 1/2 (50%) | Vertical dist. |
| Airmass below 10 km | 3/4 (75%) | Vertical dist. |
| Airmass below 35 km | 99% | Vertical dist. |
| Sensible heat transfer | 23% | Thermal |

GHOST AVIATOR

Join live classes for guided practice and doubt clearing.

| | | |
|---|---|---|
| Latent heat transfer | 77% | Thermal |
| Troposphere height (equator) | 16–18 km | Layers |
| Troposphere height (poles) | 8–10 km | Layers |
| ISA Lapse rate | 6.5°C/km (≈1.98°C/1000 ft) | Layers |
| Atmospheric mass in troposphere | 75% | Layers |
| WV+aerosols in troposphere | 99% | Layers |
| Lower Troposphere | Surface – 2.1 km | Layers |
| Mid Troposphere | 2.1 – 7.6 km | Layers |
| Upper Troposphere | 7.6 km – Tropopause | Layers |
| Tropopause lapse rate | 1–2°C/km | Tropopause |
| Tropopause temp (equator) | –70° to –75°C | Tropopause |
| Tropopause temp (poles) | –40° to –45°C | Tropopause |
| Tropopause breaks at | ~40° and ~60° Lat | Tropopause |
| Polar Tropopause pressure | ~300 hPa | Tropopause |
| Tropical Tropopause pressure | ~100 hPa | Tropopause |
| Middle Tropopause pressure | ~200 hPa | Tropopause |
| Stratosphere extent | ~18–50 km | Layers |
| Mesosphere extent | 50–80 km | Layers |
| Mesopause temperature | –100°C (173.15 K) | Layers |
| Ionosphere above | 60 km | Layers |
| Exosphere extent | Up to 10,000 km | Layers |
| ISA MSL Temperature | 15°C (288.15 K) | ISA |
| ISA MSL Pressure | 1013.25 hPa | ISA |
| ISA MSL Density | 1225 g/m³ | ISA |

GHOST AVIATOR
GHOST AVIATOR

Join live classes for guided practice and doubt clearing.

| | | |
|---|---|---|
| ISA gravity | 980.665 cm/s² | ISA |
| ISA isothermal zone | 11–20 km at –56.5°C | ISA |
| ISA isothermal start | 11 km = 36,090 ft | ISA |
| ISA isothermal end | 20 km = 65,617 ft | ISA |
| ISA temp rise zone (20–30 km) | 1°C/km, –44.5°C at 32 km | ISA |
| JSA Lapse rate | 2°C/1000 ft | JSA |
| JSA valid below | 30,000 ft (ISA ok); JSA for above | JSA |

## 2. Mnemonics Summary

GHOST AVIATOR

| Mnemonic | What it Helps Remember |
|---|---|
| **T-S-M-T-E** | Atmospheric layers bottom to top: Troposphere, Stratosphere, Mesosphere, Thermosphere, Exosphere |
| **Volume 4, Weight 3** | $N_2$:$O_2$ ratio by Volume=4:1, by Weight=3:1 |
| **Latent is Larger (77%)** | Latent heat transfer = 77%, Sensible = 23% |
| **6–10–35 = Half–3/4–All** | Vertical air mass distribution |
| **Hotter = Higher (Tropopause)** | Warmer surface → higher tropopause |
| **15–1013.25–1225** | ISA MSL: Temp, Pressure, Density |
| **11 km = 36,090 ft = –56.5°C** | ISA tropopause and isothermal start |
| **Na-Stra, Noc-Meso** | Nacreous clouds = Stratosphere; Noctilucent = Mesosphere |
| **W-C-O-M** | Greenhouse gases: Water vapour, $CO_2$, $O_3$, Methane |
| **20–25 is Ozone's zone** | Maximum ozone concentration at 20–25 km |
| **ISA Dev = Actual – ISA** | Formula for ISA deviation (positive = warmer than standard) |
| **Break at 40, Fold at 60** | Tropopause discontinuities at ~40° and ~60° latitude |

Join live classes for guided practice and doubt clearing.

## 3. Formula Sheet

### Key Formulae — Chapter 1

**ISA Temperature at altitude h (km), for h ≤ 11 km:**

$T(h) = 15 - 6.5 \times h$ [°C]

Example: h = 5 km → T = 15 − 32.5 = −17.5°C

**ISA Deviation:**

ISA Deviation = T_actual − T_ISA

Positive = warmer than ISA | Negative = colder than ISA

**ISA Isothermal zone (11–20 km):**

T = −56.5°C (constant; no lapse rate applies)

**Kelvin–Celsius conversion:**

K = °C + 273.15

−56.5°C = 216.65 K | −100°C = 173.15 K | 15°C = 288.15 K

GHOST AVIATOR

## 4. Answer Key — Chapter 1 Questions (Q1–Q45)

| Q | Ans | Q | Ans | Q | Ans | Q | Ans | Q | Ans |
|---|---|---|---|---|---|---|---|---|---|
| 1 | a | 2 | b | 3 | c | 4 | a | 5 | c |
| 6 | c | 7 | b | 8 | a | 9 | a | 10 | b |
| 11 | c | 12 | b | 13 | a | 14 | a | 15 | c |
| 16 | c | 17 | a | 18 | a | 19 | b | 20 | c |
| 21 | a | 22 | b | 23 | c | 24 | a | 25 | b |
| 26 | c | 27 | a | 28 | c | 29 | c | 30 | c |
| 31 | a | 32 | a | 33 | c | 34 | a | 35 | c |
| 36 | c | 37 | d | 38 | c | 39 | c | 40 | a |
| 41 | d | 42 | a | 43 | b | 44 | d | 45 | a |

GHOST AVIATOR