


Atoms & Nuclear Radiation.

Unit review

After Lesson 10

Mixed exam practice from the whole unit, a fix-it log, and extension pages for GCSE Physics (separate science) students.



NAME

CLASS

TEACHER

TARGET GRADE

Toolkit.

Learn these: exam questions expect you to recall them without a data sheet.

The three particles

Lesson 1

Particle	Charge	Mass	Where
Proton	+1	1	nucleus
Neutron	0	1	nucleus
Electron	-1	very small	energy levels

Reading the symbol

Lesson 1

A_ZX e.g. ${}^{23}_{11}\text{Na}$

- Z, atomic number = protons
- A, mass number = protons + neutrons
- neutrons = A - Z; in an atom, electrons = Z
- Atom radius $\approx 1 \times 10^{-10}$ m; nucleus radius $< 1/10\,000$ of that

Isotopes and ions

Lesson 2

- **Isotopes**: same number of protons, different numbers of neutrons.
- **Positive ion**: an atom that has lost one or more outer electrons. The nucleus is unchanged.

Alpha scattering

Lesson 3

Observation	Conclusion
most straight through	mostly empty space
some deflected	positive centre
a very few bounced back	mass in a tiny centre

Models and energy levels

Lesson 4

- 1897 electron $\rightarrow$ 1904 plum pudding $\rightarrow$ 1911 nuclear $\rightarrow$ 1913 Bohr $\rightarrow$ 1932 neutron (Chadwick)
- **Absorb** EM radiation $\rightarrow$ higher level, further out.
- **Emit** EM radiation $\rightarrow$ lower level, closer in.

Nuclear radiation

Lesson 5



- Unstable nuclei decay at **random** to become more stable.
- **Activity**: decays per second, in becquerel (Bq). **Count rate**: decays recorded per second by a detector.
- α : $2p + 2n$; β : fast electron, a neutron $\rightarrow$ proton; γ : EM radiation; n: a neutron

Nuclear equations

Lesson 6

Emitted	Symbol	Mass no.	Atomic no.
alpha	${}^4_2\text{He}$	- 4	- 2
beta	${}^0_{-1}\text{e}$	no change	+ 1
gamma	γ	no change	no change
neutron	${}^1_0\text{n}$	- 1	no change

Top numbers balance; bottom numbers balance.



Properties of radiation

Lesson 7

	Alpha	Beta	Gamma
Ionising	strongly	moderately	weakly
Range in air	a few cm	about 1 m	very far
Stopped by	paper, skin	a few mm of aluminium	reduced by thick lead or concrete
Typical use	smoke alarm	thickness gauge	medical tracer

More ionising → less penetrating.

Half-life from a graph

Lesson 8

1. Read the start value.
2. Halve it.
3. Across to the curve, down to the time axis.
4. Check with a second halving.

Half-life calculations

Lesson 9

- Number of half-lives = time ÷ half-life.
- After n half-lives, $(\frac{1}{2})^n$ is left: $\frac{1}{2}$, $\frac{1}{4}$, $\frac{1}{8}$, $\frac{1}{16}$...
- **HT** net decline = initial – final; as a ratio of the initial, $1 - (\frac{1}{2})^n$: $\frac{1}{2}$, $\frac{3}{4}$, $\frac{7}{8}$...

Contamination and irradiation

Lesson 10

- **Irradiation:** exposed to radiation; does not become radioactive.
- **Contamination:** radioactive atoms on or in an object.
- Outside the body γ is most hazardous; inside, α is.
- Tongs, distance, short time, lead, dosimeter.

Unit review.

Mixed questions from the whole unit, like a real exam. Try each question without looking back. Then mark it with your teacher's answers and fill in the "fix it" log.



AQA - STYLE

Mixed exam practice

42 marks, about 50 minutes

- 1 Complete the table. [3 marks]

Particle	Relative charge	Relative mass	Where in the atom?
proton		1	
neutron	0		
electron		very small	

- 2 An atom of cobalt is $^{59}_{27}\text{Co}$. (a) Give the numbers of protons, neutrons and electrons. (b) The atom loses two electrons. How many electrons does the ion have, and what is its charge? [3 marks]

- 3 Explain why $^{35}_{17}\text{Cl}$ and $^{37}_{17}\text{Cl}$ are isotopes of the same element. [2 marks]

- 4 Describe how the results of the alpha particle scattering experiment led to the nuclear model replacing the plum pudding model. [6 marks]

- 5 Describe what happens to an electron in an atom when the atom emits electromagnetic radiation. [2 marks]

- 6 Complete the two nuclear equations. [4 marks]



- 7 A student measures the count rate from a source with different absorbers. [3 marks]
The background count rate is 25 counts per minute. Which types of radiation does the source emit? Explain your answer.

Absorber	none	paper	5 mm aluminium	2 cm lead
Count rate / counts per minute	410	405	150	60

- 8 A doctor needs a radioactive tracer that is injected into a patient and detected outside the body. Which isotope should be used? Give two reasons. [3 marks]

Isotope	Radiation emitted	Half-life
A	alpha	6 hours
B	gamma	6 hours
C	gamma	30 years
D	beta	8 days

- 9 The table shows the count rate from a sample. Determine the half-life. [2 marks]

Time / hours	0	1	2	3	4	5	6
Count rate / counts per second	640	453	320	226	160	113	80

- 10 **HIGHER TIER** A source has an initial activity of 800 Bq. Calculate its activity after 4 half-lives, and the net decline as a ratio of the initial activity. [2 marks]

- 11 Explain why a sample of 50 unstable nuclei rarely has exactly 25 left after one half-life. [2 marks]

12 Compare the hazards of irradiation and contamination by an alpha source. [4 marks]

13 Explain why scientists' model of the atom has changed over time. Use one example. [3 marks]

14 Explain why beta radiation is used to monitor the thickness of paper. [3 marks]



REVIEW LATER

Fix-it log

Turn every lost mark into a target.

Q	What went wrong?	The correct physics (or method) is...	Lesson to revisit

Background radiation and medical uses.

Only for students taking GCSE Physics (separate science). Combined Science students can skip these pages.

● Background radiation

Background radiation is around us all the time. It comes from **natural sources**, such as rocks (for example granite, which gives off radon gas) and cosmic rays from space, and from **man-made sources**, such as fallout from nuclear weapons testing and nuclear accidents. The level depends on where you live and on your job.

The radiation dose a person receives is measured in **sieverts (Sv)**: 1000 millisieverts (mSv) = 1 Sv. When you measure a source, subtract the background count rate to get the corrected count rate.

● Different half-lives, different hazards

An isotope with a **short half-life** has a high activity at first but soon becomes safe. One with a **long half-life** stays radioactive for a very long time, so waste containing it must be stored safely for thousands of years.

● Nuclear radiation in medicine

Use	How it works	Radiation and half-life
Exploring (tracer)	a gamma emitter is injected or swallowed; a detector outside the body shows where it goes	gamma, short half-life (e.g. technetium-99m, 6 hours)
Treating cancer (radiotherapy)	beams of gamma rays from outside the body, or a source placed inside a tumour, kill cancer cells	gamma (or beta from an implant); beams are aimed to spare healthy cells

! Doctors weigh the **risk** (damage to healthy cells) against the **benefit** (finding or treating an illness).



PRACTISE

Check it

Answers are in the answer edition.

1 Name one natural and one man-made source of background radiation. [2 marks]


- 2 A detector records 185 counts per minute from a source. The background count rate is 25 counts per minute. Calculate the corrected count rate. [1 mark]

- 3 Explain why a tracer injected into a patient should have a short half-life and emit gamma radiation. [4 marks]

Nuclear fission and fusion.

Only for students taking GCSE Physics (separate science). These ideas build on nuclear equations (Lesson 6).

● Nuclear fission



One possible fission of uranium-235. Top numbers: $235 + 1 = 141 + 92 + 3$. Bottom numbers: $92 = 56 + 36$.

Fission is the splitting of a large, unstable nucleus, such as uranium-235 or plutonium-239. Spontaneous fission is rare: usually the nucleus must first **absorb a neutron**. It then splits into **two smaller nuclei** of roughly equal size, and gives out **two or three neutrons** and gamma rays. **Energy** is released: all the products move away with kinetic energy.

The neutrons can go on to cause more fissions: a **chain reaction**. In a nuclear reactor it is **controlled** so energy is released steadily; in a nuclear weapon it is uncontrolled, causing an explosion.

● Nuclear fusion

Fusion is the joining of two light nuclei, such as isotopes of hydrogen, to form a heavier nucleus. Some of the mass is converted into energy, which is given out as radiation. Fusion releases the energy of the Sun and other stars.



- 1 Draw a diagram to show how one fission can start a chain reaction. Label the neutrons and the new nuclei. [3 marks]

- 2 Compare nuclear fission with nuclear fusion. [4 marks]

Key words.

Learn the meaning and spelling. In exams, precise words earn marks: say “nucleus”, not “middle”, and “unstable nucleus”, not “unstable atom”.

Absorb: Take in energy. An electron that absorbs EM radiation moves to a higher energy level.

Activity: The rate at which a source of unstable nuclei decays. Unit: becquerel (Bq).

Alpha particle (α): Two protons and two neutrons: a helium nucleus. Strongly ionising; stopped by paper.

Alpha scattering: The experiment in which alpha particles fired at thin gold foil showed that atoms have a tiny, positive nucleus.

Atom: The smallest part of an element. It has a tiny nucleus with electrons around it and no overall charge.

Atomic number: The number of protons in the nucleus. It decides the element.

Becquerel (Bq): Unit of activity: 1 Bq = 1 decay per second.

Beta particle (β): A high-speed electron ejected from the nucleus when a neutron turns into a proton.

Bohr model: Electrons orbit the nucleus at specific distances, called energy levels (1913).

Contamination: Unwanted radioactive atoms on or in an object.

Count rate: The number of decays recorded each second by a detector.

Daughter nucleus: The new nucleus formed when a nucleus decays.

Dosimeter: A badge that measures how much radiation a worker receives.

Electron: A tiny particle with a relative charge of -1 , found in energy levels around the nucleus.

Emit: Give out. An electron that moves to a lower energy level emits EM radiation.

Energy level: A specific distance from the nucleus at which electrons are found.

Evidence: Observations and measurements used to test a model.

Gamma ray (γ): Electromagnetic radiation emitted from a nucleus. Weakly ionising; very penetrating.

Half-life: The time for the number of unstable nuclei in a sample, or the count rate, to halve.

Ion: An atom that has lost (or gained) electrons, so it has an overall charge.

Ionising: Able to knock electrons off atoms, turning them into ions.

Irradiation: Exposing an object to nuclear radiation. The object does not become radioactive.

Isotopes: Atoms of the same element with different numbers of neutrons.

Mass number: The total number of protons and neutrons in the nucleus.

Model: A simplified explanation of something we cannot see directly; changed when new evidence does not fit.

Net decline: (HT) How much the activity has fallen; often given as a ratio of the initial activity.

Neutron: A particle in the nucleus with no charge and a relative mass of 1.

Nuclear equation: An equation showing a decay, in which the mass numbers and atomic numbers balance.

Nuclear model: A tiny, dense, positive nucleus with electrons outside it (Rutherford, 1911).

Nucleus: The tiny centre of an atom, containing protons and neutrons and almost all the mass.

Peer review: Checking of published findings by other scientists.

Penetration: How far radiation can travel through materials before it is absorbed.

Plum pudding model: A ball of positive charge with negative electrons embedded in it (1904).

Proton: A particle in the nucleus with a relative charge of $+1$ and a relative mass of 1.

Radioactive decay: An unstable nucleus giving out radiation to become more stable. It is random.

Random: Impossible to predict for one nucleus: which one decays next, or when.

Unstable nucleus: A nucleus that will decay and give out radiation.

Progress tracker.

Colour one circle for each statement: **red** = not yet, **amber** = getting there, **green** = I can do this without help. Revisit your reds first.

I CAN...	R	A	G
01: Inside the atom			
describe the charge, mass and position of each particle	☐	☐	☐
state the size of an atom and of its nucleus	☐	☐	☐
explain why an atom is neutral	☐	☐	☐
work out protons, neutrons and electrons from nuclear notation	☐	☐	☐
02: Isotopes and ions			
explain what isotopes are	☐	☐	☐
use nuclear notation to compare isotopes	☐	☐	☐
describe how a positive ion forms	☐	☐	☐
work out the charge on an ion	☐	☐	☐
03: From plum pudding to the nucleus			
describe the plum pudding model	☐	☐	☐
describe the alpha scattering experiment	☐	☐	☐
link each result to a conclusion about the atom	☐	☐	☐
compare the plum pudding and nuclear models	☐	☐	☐
04: Energy levels and the nucleus			
describe Bohr's model of the atom	☐	☐	☐
explain what happens when electrons absorb or emit EM radiation	☐	☐	☐
describe the discovery of protons and neutrons	☐	☐	☐
explain why models of the atom have changed	☐	☐	☐
05: Radioactive decay			
explain why unstable nuclei decay	☐	☐	☐
describe decay as random	☐	☐	☐
define activity and count rate	☐	☐	☐
describe alpha, beta, gamma and neutron radiation	☐	☐	☐

06: Nuclear equations

write the symbols for alpha, beta, gamma and neutron radiation



balance a nuclear equation



find the new nucleus after alpha or beta decay



explain the effect of each decay on mass and charge

**07: Properties and uses of radiation**

compare the properties of alpha, beta and gamma



explain the link between ionising and penetration



identify radiation from absorber data



choose and justify a source for a use

**08: Half-life and random decay**

explain why decay is random but predictable



define half-life in two ways



find a half-life from a decay curve



plot a decay curve from data

**09: Half-life calculations**

calculate how much is left after a number of half-lives



find a half-life from data or a graph



find how long a source takes to fall to a value



(HT) express the net decline as a ratio

**10: Contamination and irradiation**

explain the difference between irradiation and contamination



compare the hazards inside and outside the body



describe precautions and how each reduces the risk



explain why findings are published and peer reviewed

