

Atoms & Nuclear Radiation.

Lesson 05


Radioactive decay

A smoke alarm contains a radioactive source. Nothing switches it on, yet it gives out radiation for hundreds of years. Why?

IN THIS LESSON YOU WILL

- explain why some nuclei give out radiation
- describe radioactive decay as random
- define activity and count rate
- describe what alpha, beta, gamma and neutron radiation are, and how each changes the nucleus

ANSWER EDITION FOR TEACHERS AND SELF-MARKING



NAME

CLASS

TEACHER

TARGET GRADE

How this booklet works.

This is the booklet for **Lesson 5** of the Atoms and Nuclear Radiation unit. It builds on the lessons before it; the “Do now” questions and the quick answers at the back help you check what you remember. Keep it: it is your classwork book *and* your revision guide.



DO NOW

Quick questions from earlier lessons. Remembering again is what makes it stick.



LEARN IT

Explanations, diagrams and worked examples. Read this again when you revise.



TRY IT

Practice that builds up: first with support, then on your own, then a challenge.



LAB

Record what you find in a simulation or a model experiment.



EXAM CORNER

AQA-style questions with a tip on how the marks are given.



REVISE IT

Summary, cover-and-answer questions and an “I can” check to colour in.



Revising at home? (1) Read the Learn it pages. (2) Cover them and answer the “Cover the page” questions in Revise it. (3) Check with the quick answers at the back. (4) Go through the matching slides in the teaching deck, and try the simulations.

● The route through the unit (this lesson is highlighted)

01 Inside the atom

03 From plum pudding to the nucleus

05 Radioactive decay

Unstable nuclei, activity and count rate, alpha, beta, gamma and neutron radiation (slides 22–26)

07 Properties and uses of radiation

09 Half-life calculations

02 Isotopes and ions

04 Energy levels and the nucleus

06 Nuclear equations

08 Half-life and random decay

10 Contamination and irradiation

Teaching deck: panphy.app/gcsephy/decks/atoms-and-radiation. Simulations:

panphy.app/simulations/atomic_models.html and panphy.app/simulations/nuclear_decay.html

Radioactive decay.

AQA Physics 4.4.2.1; Combined Science 6.4.2.1

BIG QUESTION

A smoke alarm contains a radioactive source. Nothing switches it on, yet it gives out radiation for hundreds of years. Why?

BY THE END YOU CAN

- explain why some nuclei give out radiation
- describe radioactive decay as random
- define activity and count rate
- describe what alpha, beta, gamma and neutron radiation are, and how each changes the nucleus

unstable nucleus

radioactive decay

random

activity

becquerel (Bq)

count rate

Geiger-Müller tube

alpha

beta

gamma

SLIDES AND SIMULATIONS

Teaching deck, slides 22–26

Radioactive decay, activity and count rate, then alpha, beta and gamma.

panphy.app/gcsephy/decks/atoms-and-radiation/#22

Nuclear Decay simulation

panphy.app/simulations/nuclear_decay.htm



RETRIEVE

Do now

From earlier lessons. No notes!

1

What is an isotope?

An atom of the same
element with a different
number of neutrons.

2

What happens when an
electron emits EM radiation?

It moves to a lower energy
level, closer to the
nucleus.

3


How many protons and
neutrons are in $^{14}_6\text{C}$?

6 protons, 8 neutrons.



When a nucleus is unstable

Unstable nuclei



Some isotopes have an **unstable nucleus**. It gives out **radiation** as it changes to become **more stable**. This is **radioactive decay**.

Decay is **random**: you cannot predict which nucleus will decay next, or when. Nothing we do changes it: heating, cooling or chemical reactions have no effect.

Quantity	Meaning	Unit
Activity	the rate at which a source of unstable nuclei decays: the number of decays each second	becquerel (Bq); 1 Bq = 1 decay per second
Count rate	the number of decays recorded each second by a detector, such as a Geiger-Müller tube	counts per second (or per minute)

THINK OF POPCORN


In a hot pan, you cannot tell which kernel will pop next, or when. But with lots of kernels, you can predict roughly how many will pop each minute.

Where the model breaks: heating makes popcorn pop faster, but nothing we do changes the rate of radioactive decay.


● Four kinds of nuclear radiation

Radiation	What it is	Symbol	Charge	What happens in the nucleus
Alpha (α)	2 protons + 2 neutrons: a helium nucleus	${}^4_2\text{He}$	+2	loses 2 protons and 2 neutrons
Beta (β)	a high-speed electron ejected from the nucleus	${}^0_{-1}\text{e}$	-1	a neutron turns into a proton
Gamma (γ)	electromagnetic radiation from the nucleus	γ	0	loses energy only
Neutron (n)	a neutron	${}^1_0\text{n}$	0	loses one neutron

● Alpha decay




● Beta decay



In beta decay, a **neutron changes into a proton**, and a fast electron is made and shoots out of the nucleus. The beta particle comes from the **nucleus**, not from the energy levels.

Gamma emission



Gamma rays are **electromagnetic waves**. The nucleus gives out extra energy, but its protons and neutrons do not change. Gamma often follows alpha or beta decay.



PRACTISE

Try it

1 Draw a line from each type of radiation to what it is.

1 alpha (α)

→ C

A a neutron

2 beta (β)

→ B

B a high-speed electron from the nucleus

3 gamma (γ)

→ D

C two protons and two neutrons

4 neutron (n)

→ A

D electromagnetic radiation

2 Which statement about radioactive decay is correct? Tick **one** box.

- ☐ Heating a source makes it decay faster.
- ☐ You can predict when a particular nucleus will decay.
- ☒ Unstable nuclei decay at random to become more stable.
- ☐ A nucleus decays when a detector is placed near it.

3 Complete the sentences.

The activity of a source is the number of decays each second. It is measured in becquerels (Bq). The number of decays a Geiger-Müller tube records each second is the count rate.

WORD BANK

activity

count rate

becquerels

half-life

seconds

4

SUPPORTED

Which radiation causes each change in the nucleus?

Change in the nucleus	Radiation
loses 2 protons and 2 neutrons	alpha
a neutron turns into a proton	beta
loses energy, but no particles change	gamma
loses one neutron	neutron

5

Explain what is wrong with Ava's statement.

Ava

Beta particles are electrons knocked out of the energy levels, so beta decay just turns the atom into an ion.

A beta particle comes from the nucleus: a neutron turns into a proton and a fast electron is emitted. The nucleus gains a proton, so the atom becomes a different element; it is not just an ion.

6

CHALLENGE

A Geiger–Müller tube records 1800 counts in 1 minute. (a) Calculate the count rate in counts per second. (b) Suggest why the activity of the source is greater than this count rate.

(a) $1800 \div 60 = 30$ counts per second. (b) Radiation goes out in all directions, so only some of it enters the tube; some is absorbed by the air or the tube wall before it is detected.



Open the **Nuclear Decay** simulation. For each tab, press **Decay this nucleus** and read the equation it shows.

7 Record what happens.

Tab	Parent nucleus	What leaves the nucleus?	Change in mass number	Change in atomic number	New nucleus
01 α	americium-241	alpha particle (He nucleus)	-4	-2	neptunium-237
02 β	carbon-14	beta particle (electron)	0	+1	nitrogen-14
03 γ	technetium-99m	gamma ray	0	0	technetium-99

8 Which decays make a new element? Explain using your table.

Alpha and beta: the atomic number (number of protons) changes. Gamma does not change the atomic number, so the element stays the same.



Get the marks: Two marks: say what the nucleus gives out *and* why.

9 Describe what is meant by radioactive decay. [2 marks]

An unstable nucleus gives out radiation (1) as it changes to become more stable (1).
(Accept: it is random.)

10 Compare an alpha particle with a beta particle. [3 marks]

Alpha is 2 protons and 2 neutrons (a helium nucleus); beta is a high-speed electron (1). Alpha has a charge of +2; beta -1 (1). Alpha is much more massive / beta is made when a neutron turns into a proton (1).

**SUMMARY**

- Unstable nuclei give out radiation to become more stable: radioactive decay.
- Decay is random: we cannot predict which nucleus decays, or when.
- Activity: decays per second, in becquerel (Bq).
Count rate: decays recorded per second by a detector.
- Alpha: $2p + 2n$ (helium nucleus). Beta: fast electron; a neutron becomes a proton. Gamma: EM radiation. Neutron: a neutron.

COVER THE PAGE AND ANSWER

1 Why do some nuclei give out radiation?

2 What is the unit of activity?

3 What is an alpha particle?

4 What happens in the nucleus in beta decay?

I CAN...

explain why unstable nuclei decay



describe decay as random



define activity and count rate



describe alpha, beta, gamma and neutron radiation



Toolkit.

The facts and methods you have met so far. Learn these: exam questions expect you to recall them without a data sheet.

The three particles

Lesson 1

Particle	Charge	Mass	Where
Proton	+1	1	nucleus
Neutron	0	1	nucleus
Electron	-1	very small	energy levels

Reading the symbol

Lesson 1

$${}^A_Z\text{X}$$
 e.g. ${}^{23}_{11}\text{Na}$

- Z, atomic number = protons
- A, mass number = protons + neutrons
- neutrons = A - Z; in an atom, electrons = Z
- Atom radius $\approx 1 \times 10^{-10}$ m; nucleus radius $< 1/10\,000$ of that

Isotopes and ions

Lesson 2

- **Isotopes**: same number of protons, different numbers of neutrons.
- **Positive ion**: an atom that has lost one or more outer electrons. The nucleus is unchanged.

Alpha scattering

Lesson 3

Observation	Conclusion
most straight through	mostly empty space
some deflected	positive centre
a very few bounced back	mass in a tiny centre

Models and energy levels

Lesson 4

- 1897 electron → 1904 plum pudding → 1911 nuclear → 1913 Bohr → 1932 neutron (Chadwick)
- **Absorb** EM radiation → higher level, further out.
- **Emit** EM radiation → lower level, closer in.

Nuclear radiation

Lesson 5

- Unstable nuclei decay at **random** to become more stable.
- **Activity**: decays per second, in becquerel (Bq). **Count rate**: decays recorded per second by a detector.
- α : $2p + 2n$; β : fast electron, a neutron → proton; γ : EM radiation; n: a neutron

Key words from this lesson.

Learn the meaning and spelling. In exams, precise words earn marks: say “nucleus”, not “middle”, and “unstable nucleus”, not “unstable atom”.

Activity: The rate at which a source of unstable nuclei decays. Unit: becquerel (Bq).

Alpha particle (α): Two protons and two neutrons: a helium nucleus. Strongly ionising; stopped by paper.

Becquerel (Bq): Unit of activity: 1 Bq = 1 decay per second.

Beta particle (β): A high-speed electron ejected from the nucleus when a neutron turns into a proton.

Count rate: The number of decays recorded each second by a detector.

Gamma ray (γ): Electromagnetic radiation emitted from a nucleus. Weakly ionising; very penetrating.

Radioactive decay: An unstable nucleus giving out radiation to become more stable. It is random.

Unstable nucleus: A nucleus that will decay and give out radiation.

CHECK YOURSELF

Quick answers.

Answers to the “Do now” and “Cover the page” questions, so you can check yourself when revising. Answers to the longer questions are in the answer edition.

DO NOW

1. An atom of the same element with a different number of neutrons.
2. It moves to a lower energy level, closer to the nucleus.
3. 6 protons, 8 neutrons.

COVER THE PAGE

1. They are unstable; they decay to become more stable.
2. Becquerel (Bq).
3. 2 protons and 2 neutrons (a helium nucleus).
4. A neutron turns into a proton (and a fast electron is emitted).