सत्यमेव जयते

Years of Service to the Nation
राष्ट्र सेवा के 150 वर्ष

राज्य समग्र कृषि-मौसम परामर्श बुलेटिन (तमिलनाडु, पुडुचेरी और कारईकाल के लिए)

State Composite Agromet Advisory Bulletin (for Tamil Nadu, Puducherry, and Karaikal)

Bulletin No: 057/2026, Issued on 17.07.2026

(अवधि: 18 जुलाई से 22 जुलाई 2026, For the period: 18th July to 22th July 2026)





संकलन एवं प्रकाशन / Compiled and issued by:

कृषि-मौसम परामर्श सेवा इकाई (AASU)

भारत मौसम विज्ञान विभाग / India Meteorological Department

प्रादेशिक मौसम विज्ञान केन्द्र चेन्नई/ Regional Meteorological Centre (RMC), Chennai

Website: https://mausam.imd.gov.in/chennai/

Email: rmcmdsaasu@gmail.com; Contact: +91-44-28246027

नोट: यह बुलेटिन तमिलनाडु, पुडुचेरी और कारईकाल के अंतर्गत सभी कृषि-मौसम प्रक्षेत्र इकाई द्वारा तैयार की गई कृषि-मौसम परामर्श बुलेटिन का संकलन है।

Note: This bulletin consolidates Agromet advisories prepared by all AMFUs under Tamil Nadu, Puducherry & Karaikal.

Weather warnings (Met-Subdivision Level)

HEAVY RAINFALL WARNING	
Day 1 (17.07.2026)	Heavy to Very Heavy rain is likely to occur at isolated places over **Keralam, Coastal Karnataka and South Interior Karnataka.**
	Heavy rain is likely to occur at isolated places over **North Interior Karnataka and Telangana.**
Day 2 (18.07.2026)	Heavy rain is likely to occur at isolated places over **Keralam and Karnataka.**
Day 3 (19.07.2026)	Heavy rain is likely to occur at isolated places over **Keralam and Coastal Karnataka.**

THUNDERSTORM WARNING	
Day 1 (18.07.2026) & Day 2 (19.07.2026)	Thunderstorm & lightning with gusty winds speed reaching **30-40 kmph** is likely to occur at one or two places over **Tamil Nadu, Puducherry & Karaikal area and Keralam.** Thunderstorm & lightning is likely to occur at one or two places over **Coastal Andhra Pradesh and Telangana.**
Day 3 (20.07.2026)	Thunderstorm & lightning with gusty winds speed reaching **30-40 kmph** is likely to occur at one or two places over **Tamil Nadu, Puducherry & Karaikal area and Keralam.**
Day 4 (21.07.2026) & Day 5 (22.07.2026)	Thunderstorm & lightning with gusty winds speed reaching **30-40 kmph** is likely to occur at one or two places over **Tamil Nadu, Puducherry and Karaikal area.**

WIND WARNING	
Day 1 (18.07.2026)	Strong surface wind speed reaching **50-60 kmph** is likely to prevail over **Telangana.** Strong surface wind speed reaching **40-50 kmph** is likely to prevail over **Interior Karnataka, Coastal Andhra Pradesh and Rayalaseema.** Strong surface wind speed reaching **30-40 kmph** is likely to prevail over **Coastal Karnataka.**
Day 2 (19.07.2026)	Strong surface wind speed reaching **50-60 kmph** is likely to prevail over **Telangana.** Strong surface wind speed reaching **40-50 kmph** is likely to prevail over **Interior Karnataka and Coastal Andhra Pradesh.** Strong surface wind speed reaching **30-40 kmph** is likely to prevail over **Coastal Karnataka and Rayalaseema.**
Day 3 (20.07.2026)	Strong surface wind speed reaching **40-50 kmph** is likely to prevail over **North Interior Karnataka and Coastal Andhra Pradesh.** Strong surface wind speed reaching **30-40 kmph** is likely to

	prevail over **Coastal Karnataka, South Interior Karnataka, Rayalaseema and Telangana.**
Day 4 (21.07.2026) & Day 5 (22.07.2026)	Strong surface wind speed reaching **40-50 kmph** is likely to prevail over **Coastal Andhra Pradesh.** Strong surface wind speed reaching **30-40 kmph** is likely to prevail over **Karnataka, Rayalaseema and Telangana.**
Day 6 (23.07.2026) & Day 7 (24.07.2026)	Strong surface wind speed reaching **30-40 kmph** is likely to prevail over **Karnataka.**

TEMPERATURE WARNING	
Day 1 (18.07.2026)	High maximum temperature combined with high humidity is likely to cause discomfort (hot & humid) weather over **North Coastal Tamil Nadu, Coastal Andhra Pradesh and Rayalaseema.**

Extended Range forecast at Met-Subdivision Level (23th July to 29th July 2026)

Meteorological Sub-Division	Rainfall	Max Temp	Min Temp
Tamilnadu, PDC & KKL	Below Normal	Above Normal	Normal
Kerala & Mahe	Below Normal	Normal	Normal
Lakshadweep	Below Normal	Normal	Normal
Coastal Karnataka	Normal	Below Normal	Normal
North Interior Karnataka	Below Normal	Below Normal	Normal
South Interior Karnataka	Below Normal	Below Normal	Normal
Coastal Andhra Pradesh & Yanam	Below Normal	Above Normal	Normal
Rayalaseema	Below Normal	Normal	Normal
Telangana	Below Normal	Below Normal	Normal

Summary of past three days rainfall over the State (14th July to 16th July 2026)

Met Subdivision	Day 1 (14.07.2026)	Day 2 (15.07.2026)	Day 3 (16.07.2026)
Tamilnadu, PDC & KKL	DRY	DRY	DRY
Keralam & Mahe	ISOL	ISOL	ISOL
Lakshadweep	DRY	DRY	DRY
Coastal Karnataka	ISOL	ISOL	ISOL
North Interior Karnataka	DRY	DRY	ISOL
South Interior Karnataka	ISOL	DRY	DRY
Coastal Andhra Pradesh & Yanam	DRY	ISOL	ISOL
Rayalaseema	DRY	DRY	ISOL
Telangana	DRY	DRY	ISOL

ISOL: Isolated *i.e.* Rain at 1 or 2 places
SCT: Scattered *i.e.* Rain at a few places
FWS: Fairly Widespread *i.e.* Rain at many places
WS: Widespread *i.e.* Rain at most places
DRY: No rain

Rainfall forecast for next 5 days

Met Subdivision	Day 1	Day 2	Day 3	Day 4	Day 5
	(18.07.2026)	(19.07.2026)	(20.07.2026)	(21.07.2026)	(22.07.2026)
Tamilnadu, PDC & KKL	ISOL	ISOL	ISOL	ISOL	ISOL
Keralam & Mahe	WS	WS	WS	FWS	FWS
Lakshadweep	FWS	FWS	FWS	FWS	FWS
Coastal Karnataka	WS	WS	WS	FWS	FWS
North Interior Karnataka	SCT	SCT	ISOL	ISOL	ISOL
South Interior Karnataka	SCT	SCT	SCT	ISOL	ISOL
CAP & Yanam	SCT	SCT	ISOL	ISOL	ISOL
Rayalaseema	ISOL	ISOL	ISOL	ISOL	ISOL
Telangana	SCT	SCT	ISOL	ISOL	ISOL

ISOL: Isolated *i.e.* Rain at 1 or 2 places **SCT:** Scattered *i.e.* Rain at a few places
FWS: Fairly Widespread *i.e.* Rain at many places

Agromet Advisories issued by different Agromet Field Units (AMFUs) of Tamil Nadu and Puducherry

CHENNAI, KANCHEEPURAM, VILLUPURAM, TIRUVANNAMALAI, TIRUVALLUR, VELLORE, CUDDALORE, CHENGALPATTU, TIRUPATTUR AND RANIPET

SMS Advisory:
(18.07.2026-22.07.2026) Light rain at isolated places with thunderstorm & lightning at one or two places is likely to occur over rest Tamil Nadu.

Livestock Advisory:
1. The farmers are advised to preserve rice straw after harvesting the paddy crop which has to be used as livestock feed.
2. The livestock especially dairy cattle, sheep and goat must not be allowed to stand in slushy and filthy area to prevent foot rot.
3. Dry fodders and feed ingredients used for the preparation of concentrate mixture should be dry and free from aflatoxin.
4. Due to high humidity, Young animals may suffer from pneumonia, get veterinary doctors advice.
5. Rabbitaries with asbestos/metal roofs will experience more heat during summer, hence covering the roof with coconut leaves or gunny bags and splashing water over them will reduce the inner micro climate of rabbit shed.
6. The weather conditions prevailing now are a conclusive environment for mosquitoes & files. Hence the farmers are advised to spray insecticides around the farm premises.
7. Fumigation of livestock shed by using neem and notchi leaves will keep the mosquitoes and other vectors away from the animals.
8. Egg laying poultry (layers) suffering from lice infestation must be looked into with proper attention. Dipping solution must be rubbed over the affected body parts and feathers using cloth or cotton, during Sunny part of the day.

Agricultural advisory:
Rice:
Prevailing weather condition is favorable for bacterial leaf blight infection in rice crop. White colour streaks appear leaf and it turns brown in colour and dried. To control spray copper hydroxide @ 300 g/ac. Due to prevailing weather condition chances of thrips infestation in paddy field. To control spray thiamethoxam 40 g/ac or azadirachtin 400 ml/ac (10000 PPM). Due to prevailing weather condition leaf folder infestation is found in paddy crop. To manage them, spray carbosulfan 400 ml/ac or cartap hydrochloride 600 g/ac during morning or evening hours. The forecasted weather may cause leaf folder infestation in rice crop. The farmers are Advised to monitor the field, if symptoms are observed it's advised to spray Chlorantraniliprole 18.5% 150 ml/ha.
Groundnut:
Treat the groundnut seeds with Trichoderma @ 4g/kg just before sowing. Groundnut growing farmers reduce the root rot infection treat the seeds with carbendazim@ 2.0 g/kg of seed 24 hours before sowing (or) Trichoderma viride @ 4.0 g/kg of seed (or) bacillus subtilis @ 10 g/kg seed.
Sugarcane:

Due to prevailing weather condition changes of sugarcane early shoot borer infection. To control spraychlorantraniliprole 18.5 SC (0.5 ml/l) or fipronil 5 % SC (2.0 ml/l).
Coconut:
To control Rugose Spiraling Whitefly in coconut, setup yellow sticky traps (size 5 x 1.5 feet) @ 8/ac in between the trees. Installation of yellow sticky traps on the palm trunk to trap the adult whiteflies. 2. Encourage buildup of parasitoids (Encarsia sp.) and reintroduce parasitized pupae of emerging zones of whitefly outbreak. 3. In sever case, spray neem oil 10 ml/lit of water or spray azadirachtin 2.0 ml/lit of water with 1.0 ml of sticking agent.
Mango:
The mango crop harvesting is in progress. The fruit may be harvested with 1cm stalk to avoid flow of latex. The fruits may be treated with 2ml /lit Ethrel along with few sodium hydroxide pellets for uniform ripening and color development.
Brinjal:
There are possibilities of mealybug infestation in brinjal crops. For Management farmers are advised to spray Profenophos 2ml/lit of water.

ARIYALUR, KARUR, NAGAPATTINAM, PERAMBALUR, THANJAVUR, TIRUVARUR AND TIRUCHIRAPALLI, MAYILADUTHURAI

Agro Advisory:

Crop	Stage	Advisory
General		Since temperature is increasing, apply mulches to conserve soil moisture.
Rice	Harvest	The prevalling weather condition favors for harvest the matured summer paddy.
Rice	Main field	The prevailing weather condition is conducive for leaf folder symptoms kuruvai paddy such as folded leaves or white scraping on the leaves are observed, spray Chlorantraniliprole or Flubendiamide at 0.3 ml per litre of water.
		At active tillering stage of Kuruvai paddy, top dressing can be done with 35 kg urea and 10 kg MOP/acre by keeping thin film of water in the field.
Cattle	Goat	Farmers are advised to apply Butox solution (2 ml in 1 litre water) externally to remove ecto - parasites in sheep and goats.

SMS advisory: Since temperature is increasing, apply mulches to conserve soil moisture.

COIMBATORE

Weather Warning on Agriculture:
Thunderstorms & lightning, squally wind is expected in some places and hence farmers are requested to stake banana crop which are more than five months old.

Agro Advisory:

Crop	Stage	Advisory
General		Drizzling rainfall are expected for the upcoming week in Western Zone districts with the wind speed of 12–16 kmph. Foothills of Western ghat regions may get more rainfall, compared to inlands. Irrigation may be given based on soil moisture.
Rice	Transplanting	By utilizing the light rainfall short duration rice planting may be continued.
Green gram	Sowing	Both green and black gram sown under irrigated conditions and also under rainfed situations may be given gap filling in order to maintain 33 plants/ m2.
Banana	5 months old	Banana plants above 5 months old may be provided with bamboo staking to avoid lodging against wind speed of above 16 kmph along with rainfall. Prevailing weather is conducive for the sigatoka leaf spot, spray Mancozeb 76% WP @ 1.5 to 2 kg / ha in 750 litre of water during non-rainy period.
Turmeric	Vegetative	Turmeric rhizome may be planted after rhizome treatment to avoid pest and disease with following chemical or biological methods. The chemical treatment includes dipping of rhizome in carbendazim solution (1 g/lit) and phosalone solution (2 ml/lit) for controlling rhizome rot and scales and biological methods include Bacillus subtilis 2.5 kg / ha + 300 kg of FYM
Sugarcane	5 months old	Propping and dethrashing is advised to protect the sugarcane from lodging against the wind speed of above 10 km with moderate rainfall. Do earthing-up and trash mulching to conserve soil moisture.
Tomato	Planting	Treat the tomato seeds with Trichoderma viride 4g with rice kanji coating before transplanting. Harvest mature vegetables during dry periods to maintain quality and shelf life.
Animals		Considering the favourable weather, wash the udder with1% Potassium permanganate solution, before and after milking. Keep animal sheds clean and dry to reduce disease incidence under humid conditions.

SMS advisory: Harvest mature vegetables during dry periods to maintain quality and shelf life.

ERODE

Weather Warning on Agriculture: Thunderstorms & lightning, squally wind is expected in some places and hence farmers are requested to stake banana crop which are more than five months old.

Agro Advisory:

Crop	Stage	Advisory
General		Drizzling rainfall are expected for the upcoming week in Western Zone districts with the wind speed of 12–16 kmph. Foothills of Western ghat regions may get more rainfall, compared to inlands. Irrigation may be given based on soil moisture.
Rice	Transplanting	By utilizing the light rainfall short duration rice planting may be continued.
Green gram	Sowing	Both green and black gram sown under irrigated conditions and also under rainfed situations may be given gap filling in order to maintain 33 plants/ m2.
Banana	5 months old	Banana plants above 5 months old may be provided with bamboo staking to avoid lodging against wind speed of above 16 kmph along with rainfall. Prevailing weather is conducive for the sigatoka leaf spot, spray Mancozeb 76% WP @ 1.5 to 2 kg / ha in 750 litre of water during non-rainy period.
Turmeric	Vegetative	Turmeric rhizome may be planted after rhizome treatment to avoid pest and disease with following chemical or biological methods. The chemical treatment includes dipping of rhizome in carbendazim solution (1 g/lit) and phosalone solution (2 ml/lit) for controlling rhizome rot and scales and biological methods include Bacillus subtilis 2.5 kg / ha + 300 kg of FYM
Sugarcane	5 months old	Propping and dethrashing is advised to protect the sugarcane from lodging against the wind speed of above 10 km with moderate rainfall. Do earthing-up and trash mulching to conserve soil moisture.
Tomato	Planting	Treat the tomato seeds with Trichoderma viride 4g with rice kanji coating before transplanting. Harvest mature vegetables during dry periods to maintain quality and shelf life.
Animals		Considering the favourable weather, wash the udder with1% Potassium permanganate solution, before and after milking. Keep animal sheds clean and dry to reduce disease incidence under humid conditions.

SMS advisory: Harvest mature vegetables during dry periods to maintain quality and shelf life.

TIRUPPUR

Weather Warning on Agriculture

Thunderstorms & lightning, squally wind is expected in some places and hence farmers are requested to stake banana crop which are more than five months old.

Agro Advisory:

Crop	Stage	Advisory
General		Drizzling rainfall are expected for the upcoming week in Western Zone districts with the wind speed of 12–16 kmph. Foothills of Western ghat regions may get more rainfall, compared to inlands. Irrigation may be given based on soil moisture.
Rice	Transplanting	By utilizing the light rainfall short duration rice planting may be continued.
Green gram	Sowing	Both green and black gram sown under irrigated conditions and also under rainfed situations may be given gap filling in order to maintain 33 plants/ m2.
Banana	5 months old	Banana plants above 5 months old may be provided with bamboo staking to avoid lodging against wind speed of above 16 kmph along with rainfall. Prevailing weather is conducive for the sigatoka leaf spot, spray Mancozeb 76% WP @ 1.5 to 2 kg / ha in 750 litre of water during non-rainy period.
Turmeric	Vegetative	Turmeric rhizome may be planted after rhizome treatment to avoid pest and disease with following chemical or biological methods. The chemical treatment includes dipping of rhizome in carbendazim solution (1 g/lit) and phosalone solution (2 ml/lit) for controlling rhizome rot and scales and biological methods include Bacillus subtilis 2.5 kg / ha + 300 kg of FYM
Sugarcane	5 months old	Propping and dethrashing is advised to protect the sugarcane from lodging against the wind speed of above 10 km with moderate rainfall. Do earthing-up and trash mulching to conserve soil moisture.
Tomato	Planting	Treat the tomato seeds with Trichoderma viride 4g with rice kanji coating before transplanting. Harvest mature vegetables during dry periods to maintain quality and shelf life.
Animals		Considering the favourable weather, wash the udder with1% Potassium permanganate solution, before and after milking. Keep animal sheds clean and dry to reduce disease incidence under humid conditions.

SMS advisory: Harvest mature vegetables during dry periods to maintain quality and shelf life.

VIRUDHUNAGAR

Crop	Stage	Agromet Advisory
General: Ploughing is recommended to conserve the available soil moisture present in the soil. Livestock Advisory: As high temperatures are expected over the next five days, ensure adequate drinking water and shade facilities for livestock to protect them from heat stress. Avoid sending livestock for grazing during the afternoon; instead, send them out only during the morning and evening hours. Provide green fodder, mineral mixture, and clean drinking water regularly to improve animal health and milk production. Ensure adequate ventilation in cattle sheds and, if necessary, reduce the heat by sprinkling water on the floor or roof. Human Advisory: Due to the high temperatures, avoid working in direct sunlight between 11:00 AM and 3:00 PM. Carry out field operations during the morning or evening hours. To prevent dehydration, drink plenty of water, buttermilk, tender coconut water, or lemon juice at regular intervals. While working under the sun, wear a hat, cap, or head covering, and use light-colored cotton clothing. If you experience dizziness, excessive fatigue, or any signs of heat-related illness, immediately move to a shaded place, take rest, and seek medical attention if necessary. Temperature-based Agricultural Advisory: Provide need-based irrigation to crops and maintain adequate soil moisture. To prevent soil moisture loss due to high temperatures, apply mulching using paddy straw, coconut husk, or other organic materials. Avoid field operations such as fertilizer application and spraying of pesticides and fungicides during the hottest part of the day. Carry out these activities only during the morning or evening hours. In horticultural crops, provide regular irrigation through drip irrigation to reduce the adverse effects of heat stress. Harvest crops that are ready for harvest during the morning or evening and store the produce safely in cool, shaded, and well-ventilated places.		
Irrigated crops		
Paddy	Maturity	To manage ear head bugs, dust/ spray any one of the following, the first during flowering and second a week later (per ha): Quinalphos 1.5% D 25 kg, Neem seed kernel extract 5% (25 kg kernel/ha),Notchi or Ipomoea or Prosopis leaf extract 10%. Spraying is advised based on local weather conditions. Irrigation to be given based on the requirement of the crop.
Maize	Grain filling stage	To manage fall army worm, apply Flubendiamide 480 SC 4 ml/10litre or Chlorantraniliprole 18.5 SC 4 ml/10lit. Spraying is advised based on local weather conditions. Irrigation to be given based on the requirement of the crop.

SMS: Rain: 18.07.2026-22.07.2026-2,4,2,5&5mm. Mulching to be done to improve soil moisture. Maize- To manage fall army worm, applyFlubendiamide 480 SC 4 ml/10litreor Chlorantraniliprole 18.5 SC 4 ml/10lit.

TIRUNELVELI

Crop	Stage	Agromet Advisory
General: Ploughing is recommended to conserve the available soil moisture present in the soil. Livestock Advisory: As high temperatures are expected over the next five days, ensure adequate drinking water and shade facilities for livestock to protect them from heat stress. Avoid sending livestock for grazing during the afternoon; instead, send them out only during the morning and evening hours. Provide green fodder, mineral mixture, and clean drinking water regularly to improve animal health and milk production. Ensure adequate ventilation in cattle sheds and, if necessary, reduce the heat by sprinkling water on the floor or roof. Human Advisory: Due to the high temperatures, avoid working in direct sunlight between 11:00 AM and 3:00 PM. Carry out field operations during the morning or evening hours. To prevent dehydration, drink plenty of water, buttermilk, tender coconut water, or lemon juice at regular intervals. While working under the sun, wear a hat, cap, or head covering, and use light-colored cotton clothing. If you experience dizziness, excessive fatigue, or any signs of heat-related illness, immediately move to a shaded place, take rest, and seek medical attention if necessary. Temperature-based Agricultural Advisory: Provide need-based irrigation to crops and maintain adequate soil moisture. To prevent soil moisture loss due to high temperatures, apply mulching using paddy straw, coconut husk, or other organic materials. Avoid field operations such as fertilizer application and spraying of pesticides and fungicides during the hottest part of the day. Carry out these activities only during the morning or evening hours. In horticultural crops, provide regular irrigation through drip irrigation to reduce the adverse effects of heat stress. Harvest crops that are ready for harvest during the morning or evening and store the produce safely in cool, shaded, and well-ventilated places.		
Irrigated crops		
Paddy	Flowering	To manage ear head bugs, dust/ spray any one of the following, the first during flowering and second a week later (per ha): Quinalphos 1.5% D 25 kg, Neem seed kernel extract 5% (25 kg kernel/ha), Notchi or Ipomoea or Prosopis leaf extract 10%. Spraying is advised based on local weather conditions. Need based irrigation to be given.
Cotton	Flowering	To manage leaf hoppers, spray any one of the following insecticides, Imidacloprid 200SL at 100ml/ha; Imidacloprid 70% WG @30-35 g/ha; Imidacloprid 17.8% SL @ 100 – 125 ml/ha; Azadirachtin 0.03%WSP 2500- 5000g/ha.Spraying is advised based on local weather conditions. Need based irrigation to be given.
Banana	All stages	Provide support (staking) to protect banana trees which are over five months old.

		Apply press mud at 5 kg per plant to reduce the wilt incidence or apply Pseudomonas fluorescens (Pf1) liquid formulation @ 4 l/ha at 2nd, 4th and 6th months after planting through drip system to manage panama wilt and nematode complex. Need based irrigation to be given.

SMS: Rain: 18.07.2026-22.07.2026-4,1,2,2&2mm.Mulching to be done to improve soil moisture. Paddy -To manage ear head bugs, dust/ spray (per ha): Quinalphos 1.5% D 25 kg or Neem seed kernel extract 5% (25 kg kernel/ha) or Notchi or Ipomoea or Prosopis leaf extract 10%.

THOOTHUKUDI

Crop	Stage	Agromet Advisory
General: Ploughing is recommended to conserve the available soil moisture present in the soil. Livestock Advisory: As high temperatures are expected over the next five days, ensure adequate drinking water and shade facilities for livestock to protect them from heat stress. Avoid sending livestock for grazing during the afternoon; instead, send them out only during the morning and evening hours. Provide green fodder, mineral mixture, and clean drinking water regularly to improve animal health and milk production. Ensure adequate ventilation in cattle sheds and, if necessary, reduce the heat by sprinkling water on the floor or roof. Human Advisory: Due to the high temperatures, avoid working in direct sunlight between 11:00 AM and 3:00 PM. Carry out field operations during the morning or evening hours. To prevent dehydration, drink plenty of water, buttermilk, tender coconut water, or lemon juice at regular intervals. While working under the sun, wear a hat, cap, or head covering, and use light-colored cotton clothing. If you experience dizziness, excessive fatigue, or any signs of heat-related illness, immediately move to a shaded place, take rest, and seek medical attention if necessary. Temperature-based Agricultural Advisory: Provide need-based irrigation to crops and maintain adequate soil moisture. To prevent soil moisture loss due to high temperatures, apply mulching using paddy straw, coconut husk, or other organic materials. Avoid field operations such as fertilizer application and spraying of pesticides and fungicides during the hottest part of the day. Carry out these activities only during the morning or evening hours. In horticultural crops, provide regular irrigation through drip irrigation to reduce the adverse effects of heat stress. Harvest crops that are ready for harvest during the morning or evening and store the produce safely in cool, shaded, and well-ventilated places.		
Irrigated crops		

Paddy	Flowering	To manage ear head bugs, dust/ spray any one of the following, the first during flowering and second a week later (per ha): Quinalphos 1.5% D 25 kg, Neem seed kernel extract 5% (25 kg kernel/ha), Notchi or Ipomoea or Prosopis leaf extract 10%. Spraying is advised based on local weather conditions. Need based irrigation to be given.
Banana	All stages	Provide support (staking) to protect banana trees which are over five months old. Apply press mud at 5 kg per plant to reduce the wilt incidence or apply Pseudomonas fluorescens (Pf1) liquid formulation @ 4 l/ha at 2nd, 4th and 6th months after planting through drip system to manage panama wilt and nematode complex. Need based irrigation to be given.

SMS: Rain:20.07.2026&22.7.2026 – 1&1mm.Mulching to be done to improve soil moisture. Paddy - To manage ear head bugs, dust/ spray (per ha): Quinalphos 1.5% D 25 kg or Neem seed kernel extract 5% (25 kg kernel/ha) or Notchi or Ipomoea or Prosopis leaf extract 10%.

TENKASI

Crop	Stage	Agromet Advisory

General:

Ploughing is recommended to conserve the available soil moisture present in the soil.

Livestock Advisory:

As high temperatures are expected over the next five days, ensure adequate drinking water and shade facilities for livestock to protect them from heat stress.

Avoid sending livestock for grazing during the afternoon; instead, send them out only during the morning and evening hours.

Provide green fodder, mineral mixture, and clean drinking water regularly to improve animal health and milk production.

Ensure adequate ventilation in cattle sheds and, if necessary, reduce the heat by sprinkling water on the floor or roof.

Human Advisory:

Due to the high temperatures, avoid working in direct sunlight between 11:00 AM and 3:00 PM.

Carry out field operations during the morning or evening hours.

To prevent dehydration, drink plenty of water, buttermilk, tender coconut water, or lemon juice at regular intervals.

While working under the sun, wear a hat, cap, or head covering, and use light-colored cotton clothing.

If you experience dizziness, excessive fatigue, or any signs of heat-related illness, immediately move to a shaded place, take rest, and seek medical attention if necessary.

Temperature-based Agricultural Advisory:

Provide need-based irrigation to crops and maintain adequate soil moisture.

To prevent soil moisture loss due to high temperatures, apply mulching using paddy straw, coconut husk, or other organic materials.

Avoid field operations such as fertilizer application and spraying of pesticides and fungicides during the hottest part of the day. Carry out these activities only during the morning or evening hours.

In horticultural crops, provide regular irrigation through drip irrigation to reduce the adverse

effects of heat stress. Harvest crops that are ready for harvest during the morning or evening and store the produce safely in cool, shaded, and well-ventilated places.		
Irrigated crops		
Sorghum	Maturity	Harvest early and safeguard the produce from rain.
Paddy	Flowering	To manage ear head bugs, dust/ spray any one of the following, the first during flowering and second a week later (per ha): Quinalphos 1.5% D 25 kg, Neem seed kernel extract 5% (25 kg kernel/ha),Notchi or Ipomoea or Prosopis leaf extract 10%.Spraying is advised based on local weather conditions. Need based irrigation to be given.
Cotton	Maturity	Harvest early and safeguard the produce from rain.
Citrus	Fruiting	The manage citrus canker disease, spray of Copper Oxy Chloride (COC) @ 0.3% followed by 4 sprayings with Streptocyclin 100 ppm + COC @ 0.3 % at monthly intervals after pruning. Spraying is advised based on local weather conditions. Need based irrigation to be given.
Banana	All stages	Provide support (staking) to protect banana trees which are over five months old. Apply press mud at 5 kg per plant to reduce the wilt incidence or apply Pseudomonas fluorescens (Pf1) liquid formulation @ 4 l/ha at 2nd, 4th and 6th months after planting through drip system to manage panama wilt and nematode complex. Need based irrigation to be given.

SMS: Rain:18.07.2026-22.07.2026-3,3,2,2&2mm; Mulching to be done to improve soil moisture.Paddy -To manage ear head bugs, dust/ spray (per ha): Quinalphos 1.5% D 25 kg or Neem seed kernel extract 5% (25 kg kernel/ha) or Notchi or Ipomoea or Prosopis leaf extract 10%.

SALEM

WEATHER ADVISORY BULLETIN FOR LIVESTOCK, POULTRY AND AGRICULTURE

Livestock, Poultry & Crop	Stage	Weather advisory
Livestock	Sheep	Bluetongue Disease and its prevention measures in Sheep Bluetongue disease is an important viral disease that severely affects sheep. It is a virus-borne disease transmitted by the bites of infected biting midges. Although the disease can affect other livestock, it causes the most severe illness in sheep, resulting in significant economic losses for sheep farmers. The major clinical signs include high fever, painful mouth ulcers, swelling of the face and lameness. The tongue may become swollen, turn bluish in color (cyanosis) and may protrude from the mouth. To prevent this disease, a Bluetongue vaccine has been developed in India.

		So, the farmers in Tamil Nadu are advised to administer both the primary and booster doses of the Bluetongue vaccine to their sheep, ensuring that the booster dose is completed by the 2nd week of August. Sheep older than 3 months of age should receive the primary vaccination in July, followed by a booster vaccination 28 days later, preferably by the 2nd week of August. Thereafter, the Bluetongue vaccine should be administered once every year without fail to provide continued protection
Poultry Disease Diagnosis and Surveillance Laboratory, Namakkal		
Poultry	General	Post mortem observations in the birds brought for diagnosis at PDDSL, VCRI, Namakkal revealed death of birds due to Chicken Infectious Anaemia(CIA) and Heat Stroke infection. Hence, the farmers are advised to use appropriate antibiotics, follow feed management techniques and also get consultation from nearby Veterinary Doctor to take appropriate disinfection measures and to implement the biosecurity measures.
Agro advisory from Krishi Vigyan Kendra, Namakkal		
Small Onion	Vegetati-ve Stage	Farmers cultivating small onion are advised to apply a vegetable MN mixture or Zinc sulphate as a foliar spray when the crop is 30 days old. Mix 3g/litre of water of zinc sulphate or vegetable MN mixture + 1 ml of sticking agent and spray during the morning or evening hours. Repeat the spray three times at 15-days interval. This foliar application helps prevent tip blight caused by Zn deficiency. It also improves the yield and quality of the onion.
Horticultu-ral crops	General	Installing drip irrigation for sugarcane and horticultural crops helps to maintain uniform soil moisture and ensures that nutrients are delivered directly to the root zone without being wasted. Since Namakkal has a large no. of poultry farms, poultry manure and vermicompost are readily available. Incorporating these organic manures into the soil during land preparation improves the soil's structure, fertility and overall health.

KRISHNAGIRI

WEATHER ADVISORY BULLETIN FOR LIVESTOCK, POULTRY AND AGRICULTURE

Livestock, Poultry & Crop	Stage	Weather advisory
Livestock	Sheep	Bluetongue Disease and its prevention measures in Sheep Bluetongue disease is an important viral disease that severely affects sheep. It is a virus-borne disease transmitted by the bites of infected biting midges. Although the disease can affect other livestock, it causes the most severe illness in sheep, resulting in significant economic losses for sheep farmers. The major clinical signs include high fever, painful mouth ulcers, swelling of the face and lameness. The tongue may become swollen, turn bluish in color (cyanosis) and may protrude from the mouth. To prevent this disease, a Bluetongue vaccine has been developed in India. So, the farmers in Tamil Nadu are advised to administer both the primary and booster doses of the Bluetongue vaccine to their sheep, ensuring that the booster dose is completed by the 2nd week of August. Sheep older than 3 months of age should receive the primary vaccination in July, followed by a booster vaccination 28 days later, preferably by the 2nd week of August. Thereafter, the Bluetongue vaccine should be administered once every year without fail to provide continued protection
Agro advisory from Krishi Vigyan Kendra, Namakkal		
Small Onion	Vegetati-ve Stage	Farmers cultivating small onion are advised to apply a vegetable MN mixture or Zinc sulphate as a foliar spray when the crop is 30 days old. Mix 3g/litre of water of zinc sulphate or vegetable MN mixture + 1 ml of sticking agent and spray during the morning or evening hours. Repeat the spray three times at 15-days interval. This foliar application helps prevent tip blight caused by Zn deficiency. It also improves the yield and quality of the onion.
Horticultu-ral crops	General	Installing drip irrigation for sugarcane and horticultural crops helps to maintain uniform soil moisture and ensures that nutrients are delivered directly to the root zone without being wasted. Since Namakkal has a large no. of poultry farms, poultry manure and vermicompost are readily available. Incorporating these organic manures into the soil during land preparation improves the soil's structure, fertility and overall health.
Castor	Flowering Stage	The Castor crop was initiated flowering in the current season. Farmers are advised to spray Growth Hormone (GH) namely “Castor gold” at the rate of 0.5 ml/litre of water along with Sticking agent 1 ml/litre of water on 25th and 60th Days after Sowing (DAS) during morning or evening hours to increase the female flowers in spikes and improve the yield in the Castor crop.

DHARMAPURI

WEATHER ADVISORY BULLETIN FOR LIVESTOCK, POULTRY AND AGRICULTURE

Livestock, Poultry & Crop	Stage	Weather advisory
Livestock	Sheep	Bluetongue Disease and its prevention measures in Sheep Bluetongue disease is an important viral disease that severely affects sheep. It is a virus-borne disease transmitted by the bites of infected biting midges. Although the disease can affect other livestock, it causes the most severe illness in sheep, resulting in significant economic losses for sheep farmers. The major clinical signs include high fever, painful mouth ulcers, swelling of the face and lameness. The tongue may become swollen, turn bluish in color (cyanosis) and may protrude from the mouth. To prevent this disease, a Bluetongue vaccine has been developed in India. So, the farmers in Tamil Nadu are advised to administer both the primary and booster doses of the Bluetongue vaccine to their sheep, ensuring that the booster dose is completed by the 2nd week of August. Sheep older than 3 months of age should receive the primary vaccination in July, followed by a booster vaccination 28 days later, preferably by the 2nd week of August. Thereafter, the Bluetongue vaccine should be administered once every year without fail to provide continued protection
Agro advisory from Krishi Vigyan Kendra, Namakkal		
Small Onion	Vegetati-ve Stage	Farmers cultivating small onion are advised to apply a vegetable MN mixture or Zinc sulphate as a foliar spray when the crop is 30 days old. Mix 3g/litre of water of zinc sulphate or vegetable MN mixture + 1 ml of sticking agent and spray during the morning or evening hours. Repeat the spray three times at 15-days interval. This foliar application helps prevent tip blight caused by Zn deficiency. It also improves the yield and quality of the onion.
Horticultu-ral crops	General	Installing drip irrigation for sugarcane and horticultural crops helps to maintain uniform soil moisture and ensures that nutrients are delivered directly to the root zone without being wasted. Since Namakkal has a large no. of poultry farms, poultry manure and vermicompost are readily available. Incorporating these organic manures into the soil during land preparation improves the soil's structure, fertility and overall health.
Castor	Flowering Stage	The Castor crop was initiated flowering in the current season. Farmers are advised to spray Growth Hormone (GH) namely “Castor gold” at the rate of 0.5 ml/litre of water along with Sticking agent 1 ml/litre of water on 25th and 60th Days after Sowing (DAS) during morning or evening hours to increase the

		female flowers in spikes and improve the yield in the Castor crop.

NAMAKKAL

WEATHER ADVISORY BULLETIN FOR LIVESTOCK, POULTRY AND AGRICULTURE

Livestock, Poultry & Crop	Stage	Weather advisory
Livestock	Sheep	Bluetongue Disease and its prevention measures in Sheep Bluetongue disease is an important viral disease that severely affects sheep. It is a virus-borne disease transmitted by the bites of infected biting midges. Although the disease can affect other livestock, it causes the most severe illness in sheep, resulting in significant economic losses for sheep farmers. The major clinical signs include high fever, painful mouth ulcers, swelling of the face and lameness. The tongue may become swollen, turn bluish in color (cyanosis) and may protrude from the mouth. To prevent this disease, a Bluetongue vaccine has been developed in India. So, the farmers in Tamil Nadu are advised to administer both the primary and booster doses of the Bluetongue vaccine to their sheep, ensuring that the booster dose is completed by the 2nd week of August. Sheep older than 3 months of age should receive the primary vaccination in July, followed by a booster vaccination 28 days later, preferably by the 2nd week of August. Thereafter, the Bluetongue vaccine should be administered once every year without fail to provide continued protection
Poultry Disease Diagnosis and Surveillance Laboratory (PDDSL), Namakkal		
Poultry	General	Post mortem observations in the birds brought for diagnosis at PDDSL, VCRI, Namakkal revealed death of birds due to Chicken Infectious Anaemia(CIA) and Heat Stroke infection. Hence, the farmers are advised to use appropriate antibiotics, follow feed management techniques and also get consultation from nearby Veterinary Doctor to take appropriate disinfection measures and to implement the biosecurity measures.
Agro advisory from Krishi Vigyan Kendra, Namakkal		
Small Onion	Vegetati-ve Stage	Farmers cultivating small onion are advised to apply a vegetable MN mixture or Zinc sulphate as a foliar spray when the crop is 30 days old. Mix 3g/litre of water of zinc sulphate or vegetable MN mixture + 1 ml of sticking agent and spray during the morning or evening hours. Repeat the spray three times at 15-days interval. This foliar application helps prevent tip blight caused by Zn

		deficiency. It also improves the yield and quality of the onion.
Horticultu-ral crops	General	Installing drip irrigation for sugarcane and horticultural crops helps to maintain uniform soil moisture and ensures that nutrients are delivered directly to the root zone without being wasted. Since Namakkal has a large no. of poultry farms, poultry manure and vermicompost are readily available. Incorporating these organic manures into the soil during land preparation improves the soil's structure, fertility and overall health.

KANYAKUMARI

AGRO ADVISORIES

S. N.	CROP	STAGES	ADVISORIES
1.	Fruit trees and Tree spices	All stages	Strengthening the basins may be taken up for better harvesting of rains.
2.	Clove and Cinnamon	All stages	Avoid moisture loss by the application of mulching in clove and cinnamon field through farm waste materials.
3.	Coconut	All stages	Green manure crops sown in the coconut basin or in the interspace of coconut gardens have to be incorporated into the soil if they have attained 50% flowering.
4.	Horticultural crops.	Planting stage.	As 12 - 16 km/hr wind speed is anticipated, it is advised to provide staking to newly planted horticultural plants.
5.	Rubber	Tapping Stage	Fix a polythene rain guard to the trunk of the tree above the tapping panel so that tapping can be done during rainy season also.
6.	Agricultural and horticultural crops	Planting Stage	Preparatory cultivation can be started for planting of paddy and horticultural crops such as vegetables, rubber, coconut etc

SMS Advisory: Trim the roots and decayed portion of the corm and paring and pralinage with 40g carbofuran 3G granules per sucker to control the nematode infestation.

KARAIKAL

Agro Advisory:

Crop/ livestock	Stage/ Particulars	Agro Advisory
Cotton	Flowering to picking stage	To manage mealy bug in cotton, spray Thiamethoxam 25 WG @ 0.2 g/l or Buprofezin 25 SC @ 2 ml/l.

		To manage white fly in cotton, spray Azadirachtin 0.03 (300 ppm) @ 5-10 ml/l or Thiamethoxam 25 WG 0.4 g/l.
Banana	Bearing stage	To protect banana plants from toppling or snapping in strong winds, provide firm support (propping or staking) using double bamboo poles. This practice is especially critical for plants bearing heavy bunches and those nearing harvest.
Brinjal	Phomopsis blight	To manage Phomopsis blight in brinjal, spray Pseudomonas fluorescens or Bacillus subtilis @ 5g or 5ml/ l.
	Shoot borer	To manage Brinjal shoot borer, spray Emamectin Benzoate 4% SG @ 0.4g/l.
Coconut	Bearing stage	Root feeding of coconut tonic (200 ml / tree) in 3 months interval to avoid button shedding in coconut.

SMS: Light drizzle to light rain is expected for next five days.

PUDUCHERRY

Agro Advisory

Crop/livestock	Stage/Particulars	Agro Advisory
Rice	Tillering stage	To manage rice stem borer, release parasitoid Trichogramma japonicum @1,00,000/ha at active tillering stage (37, 44 and 51 DAP) or Azadirachtin 0.03% @ 4 ml/l or cartaphydrochloride 50% SP @ 2g/l or flubendiamide 20 WG @ 0.25 g/l or thiamethoxam 25 WG @ 0.2 g/l. To manage rice leaf folder release Trichogramma chilonis 5cc (1,00,000/ha) at 37, 44, 51 day after planting. Avoid excess application of nitrogen fertilizers. Setup light trap to attract adult moths. Setup T- shaped bird perches for predatory birds to control these pests. Apply cartaphydrochloride 50% SP @ 2g/l or flubendiamide 20 WG @ 0.25 g/l or thiamethoxam 25 WG @ 0.2 g/l.
Sugarcane	Vegetative stage	To protect sugarcane crops older than 5 months (over 150 days) from strong winds, carry out 'trash twisting' or 'double-line propping' to prevent lodging. To control shoot borer in sugarcane, soil application of Cartaphydrochloride 4G 10kg/hectare may be done.
Banana	Bearing stage	To protect banana plants from toppling or snapping in strong winds, provide firm support (propping or staking) using double bamboo poles. This practice is especially critical for plants bearing heavy bunches and those nearing harvest.
Cassava	Vegetative stage	To manage red mite in Cassava, apply Pongamia oil 1 ml/l or spray Propargite 1.5 ml and Sulphur 2.5 g/ l.
Coconut	Bearing stage	Root feeding of coconut tonic (200 ml / tree) in 3 months interval to avoid button shedding in coconut.

Livestock	Heat stress	Heat stress is likely to occur in cattle due to prevailing hot weather. Therefore, following steps can be followed to protect livestock from summer heat. Cattle can be protected from heat by spreading straw, dry grass, palm fronds, coconut leaves or coir waste on the surface of cattle sheds and sprinkling water over it. Cattle can be sent for grazing between 06 am to 10 am and 04 pm to 06 pm. Roofs should be erected over the drinking troughs for livestock to provide cold drinking water to the livestock. Cattle should be bathed twice daily. If cattle show signs of heat stress such as excessive drinking of water, rapid breathing, excessive salivation, loss of appetite, shivering and falling down, then the affected animals should be treated at the nearest veterinary hospital.

SMS: Light drizzle to light rain is expected for next four days.


Annexure I

District level Heavy Rainfall Warning for next 5 days

NIL.

Annexure II

Meteorological Subdivision Level Rainfall Weather Forecast for next 5 days of RMC Chennai



Annexure III

District Forecast & Warning for Next 5 Days issued on: 17.07.2026

NAME OF THE DISTRICT	RAINFALL INTENSITY	18.07.2026	19.07.2026	20.07.2026	21.07.2026	22.07.2026
		SAT	SUN	MON	TUE	WED
TIRUVALLUR	Intensity	Moderate	Moderate	Light	Light	Light
	Probability	Likely	Likely	Likely	Likely	Likely
CHENNAI	Intensity	Moderate	Moderate	Light	Light	Light
	Probability	Likely	Likely	Likely	Likely	Likely
CHENGALPATTU	Intensity	Moderate	Moderate	Light	Light	Light
	Probability	Likely	Likely	Likely	Likely	Likely
KANCHEEPURAM	Intensity	Moderate	Moderate	Light	Light	Light
	Probability	Likely	Likely	Likely	Likely	Likely
VILLUPURAM	Intensity	Moderate	Moderate	Light	Light	Light
	Probability	Likely	Likely	Likely	Likely	Likely
PUDUCHERRY	Intensity	Moderate	Moderate	Light	Light	Light
	Probability	Likely	Likely	Likely	Likely	Likely
CUDDALORE	Intensity	Moderate	Moderate	Light	Light	Light
	Probability	Likely	Likely	Likely	Likely	Likely
MAYILADUTHURAI	Intensity	Moderate	Moderate	Light	Light	Light
	Probability	Likely	Likely	Likely	Likely	Likely
NAGAPATTINAM	Intensity	Moderate	Moderate	Light	Light	Light
	Probability	Likely	Likely	Likely	Likely	Likely
KARAIKAL	Intensity	Light	Light	Light	Light	Light
	Probability	Likely	Likely	Likely	Likely	Likely
TIRUVARUR	Intensity	Light	Light	Light	Light	Light
	Probability	Likely	Likely	Likely	Likely	Likely
THANJAVUR	Intensity	Light	Light	Light	Light	Light
	Probability	Likely	Likely	Likely	Likely	Likely
PUDUKKOTTAI	Intensity	Light	Light	Light	Light	Light
	Probability	Likely	Likely	Likely	Likely	Likely
KALLAKURUCHI	Intensity	Light	Light	Light	Light	Light
	Probability	Likely	Likely	Likely	Likely	Likely
VELLORE	Intensity	Light	Light	Light	Light	Light
	Probability	Likely	Likely	Likely	Likely	Likely
RANIPET	Intensity	Light	Light	Light	Light	Light

	Probability	Likely	Likely	Likely	Likely	Likely
TIRUPPATTUR	Intensity	Light	Light	Light	Light	Light
	Probability	Likely	Likely	Likely	Likely	Likely
TIRUVANNAMALAI	Intensity	Light	Light	Light	Light	Light
	Probability	Likely	Likely	Likely	Likely	Likely
ARIYALUR	Intensity	Light	Light	Light	Light	Light
	Probability	Likely	Likely	Likely	Likely	Likely
PERAMBALUR	Intensity	Light	Light	Light	Light	Light
	Probability	Likely	Likely	Likely	Likely	Likely
TIRUCHIRAPALLI	Intensity	Light	Light	Light	Light	Light
	Probability	Likely	Likely	Likely	Likely	Likely
KARUR	Intensity	Light	Light	Light	Light	Light
	Probability	Likely	Likely	Likely	Likely	Likely
NAMAKKAL	Intensity	Light	Light	Light	Light	Light
	Probability	Likely	Likely	Likely	Likely	Likely
SALEM	Intensity	Light	Light	Light	Light	Light
	Probability	Likely	Likely	Likely	Likely	Likely
DHARMAPURI	Intensity	Light	Light	Light	Light	Light
	Probability	Likely	Likely	Likely	Likely	Likely
KRISHNAGIRI	Intensity	Light	Light	Light	Light	Light
	Probability	Likely	Likely	Likely	Likely	Likely
ERODE	Intensity	Light	Light	Light	Light	Light
	Probability	Likely	Likely	Likely	Likely	Likely
NILGIRIS	Intensity	Moderate	Moderate	Moderate	Light-Moderate	Light-Moderate
	Probability	Likely	Likely	Likely	Likely	Likely
COIMBATORE	Intensity	Moderate	Moderate	Moderate	Light-Moderate	Light-Moderate
	Probability	Likely	Likely	Likely	Likely	Likely
TIRUPPUR	Intensity	Moderate	Moderate	Moderate	Light-Moderate	Light-Moderate
	Probability	Likely	Likely	Likely	Likely	Likely
THENI	Intensity	Moderate	Moderate	Moderate	Light-Moderate	Light-Moderate
	Probability	Likely	Likely	Likely	Likely	Likely
DINDIGUL	Intensity	Moderate	Moderate	Moderate	Light-Moderate	Light-Moderate
	Probability	Likely	Likely	Likely	Likely	Likely
MADURAI	Intensity	Moderate	Moderate	Moderate	Light	Light
	Probability	Likely	Likely	Likely	Likely	Likely
VIRUDHUNAGAR	Intensity	Moderate	Moderate	Moderate	Light-Moderate	Light-Moderate
	Probability	Likely	Likely	Likely	Likely	Likely
SIVAGANGA	Intensity	Dry	Dry	Dry	Light	Light
	Probability	Likely	Likely	Likely	Likely	Likely

TENKASI	Intensity	Moderate	Moderate	Moderate	Light-Moderate	Light-Moderate
	Probability	Likely	Likely	Likely	Likely	Likely
RAMANATHAPURAM	Intensity	Dry	Dry	Dry	Light	Light
	Probability	Likely	Likely	Likely	Likely	Likely
THOOTHUKUDI	Intensity	Dry	Dry	Dry	Light	Light
	Probability	Likely	Likely	Likely	Likely	Likely
TIRUNELVELI	Intensity	Moderate	Moderate	Moderate	Light-Moderate	Light-Moderate
	Probability	Likely	Likely	Likely	Likely	Likely
KANYAKUMARI	Intensity	Moderate	Moderate	Moderate	Light-Moderate	Light-Moderate
	Probability	Likely	Likely	Likely	Likely	Likely

Warning Colour Codes	हिन्दी	मराठी
WARNING (TAKE ACTION)	चेतावनी	चेतावणी
ALERT (BE PREPARED)	सतर्क रहें	सतर्क असावे
WATCH (BE UPDATED)	निगरानी रखें	लक्ष असावे
NO WARNING (NO ACTION)	कोई चेतावनी नहीं	धोक्याची सूचना नाही

