Gramin Krishi Mausam Sewa (GKMS)
ALL INDIA COORDINATED RESEARCH PROJECT ON AGROMETEOROLOGY
Vasantrao Naik Marathwada Krishi Vidyapeeth Parbhani 431402
Email : amfuparbhani@gmail.com

Beed District Level AAB No. : 33/ 2026-2027 Day & Date : Friday, 24.07.2026

Observed weather during last week (Dated 18th to 24th July, 2026)							Weather Parameters		Weather Forecast (Valid 25th to 29th July, 2026)				
18/07	19/07	20/07	21/07	22/07	23/07	24/07	Date		25/07	26/07	27/07	28/07	29/07
							Rainfall (mm)		4.0	3.0	3.0	2.0	5.0
							T_{max} (^{0}C)		33.0	33.0	34.0	34.0	33.0
							T_{min} (^{0}C)		22.0	23.0	24.0	25.0	25.0
							AM	Cloud Cover	Fully Cloudy	Fully Cloudy	Cloudy	Fully Cloudy	Fully Cloudy
							PM						
							RH-I (%)		83	82	82	82	82
							RH-II (%)		61	56	56	56	58
							Wind Speed (km/hr)		25	23	18	20	20
							AM	Wind Direction	SW	WSW	WSW	WSW	WSW
							PM						

Weather Summary / Alert :

As per the forecast from RMC Mumbai, Thundershower along with light rainfall may occur at isolated places in Beed District on dated 26th July. Light to moderate rainfall may occur at isolated places in Beed District on dated 24th July. Light rainfall may occur at scattered places on dated 24th & 25th July & at isolated places during 26th to 28th July in Beed District. Gradual rise in maximum temperature by 2 to 3 ^{O}C during next four to five days in Beed District.

General Advisory :

As per ERFS products Rainfall may be below normal, maximum temperature & minimum temperature may be Normal during 24th to 30th July 2026 & Rainfall may be below normal, maximum temperature may be Normal to slightly above Normal & minimum temperature mey be slightly above Normal during 31st July to 06th August 2026.

In Marathwada region, sowing may be initiated in selected circles: Chh. Sambhaji Nagar district: Chh. Sambhaji Nagar Block (Chh. Sambhaji Nagar), Paithan Block (Lohagaon) ; Beed district: Ambejogai Block (Lokhandi), Kej Block (Usuf) : Hingoli District: Kalamnuri Block (Akhada), Wasmat Block (Wasmat, Hayatnagar, Girgaon, Hatta), Sengaon Block (Sengaon, Ajegaon, Sakhara, Hatta) Farmers in other revenue circles are advised not to rush for sowing and wait for favorable rainfall conditions.

SMS Advisory: The areas where sowing is not completed sowing of Sunflower, Pigeonpea, Hybrid Pearl millet, Soybean Pigeonpea (4:2), Pearl millet + Pigeonpea (3:3), Caster, Niger and Sesamum should be done after receipt of (75 to 100 mm) rainfall.

Name of Crop	Stage	Agromet Advisory
Soybean	Growth	Intercultural operation should be done for weed management in soybean crop. If infestation of leaf eating catter piller & stem borer may observed in soybean crop, For management take a spray of any one chemical as given below Profenofos 50% @ 400 ml or Ethion 50% @ 600 ml or Thimethoxam 12.6% + Lambda Cyhalothrin 9.5% @ 50 ml or if incidence is more take a spray of Chlorantraniliprole 18.5% @ 60 ml per acre during clear weather condition. If yellow leaf observed in soybean crop due to micronutrient deficiency for management take a spray of EDTA Chilated mix Micronutrients grade-2 @ 50 g or 50 ml with 19:19:19 @ 100 g per 10 liter of water during clear weather condition. If infestatin of yellow mosaic observed in soybean crop for management, Remove and destroy yellow mosaic infected plants.
Kharif sorghum	Growth	Intercultural operation should be done for weed management in kharif sorghum crop. For management of fall armyworm in kharif sorghumcrop take a spray of Emamectine Benzoate 5 % @ 4 g or Spinetoram 11.7 SC @ 4 ml per 10 liter of water during clear weather condition.
Pearl millet	Growth	Light hoeing should be done in pearl millet crop for weed management and for soil moisture conservation. Light hoeing should be done in soybean crop for weed management and for soil moisture conservation.
Sugarcane	Vegetative Growth	Due to cloudy weather, an infestation of the 'Payrila' (moth) pest is being observed in the sugarcane crop. To monitor the pest and to trap and destroy large numbers of adult moths, a light trap should be installed at a height of six inches above the crop canopy—at a ratio of one trap per five acres—and operated between 6:00 PM and 10:00 PM. For management, take a spray of biological fungus Metarhizium anisopliae @ 40 grams in 10 liters of water or take a spray a 5% NSKE of for chemical control, take a spray Chlorpyrifos 20% @ 600 ml or Monocrotophos 36% @ 200 ml of per acre during clear weather condition.
Turmeric	Growth	For management of Rhizome fly in turmeric take a spray of Quinalphos 25 % @ 20 ml or Diamethoate 30 % @ 10 ml per 10 liter of water with an interval of 15 days during clear weather condition. (As label claim was not given by Central Insecticides Board for Rhizome fly in turmeric the research recommendation of university is mentioned here). Intercultural operations should be done for weed management in turmeric crop during wapsa condition.
Citrus	Growth	As the *Mrig Bahar* orange and sweet lime orchards are currently in the flowering & Fruit Initiation stage, do not spray any insecticide in Mrug bahar citrus orchard. Intercultural operations should be done for weed management in citrus orchard.
Pomegranate	Growth	As the *Mrig Bahar* pomegranate orchards Given the possibility of a disease outbreak in the orchard, spray a 1% Bordeaux mixture during clear weather condition. Intercultural operations should be done for weed management in pomegranate orchard.
Sapota	Growth	Intercultural operations should be done for weed management in Sapota orchard.
Vegetable	Transplanting/ Harvesting	If sufficient moisture is available in soil sowing of direct sown vegetable crops (i.e. by seeds) for example Okra, Bitter gourd, Ridge gourd, Pumpkin can be done. Transplanting of 45 days old seedlings of vegetable crops (Brinjal, Tomato, Chilli etc.) should be done at the places where rainfall was received by considering soil moisture. Harvesting of mature vegetable crops should be done. Weed management should be done in vegetable crops.
Floriculture	Transplanting/ Harvesting	Transplanting of Floriculture crops should be done at the places where rainfall was received by considering soil moisture. Harvesting of mature flowers should be done. Weed management should be done in floricultural crops.
Animal husbandry	--	During the monsoon season, the prevalence of stable flies, mosquitoes, and ticks around livestock increases; therefore, as per advice of veternary doctor spray the insecticides. Livestock should be vaccinated against monsoon-related diseases as recommended by a veternary doctor. In particular, goats and sheep should be vaccinated against enterotoxaemia at the appropriate time to prevent the disease during the monsoon. During monsoon season, avoid feeding livestock excessive amounts of tender, green fodder.

This Agro Advisory Bulletin (AAB) is prepared and published with the consolation and recommendation of SMS committee of "Gramin Krishi Mausam Sewa (GKMS)", Vasantrao Naik Marathwada Krishi Vidyapeeth, Parbhani 431 402 (MS).

Principal Nodal Officer
Gramin Krishi Mausam Sewa (GKMS)
VNMKV, Parbhani