Gramin Krishi Mausam Sewa, AMFU, Bulletin District Yavatmal.

(Issued jointly by GKMS, Akola, Dr Panjabrao Deshmukh Krishi Vidyapeeth, Akola &
(India Meteorology Department) Email: gkmsakola@yahoo.com.

Yavatmal District Agromet Advisory Bulletin No. : 16 / 2026-27 **Day & Date : Friday 24.07.2026**

Weather Parameters	Weather Forecast(Valid for 24th July 2026 To 28th July 2026)				
Date	24/07	25/07	26/07	27/07	28/07
Rainfall (mm)	5.1	2.6	2.5	8.6	4.4
Tmax (0C)	29.7	31.8	31.7	32.1	32.4
Tmin (0C)	24.7	24.7	25.4	25.9	25.9
RH-I (%)	87	86	87	86	84
RH-II (%)	68	57	60	59	57
Wind Speed (km/hr)	16.3	12.4	14.3	15.2	19.1
Wind Direction	225	240	260	266	273
Cloud cover	Cloudy	Cloudy	Cloudy	Cloudy	Cloudy

Crop	Stage	Weather Based Agro Advisory
Forecast	-	• Today's forecast for the district indicates light to moderate rainfall at many places on 24 and 27- 28 July and at few places on 25-26 July There will be rise in maximum temperature by 2-4oC in 48 hours, thereafter no large change in subsequent days over vidharbha region. Download DAMINI App for updated warning alert of lightning. • Warning includes thunderstorm with lightning at one or two places during 27 July.
General	Growth	• In timely sown all kharif crops undertake inter culture operation (weeding / hoeing) during clear weather, for management of weeds and improved soil aeration. • For weed control in timely sown soybean and cotton, alternative option of recommended early post emergence herbicidal spray (with label claim) at proper stage and proper soil moisture condition should be undertaken. • Need based Gap filling and thinning operation may be carried out under sufficient soil moisture condition.
Soybean	Growth	• In timely sown soybean crop undertake inter culture operation (weeding / hoeing) during clear weather for management of weeds and improved soil aeration. • For weed control in timely sown soybean, alternative option of recommended early post emergence herbicidal spray (with label claim) at proper stage and proper soil moisture condition should be undertaken.
Cotton	Growth	• Need based gap filling may be undertaken to maintain optimum plant population in irrigated fields. • In timely sown cotton crop undertake inter culture operation (weeding / hoeing) during clear weather for management of weeds and improved soil aeration. • For weed control in timely sown cotton, alternative option of recommended early post emergence herbicidal spray (with label claim) at proper stage and proper soil moisture condition should be undertaken.
Pigeon pea	Growth	• Need based gap filling may be undertaken to maintain optimum plant population in irrigated fields. • In timely sown pigeon pea crop undertake inter culture operation (weeding / hoeing) during clear weather for management of weeds and improved soil aeration.
Green gram / Black gram	Growth	• In timely sown green gram crop undertake inter culture operation (weeding / hoeing) during clear weather for management of weeds and improved soil aeration.
Sorghum	Growth	• Need based gap filling may be undertaken to maintain optimum plant population in irrigated fields. • In timely sown sorghum crop undertake inter culture operation (weeding / hoeing) during clear weather for management of weeds and improved soil aeration.
Maize	Growth	• Need based gap filling may be undertaken to maintain optimum plant population in irrigated fields. In timely sown maize crop undertake inter culture operation (weeding / hoeing) during clear weather for management of weeds and improved soil aeration.
Fruit crops	Plantation	• Complete the plantation of all fruit crops in the area with sufficient monsoon rains. • In ambia bahar mandarin and sweet orange orchards, damaged twigs from sudden rain/thunderstorm be removed and cut ends be smeared with Bordeaux paste (1 kg lime + 1 kg CuSo4 + 10 liter water). • During this month there is rapid spread of citrus canker in acid lime (lemon) hence the affected leaves, branches should be pruned and burnt. Spray copper oxychloride180 gm with streptocycline 6 gm in 60 liters of water before the onset of the monsoon. Repeat the spray 3-4 times at an interval of 30 days.
Vegetable	Trans plantation / Nursery	• Before transplanting chilly seedlings, dip the tops of the seedlings in a solution of 10 ml of Dimethoate 30% + 3 g of water soluble sulphur (80%) + 2.5 g of Mancozeb in one liter of water. • Before transplanting brinjal seedlings, dip the tops of the seedlings in a solution of 10 ml of 30% dimethoate per 10 liters of water.
Livestock	-	• Ensure sufficient feed and medication for livestock to last through prevailing intermittent disruption. • Undertake maintenance and repair of animal shelter and water points. Also ensure proper care and management of livestock during rainy season. • External and Internal parasite cases increase in rainy season so take care of it with consulting the veterinary doctor. • Enrichment of poultry feed with vitamin "C" and "E" is advisable.

This Agro Advisory Bulletin (AAB) is prepared & published with the consultation and recommendation of SMS committee of "Gramin Krishi Mausam Sewa" (GKMS) Dr. Panjabrao Deshmukh Krishi Vidyapeeth, Akola. 444005 (MS).

Nodal Officer
Gramin Krishi Mausam Sewa (GKMS)
Dr Panjabrao Deshmukh Krishi Vidyapeeth, Akola

Technical Officer
Gramin Krishi Mausam Sewa (GKMS)
Dr Panjabrao Deshmukh Krishi Vidyapeeth, Akola