

Taylor's Method

Taylor's Method

First Order ODEs: $y' = f(x, y)$

Taylor's Method

First Order ODEs: $y' = f(x, y)$

Initial Condition: $y(x_i) = y_i$

Taylor's Method

First Order ODEs: $y' = f(x, y)$

Initial Condition: $y(x_i) = y_i$

$y(x_i)$ = Exact Solution

y_i = Numerical Solution


Taylor's Method

First Order ODEs: $y' = f(x, y)$

Initial Condition: $y(x_i) = y_i$

$y(x_i)$ = Exact Solution

y_i = Numerical Solution



Taylor's Method

First Order ODEs:

$$y' = f(x, y)$$

Initial Condition:


$$y(x_i) = y_i$$

$y(x_i)$ = Exact Solution

y_i = Numerical Solution

Taylor Series:

$$y(x_{i+1}) = y(x_i + h)$$



Taylor's Method

First Order ODEs: $y' = f(x, y)$


Initial Condition: $y(x_i) = y_i$

$y(x_i)$ = Exact Solution

y_i = Numerical Solution

Taylor Series:

$$y(x_{i+1}) = y(x_i + h)$$



Taylor's Method

First Order ODEs: $y' = f(x, y)$


Initial Condition: $y(x_i) = y_i$

$y(x_i)$ = Exact Solution

y_i = Numerical Solution

Taylor Series:

$$\begin{aligned} y(x_{i+1}) &= y(x_i + h) \\ &= y(x_i) + h \cdot y'(x_i) + \frac{h^2}{2!} \cdot y''(x_i) + \frac{h^3}{3!} \cdot y^{(3)}(x_i) + \dots \end{aligned}$$



Taylor's Method

First Order ODEs: $y' = f(x, y)$


Initial Condition: $y(x_i) = y_i$

$y(x_i)$ = Exact Solution

y_i = Numerical Solution

Taylor Series:

$$\begin{aligned} y(x_{i+1}) &= y(x_i + h) \\ &= y(x_i) + h \cdot y'(x_i) + \frac{h^2}{2!} \cdot y''(x_i) + \frac{h^3}{3!} \cdot y^{(3)}(x_i) + \dots \end{aligned}$$



Taylor's Method

First Order ODEs: $y' = f(x, y)$

Initial Condition: $y(x_i) = y_i$

$y(x_i)$ = Exact Solution


y_i = Numerical Solution

Taylor Series:

$$\begin{aligned} y(x_{i+1}) &= y(x_i + h) \\ &= y(x_i) + h \cdot y'(x_i) + \frac{h^2}{2!} \cdot y''(x_i) + \frac{h^3}{3!} \cdot y^{(3)}(x_i) + \dots \end{aligned}$$

Taylor's Method:

$$y_{i+1} = y_i + h \cdot y'_i + \frac{h^2}{2!} \cdot y''_i + \frac{h^3}{3!} \cdot y^{(3)}_i + \dots$$



Taylor's Method

First Order ODEs: $y' = f(x, y)$

Initial Condition: $y(x_i) = y_i$

$y(x_i)$ = Exact Solution


y_i = Numerical Solution

Taylor Series:

$$\begin{aligned} y(x_{i+1}) &= y(x_i + h) \\ &= y(x_i) + h \cdot y'(x_i) + \frac{h^2}{2!} \cdot y''(x_i) + \frac{h^3}{3!} \cdot y^{(3)}(x_i) + \dots \end{aligned}$$

Taylor's Method:

$$\begin{aligned} y_{i+1} &= y_i + h \cdot y'_i + \frac{h^2}{2!} \cdot y''_i + \frac{h^3}{3!} \cdot y^{(3)}_i + \dots \\ &= y_i + h \cdot f(x_i, y_i) + \frac{h^2}{2!} \cdot f'(x_i, y_i) + \frac{h^3}{3!} \cdot f''(x_i, y_i) + \dots \end{aligned}$$



Taylor's Method

First Order ODEs: $y' = f(x, y)$

Initial Condition: $y(x_i) = y_i$

$y(x_i)$ = Exact Solution


y_i = Numerical Solution

Taylor Series:

$$\begin{aligned} y(x_{i+1}) &= y(x_i + h) \\ &= y(x_i) + h \cdot y'(x_i) + \frac{h^2}{2!} \cdot y''(x_i) + \frac{h^3}{3!} \cdot y^{(3)}(x_i) + \dots \end{aligned}$$

Taylor's Method:

$$\begin{aligned} y_{i+1} &= y_i + h \cdot y'_i + \frac{h^2}{2!} \cdot y''_i + \frac{h^3}{3!} \cdot y^{(3)}_i + \dots \\ &= y_i + h \cdot f(x_i, y_i) + \frac{h^2}{2!} \cdot f'(x_i, y_i) + \frac{h^3}{3!} \cdot f''(x_i, y_i) + \dots \end{aligned}$$



First Order: $y_{i+1} = y_i + h \cdot f(x_i, y_i)$

Taylor's Method

First Order ODEs: $y' = f(x, y)$

Initial Condition: $y(x_i) = y_i$

$y(x_i)$ = Exact Solution


y_i = Numerical Solution

Taylor Series:

$$\begin{aligned} y(x_{i+1}) &= y(x_i + h) \\ &= y(x_i) + h \cdot y'(x_i) + \frac{h^2}{2!} \cdot y''(x_i) + \frac{h^3}{3!} \cdot y^{(3)}(x_i) + \dots \end{aligned}$$

Taylor's Method:

$$\begin{aligned} y_{i+1} &= y_i + h \cdot y'_i + \frac{h^2}{2!} \cdot y''_i + \frac{h^3}{3!} \cdot y^{(3)}_i + \dots \\ &= y_i + h \cdot f(x_i, y_i) + \frac{h^2}{2!} \cdot f'(x_i, y_i) + \frac{h^3}{3!} \cdot f''(x_i, y_i) + \dots \end{aligned}$$



First Order: $y_{i+1} = y_i + h \cdot f(x_i, y_i)$

Taylor's Method

First Order ODEs: $y' = f(x, y)$

Initial Condition: $y(x_i) = y_i$

$y(x_i)$ = Exact Solution


y_i = Numerical Solution

Taylor Series:

$$\begin{aligned} y(x_{i+1}) &= y(x_i + h) \\ &= y(x_i) + h \cdot y'(x_i) + \frac{h^2}{2!} \cdot y''(x_i) + \frac{h^3}{3!} \cdot y^{(3)}(x_i) + \dots \end{aligned}$$

Taylor's Method:

$$\begin{aligned} y_{i+1} &= y_i + h \cdot y'_i + \frac{h^2}{2!} \cdot y''_i + \frac{h^3}{3!} \cdot y^{(3)}_i + \dots \\ &= y_i + h \cdot f(x_i, y_i) + \frac{h^2}{2!} \cdot f'(x_i, y_i) + \frac{h^3}{3!} \cdot f''(x_i, y_i) + \dots \end{aligned}$$



First Order: $y_{i+1} = y_i + h \cdot f(x_i, y_i)$

Taylor's Method

First Order ODEs: $y' = f(x, y)$

Initial Condition: $y(x_i) = y_i$

$y(x_i)$ = Exact Solution


y_i = Numerical Solution

Taylor Series:

$$\begin{aligned} y(x_{i+1}) &= y(x_i + h) \\ &= y(x_i) + h \cdot y'(x_i) + \frac{h^2}{2!} \cdot y''(x_i) + \frac{h^3}{3!} \cdot y^{(3)}(x_i) + \dots \end{aligned}$$

Taylor's Method:

$$\begin{aligned} y_{i+1} &= y_i + h \cdot y'_i + \frac{h^2}{2!} \cdot y''_i + \frac{h^3}{3!} \cdot y^{(3)}_i + \dots \\ &= y_i + h \cdot f(x_i, y_i) + \frac{h^2}{2!} \cdot f'(x_i, y_i) + \frac{h^3}{3!} \cdot f''(x_i, y_i) + \dots \end{aligned}$$



First Order: $y_{i+1} = y_i + h \cdot f(x_i, y_i)$

Second Order: $y_{i+1} = y_i + h \cdot f(x_i, y_i) + \frac{h^2}{2!} \cdot f'(x_i, y_i)$

Taylor's Method

First Order ODEs: $y' = f(x, y)$

Initial Condition: $y(x_i) = y_i$

$y(x_i)$ = Exact Solution


y_i = Numerical Solution

Taylor Series:

$$\begin{aligned} y(x_{i+1}) &= y(x_i + h) \\ &= y(x_i) + h \cdot y'(x_i) + \frac{h^2}{2!} \cdot y''(x_i) + \frac{h^3}{3!} \cdot y^{(3)}(x_i) + \dots \end{aligned}$$

Taylor's Method:

$$\begin{aligned} y_{i+1} &= y_i + h \cdot y'_i + \frac{h^2}{2!} \cdot y''_i + \frac{h^3}{3!} \cdot y^{(3)}_i + \dots \\ &= y_i + h \cdot f(x_i, y_i) + \frac{h^2}{2!} \cdot f'(x_i, y_i) + \frac{h^3}{3!} \cdot f''(x_i, y_i) + \dots \end{aligned}$$



First Order: $y_{i+1} = y_i + h \cdot f(x_i, y_i)$

Second Order: $y_{i+1} = y_i + h \cdot f(x_i, y_i) + \frac{h^2}{2!} \cdot f'(x_i, y_i)$

$\vdots$

Taylor's Method

Example 07-01: Approximate the numerical solution of the following initial value problem using Taylor's method of order 1,2,4 for $0 \leq x \leq 1$ with step size $h = 0.2$.

$$y' = -2xy^2, y(0) = 1; \quad 0 \leq x \leq 1$$

Taylor's Method

Example 07-01: Approximate the numerical solution of the following initial value problem using Taylor's method of order 1,2,4 for $0 \leq x \leq 1$ with step size $h = 0.2$.

$$y' = -2xy^2, y(0) = 1; \quad 0 \leq x \leq 1$$

Exact Solution: $y(x) = \frac{1}{1+x^2}$

Taylor's Method

Example 07-01: Approximate the numerical solution of the following initial value problem using Taylor's method of order 1,2,4 for $0 \leq x \leq 1$ with step size $h = 0.2$.

$$y' = -2xy^2, y(0) = 1; \quad 0 \leq x \leq 1$$

Exact Solution: $y(x) = \frac{1}{1+x^2}$

$$f(x, y) = -2xy^2$$

Taylor's Method

Example 07-01: Approximate the numerical solution of the following initial value problem using Taylor's method of order 1,2,4 for $0 \leq x \leq 1$ with step size $h = 0.2$.

$$y' = -2xy^2, y(0) = 1; \quad 0 \leq x \leq 1$$

Exact Solution: $y(x) = \frac{1}{1+x^2}$

$$f(x, y) = -2xy^2, N = \frac{1 - 0}{0.2} = 5$$

Taylor's Method

Example 07-01: Approximate the numerical solution of the following initial value problem using Taylor's method of order 1,2,4 for $0 \leq x \leq 1$ with step size $h = 0.2$.

$$y' = -2xy^2, y(0) = 1; \quad 0 \leq x \leq 1$$

Exact Solution: $y(x) = \frac{1}{1+x^2}$

$$f(x, y) = -2xy^2, N = \frac{1 - 0}{0.2} = 5$$

Taylor's Method

Example 07-01: Approximate the numerical solution of the following initial value problem using Taylor's method of order 1,2,4 for $0 \leq x \leq 1$ with step size $h = 0.2$.

$$y' = -2xy^2, y(0) = 1; \quad 0 \leq x \leq 1$$

Exact Solution: $y(x) = \frac{1}{1+x^2}$

$$f(x, y) = -2xy^2, N = \frac{1 - 0}{0.2} = 5$$

Taylor's Method

Example 07-01: Approximate the numerical solution of the following initial value problem using Taylor's method of order 1,2,4 for $0 \leq x \leq 1$ with step size $h = 0.2$.

$$y' = -2xy^2, y(0) = 1; \quad 0 \leq x \leq 1$$

Exact Solution: $y(x) = \frac{1}{1+x^2}$

$$f(x, y) = -2xy^2, N = \frac{1 - 0}{0.2} = 5$$

First Order:

($n = 1$)

Taylor's Method

Example 07-01: Approximate the numerical solution of the following initial value problem using Taylor's method of order 1,2,4 for $0 \leq x \leq 1$ with step size $h = 0.2$.

$$y' = -2xy^2, y(0) = 1; \quad 0 \leq x \leq 1$$

Exact Solution: $y(x) = \frac{1}{1+x^2}$

$$f(x, y) = -2xy^2, N = \frac{1 - 0}{0.2} = 5$$

First Order:

$$y_{i+1} = y_i + h \cdot f(x_i, y_i)$$

($n = 1$)

Taylor's Method

Example 07-01: Approximate the numerical solution of the following initial value problem using Taylor's method of order 1,2,4 for $0 \leq x \leq 1$ with step size $h = 0.2$.

$$y' = -2xy^2, y(0) = 1; \quad 0 \leq x \leq 1$$

Exact Solution: $y(x) = \frac{1}{1+x^2}$

$$f(x, y) = -2xy^2, \quad N = \frac{1 - 0}{0.2} = 5$$

First Order:

($n = 1$)

$$y_{i+1} = y_i + h \cdot f(x_i, y_i)$$

$$y_{i+1} = y_i + h \cdot (-2x_i y_i^2)$$

Taylor's Method

Example 07-01: Approximate the numerical solution of the following initial value problem using Taylor's method of order 1,2,4 for $0 \leq x \leq 1$ with step size $h = 0.2$.

$$y' = -2xy^2, y(0) = 1; \quad 0 \leq x \leq 1$$

Exact Solution: $y(x) = \frac{1}{1+x^2}$

$$f(x, y) = -2xy^2, \quad N = \frac{1 - 0}{0.2} = 5$$

First Order:

($n = 1$)

$$y_{i+1} = y_i + h \cdot f(x_i, y_i)$$

$$y_1 = y_0 + h \cdot (-2x_0y_0^2)$$

$$y_{i+1} = y_i + h \cdot (-2x_iy_i^2)$$

Taylor's Method

Example 07-01: Approximate the numerical solution of the following initial value problem using Taylor's method of order 1,2,4 for $0 \leq x \leq 1$ with step size $h = 0.2$.

$$y' = -2xy^2, y(0) = 1; \quad 0 \leq x \leq 1$$

Exact Solution: $y(x) = \frac{1}{1+x^2}$

$$f(x, y) = -2xy^2, \quad N = \frac{1 - 0}{0.2} = 5$$

First Order:

($n = 1$)

$$y_{i+1} = y_i + h \cdot f(x_i, y_i)$$

$$y_{i+1} = y_i + h \cdot (-2x_i y_i^2)$$

$$y_1 = y_0 + h \cdot (-2x_0 y_0^2)$$

$$= 1 + 0.2 \cdot [-2 \cdot (0) \cdot (1)^2]$$

Taylor's Method

Example 07-01: Approximate the numerical solution of the following initial value problem using Taylor's method of order 1,2,4 for $0 \leq x \leq 1$ with step size $h = 0.2$.

$$y' = -2xy^2, y(0) = 1; \quad 0 \leq x \leq 1$$

Exact Solution: $y(x) = \frac{1}{1+x^2}$

$$f(x, y) = -2xy^2, N = \frac{1-0}{0.2} = 5$$

First Order:

($n = 1$)

$$y_{i+1} = y_i + h \cdot f(x_i, y_i)$$

$$y_{i+1} = y_i + h \cdot (-2x_i y_i^2)$$

$$\begin{aligned} y_1 &= y_0 + h \cdot (-2x_0 y_0^2) \\ &= 1 + 0.2 \cdot [-2 \cdot (0) \cdot (1)^2] \\ &= 1 \end{aligned}$$

Taylor's Method

Example 07-01: Approximate the numerical solution of the following initial value problem using Taylor's method of order 1,2,4 for $0 \leq x \leq 1$ with step size $h = 0.2$.

$$y' = -2xy^2, y(0) = 1; \quad 0 \leq x \leq 1$$

Exact Solution: $y(x) = \frac{1}{1+x^2}$

$$f(x, y) = -2xy^2, N = \frac{1-0}{0.2} = 5$$

First Order:

($n = 1$)

$$y_{i+1} = y_i + h \cdot f(x_i, y_i)$$

$$y_{i+1} = y_i + h \cdot (-2x_i y_i^2)$$

$$\begin{aligned} y_1 &= y_0 + h \cdot (-2x_0 y_0^2) \\ &= 1 + 0.2 \cdot [-2 \cdot (0) \cdot (1)^2] \\ &= 1 \end{aligned}$$

$$y_2 = y_1 + h \cdot (-2x_1 y_1^2)$$

Taylor's Method

Example 07-01: Approximate the numerical solution of the following initial value problem using Taylor's method of order 1,2,4 for $0 \leq x \leq 1$ with step size $h = 0.2$.

$$y' = -2xy^2, y(0) = 1; \quad 0 \leq x \leq 1$$

Exact Solution: $y(x) = \frac{1}{1+x^2}$

$$f(x, y) = -2xy^2, N = \frac{1-0}{0.2} = 5$$

First Order:

($n = 1$)

$$y_{i+1} = y_i + h \cdot f(x_i, y_i)$$

$$y_{i+1} = y_i + h \cdot (-2x_i y_i^2)$$

$$\begin{aligned} y_1 &= y_0 + h \cdot (-2x_0 y_0^2) \\ &= 1 + 0.2 \cdot [-2 \cdot (0) \cdot (1)^2] \\ &= 1 \end{aligned}$$

$$\begin{aligned} y_2 &= y_1 + h \cdot (-2x_1 y_1^2) \\ &= 1 + 0.2 \cdot [-2 \cdot (0.2) \cdot (1)^2] \end{aligned}$$

Taylor's Method

Example 07-01: Approximate the numerical solution of the following initial value problem using Taylor's method of order 1,2,4 for $0 \leq x \leq 1$ with step size $h = 0.2$.

$$y' = -2xy^2, y(0) = 1; \quad 0 \leq x \leq 1$$

Exact Solution: $y(x) = \frac{1}{1+x^2}$

$$f(x, y) = -2xy^2, N = \frac{1-0}{0.2} = 5$$

First Order:

($n = 1$)

$$y_{i+1} = y_i + h \cdot f(x_i, y_i)$$

$$y_{i+1} = y_i + h \cdot (-2x_i y_i^2)$$

$$\begin{aligned} y_1 &= y_0 + h \cdot (-2x_0 y_0^2) \\ &= 1 + 0.2 \cdot [-2 \cdot (0) \cdot (1)^2] \\ &= 1 \end{aligned}$$

$$\begin{aligned} y_2 &= y_1 + h \cdot (-2x_1 y_1^2) \\ &= 1 + 0.2 \cdot [-2 \cdot (0.2) \cdot (1)^2] \\ &= 0.92 \end{aligned}$$

Taylor's Method

Example 07-01: Approximate the numerical solution of the following initial value problem using Taylor's method of order 1,2,4 for $0 \leq x \leq 1$ with step size $h = 0.2$.

$$y' = -2xy^2, y(0) = 1; \quad 0 \leq x \leq 1$$

Exact Solution: $y(x) = \frac{1}{1+x^2}$

$$f(x, y) = -2xy^2, \quad N = \frac{1 - 0}{0.2} = 5$$

First Order:

($n = 1$)

$$y_{i+1} = y_i + h \cdot f(x_i, y_i)$$

$$y_{i+1} = y_i + h \cdot (-2x_i y_i^2)$$

$$\begin{aligned} y_1 &= y_0 + h \cdot (-2x_0 y_0^2) \\ &= 1 + 0.2 \cdot [-2 \cdot (0) \cdot (1)^2] \\ &= 1 \end{aligned}$$

$$\begin{aligned} y_2 &= y_1 + h \cdot (-2x_1 y_1^2) \\ &= 1 + 0.2 \cdot [-2 \cdot (0.2) \cdot (1)^2] \\ &= 0.92 \end{aligned}$$

Second Order:

($n = 2$)

Taylor's Method

Example 07-01: Approximate the numerical solution of the following initial value problem using Taylor's method of order 1,2,4 for $0 \leq x \leq 1$ with step size $h = 0.2$.

$$y' = -2xy^2, y(0) = 1; \quad 0 \leq x \leq 1$$

Exact Solution: $y(x) = \frac{1}{1+x^2}$

$$f(x, y) = -2xy^2, N = \frac{1 - 0}{0.2} = 5$$

First Order:

($n = 1$)

$$y_{i+1} = y_i + h \cdot f(x_i, y_i)$$

$$y_{i+1} = y_i + h \cdot (-2x_i y_i^2)$$

$$y_1 = y_0 + h \cdot (-2x_0 y_0^2)$$

$$= 1 + 0.2 \cdot [-2 \cdot (0) \cdot (1)^2]$$

$$= 1$$

$$y_2 = y_1 + h \cdot (-2x_1 y_1^2)$$

$$= 1 + 0.2 \cdot [-2 \cdot (0.2) \cdot (1)^2]$$

$$= 0.92$$

Second Order:

($n = 2$)

$$y_{i+1} = y_i + h \cdot f(x_i, y_i) + \frac{h^2}{2!} \cdot f'(x_i, y_i)$$

Taylor's Method

Example 07-01: Approximate the numerical solution of the following initial value problem using Taylor's method of order 1,2,4 for $0 \leq x \leq 1$ with step size $h = 0.2$.

$$y' = -2xy^2, y(0) = 1; \quad 0 \leq x \leq 1$$

$$\text{Exact Solution: } y(x) = \frac{1}{1+x^2}$$

$$f(x, y) = -2xy^2, N = \frac{1 - 0}{0.2} = 5$$

First Order:

($n = 1$)

$$y_{i+1} = y_i + h \cdot f(x_i, y_i)$$

$$y_{i+1} = y_i + h \cdot (-2x_i y_i^2)$$

$$y_1 = y_0 + h \cdot (-2x_0 y_0^2)$$

$$= 1 + 0.2 \cdot [-2 \cdot (0) \cdot (1)^2]$$

$$= 1$$

$$y_2 = y_1 + h \cdot (-2x_1 y_1^2)$$

$$= 1 + 0.2 \cdot [-2 \cdot (0.2) \cdot (1)^2]$$

$$= 0.92$$

Second Order:

($n = 2$)

$$y_{i+1} = y_i + h \cdot f(x_i, y_i) + \frac{h^2}{2!} \cdot f'(x_i, y_i)$$

$$f'(x, y) = \frac{df}{dx} = \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} \cdot \frac{dy}{dx} = \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} \cdot f$$

Taylor's Method

Example 07-01: Approximate the numerical solution of the following initial value problem using Taylor's method of order 1,2,4 for $0 \leq x \leq 1$ with step size $h = 0.2$.

$$y' = -2xy^2, y(0) = 1; \quad 0 \leq x \leq 1$$

$$\text{Exact Solution: } y(x) = \frac{1}{1+x^2}$$

$$f(x, y) = -2xy^2, \quad N = \frac{1 - 0}{0.2} = 5$$

First Order:

($n = 1$)

$$y_{i+1} = y_i + h \cdot f(x_i, y_i)$$

$$y_{i+1} = y_i + h \cdot (-2x_i y_i^2)$$

$$y_1 = y_0 + h \cdot (-2x_0 y_0^2)$$

$$= 1 + 0.2 \cdot [-2 \cdot (0) \cdot (1)^2]$$

$$= 1$$

$$y_2 = y_1 + h \cdot (-2x_1 y_1^2)$$

$$= 1 + 0.2 \cdot [-2 \cdot (0.2) \cdot (1)^2]$$

$$= 0.92$$

Second Order:

($n = 2$)

$$y_{i+1} = y_i + h \cdot f(x_i, y_i) + \frac{h^2}{2!} \cdot f'(x_i, y_i)$$

$$\begin{aligned} f'(x, y) &= \frac{df}{dx} = \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} \cdot \frac{dy}{dx} = \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} \cdot f \\ &= \frac{\partial}{\partial x}(-2xy^2) + \left[\frac{\partial}{\partial y}(-2xy^2) \right] \cdot (-2xy^2) \end{aligned}$$

Taylor's Method

Example 07-01: Approximate the numerical solution of the following initial value problem using Taylor's method of order 1,2,4 for $0 \leq x \leq 1$ with step size $h = 0.2$.

$$y' = -2xy^2, y(0) = 1; \quad 0 \leq x \leq 1$$

$$\text{Exact Solution: } y(x) = \frac{1}{1+x^2}$$

$$f(x, y) = -2xy^2, N = \frac{1-0}{0.2} = 5$$

First Order:

($n = 1$)

$$y_{i+1} = y_i + h \cdot f(x_i, y_i)$$

$$y_{i+1} = y_i + h \cdot (-2x_i y_i^2)$$

$$y_1 = y_0 + h \cdot (-2x_0 y_0^2)$$

$$= 1 + 0.2 \cdot [-2 \cdot (0) \cdot (1)^2]$$

$$= 1$$

$$y_2 = y_1 + h \cdot (-2x_1 y_1^2)$$

$$= 1 + 0.2 \cdot [-2 \cdot (0.2) \cdot (1)^2]$$

$$= 0.92$$

Second Order:

($n = 2$)

$$y_{i+1} = y_i + h \cdot f(x_i, y_i) + \frac{h^2}{2!} \cdot f'(x_i, y_i)$$

$$\begin{aligned} f'(x, y) &= \frac{df}{dx} = \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} \cdot \frac{dy}{dx} = \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} \cdot f \\ &= \frac{\partial}{\partial x}(-2xy^2) + \left[\frac{\partial}{\partial y}(-2xy^2) \right] \cdot (-2xy^2) \\ &= -2y^2 + (-4xy) \cdot (-2xy^2) \end{aligned}$$

Taylor's Method

Example 07-01: Approximate the numerical solution of the following initial value problem using Taylor's method of order 1,2,4 for $0 \leq x \leq 1$ with step size $h = 0.2$.

$$y' = -2xy^2, y(0) = 1; \quad 0 \leq x \leq 1$$

$$\text{Exact Solution: } y(x) = \frac{1}{1+x^2}$$

$$f(x, y) = -2xy^2, N = \frac{1-0}{0.2} = 5$$

First Order:

($n = 1$)

$$y_{i+1} = y_i + h \cdot f(x_i, y_i)$$

$$y_{i+1} = y_i + h \cdot (-2x_i y_i^2)$$

$$y_1 = y_0 + h \cdot (-2x_0 y_0^2)$$

$$= 1 + 0.2 \cdot [-2 \cdot (0) \cdot (1)^2]$$

$$= 1$$

$$y_2 = y_1 + h \cdot (-2x_1 y_1^2)$$

$$= 1 + 0.2 \cdot [-2 \cdot (0.2) \cdot (1)^2]$$

$$= 0.92$$

Second Order:

($n = 2$)

$$y_{i+1} = y_i + h \cdot f(x_i, y_i) + \frac{h^2}{2!} \cdot f'(x_i, y_i)$$

$$\begin{aligned} f'(x, y) &= \frac{df}{dx} = \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} \cdot \frac{dy}{dx} = \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} \cdot f \\ &= \frac{\partial}{\partial x}(-2xy^2) + \left[\frac{\partial}{\partial y}(-2xy^2) \right] \cdot (-2xy^2) \\ &= -2y^2 + (-4xy) \cdot (-2xy^2) = -2y^2 + 8x^2y^3 \end{aligned}$$

Taylor's Method

Example 07-01: Approximate the numerical solution of the following initial value problem using Taylor's method of order 1,2,4 for $0 \leq x \leq 1$ with step size $h = 0.2$.

$$y' = -2xy^2, y(0) = 1; \quad 0 \leq x \leq 1$$

$$\text{Exact Solution: } y(x) = \frac{1}{1+x^2}$$

$$f(x, y) = -2xy^2, N = \frac{1-0}{0.2} = 5$$

First Order:

($n = 1$)

$$y_{i+1} = y_i + h \cdot f(x_i, y_i)$$

$$y_{i+1} = y_i + h \cdot (-2x_i y_i^2)$$

$$y_1 = y_0 + h \cdot (-2x_0 y_0^2)$$

$$= 1 + 0.2 \cdot [-2 \cdot (0) \cdot (1)^2]$$

$$= 1$$

$$y_2 = y_1 + h \cdot (-2x_1 y_1^2)$$

$$= 1 + 0.2 \cdot [-2 \cdot (0.2) \cdot (1)^2]$$

$$= 0.92$$

Second Order:

($n = 2$)

$$y_{i+1} = y_i + h \cdot f(x_i, y_i) + \frac{h^2}{2!} \cdot f'(x_i, y_i)$$

$$y_{i+1} = y_i + h \cdot (-2x_i y_i^2) + \frac{h^2}{2!} \cdot (-2y_i^2 + 8x_i^2 y_i^3)$$

$$f'(x, y) = \frac{df}{dx} = \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} \cdot \frac{dy}{dx} = \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} \cdot f$$

$$= \frac{\partial}{\partial x}(-2xy^2) + \left[\frac{\partial}{\partial y}(-2xy^2) \right] \cdot (-2xy^2)$$

$$= -2y^2 + (-4xy) \cdot (-2xy^2) = -2y^2 + 8x^2 y^3$$

Taylor's Method

Example 07-01: Approximate the numerical solution of the following initial value problem using Taylor's method of order 1,2,4 for $0 \leq x \leq 1$ with step size $h = 0.2$.

$$y' = -2xy^2, y(0) = 1; \quad 0 \leq x \leq 1$$

$$\text{Exact Solution: } y(x) = \frac{1}{1+x^2}$$

$$f(x, y) = -2xy^2, N = \frac{1-0}{0.2} = 5$$

First Order:

($n = 1$)

$$y_{i+1} = y_i + h \cdot f(x_i, y_i)$$

$$y_{i+1} = y_i + h \cdot (-2x_i y_i^2)$$

$$y_1 = y_0 + h \cdot (-2x_0 y_0^2)$$

$$= 1 + 0.2 \cdot [-2 \cdot (0) \cdot (1)^2]$$

$$= 1$$

$$y_2 = y_1 + h \cdot (-2x_1 y_1^2)$$

$$= 1 + 0.2 \cdot [-2 \cdot (0.2) \cdot (1)^2]$$

$$= 0.92$$

Second Order:

($n = 2$)

$$y_{i+1} = y_i + h \cdot f(x_i, y_i) + \frac{h^2}{2!} \cdot f'(x_i, y_i)$$

$$y_{i+1} = y_i + h \cdot (-2x_i y_i^2) + \frac{h^2}{2!} \cdot (-2y_i^2 + 8x_i^2 y_i^3)$$

$$y_1 = y_0 + h \cdot (-2x_0 y_0^2) + \frac{h^2}{2!} \cdot (-2y_0^2 + 8x_0^2 y_0^3)$$

$$f'(x, y) = \frac{df}{dx} = \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} \cdot \frac{dy}{dx} = \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} \cdot f$$

$$= \frac{\partial}{\partial x} (-2xy^2) + \left[\frac{\partial}{\partial y} (-2xy^2) \right] \cdot (-2xy^2)$$

$$= -2y^2 + (-4xy) \cdot (-2xy^2) = -2y^2 + 8x^2 y^3$$

Taylor's Method

Example 07-01: Approximate the numerical solution of the following initial value problem using Taylor's method of order 1,2,4 for $0 \leq x \leq 1$ with step size $h = 0.2$.

$$y' = -2xy^2, y(0) = 1; \quad 0 \leq x \leq 1$$

$$\text{Exact Solution: } y(x) = \frac{1}{1+x^2}$$

$$f(x, y) = -2xy^2, N = \frac{1-0}{0.2} = 5$$

First Order:

($n = 1$)

$$y_{i+1} = y_i + h \cdot f(x_i, y_i)$$

$$y_{i+1} = y_i + h \cdot (-2x_i y_i^2)$$

$$y_1 = y_0 + h \cdot (-2x_0 y_0^2)$$

$$= 1 + 0.2 \cdot [-2 \cdot (0) \cdot (1)^2]$$

$$= 1$$

$$y_2 = y_1 + h \cdot (-2x_1 y_1^2)$$

$$= 1 + 0.2 \cdot [-2 \cdot (0.2) \cdot (1)^2]$$

$$= 0.92$$

Second Order:

($n = 2$)

$$y_{i+1} = y_i + h \cdot f(x_i, y_i) + \frac{h^2}{2!} \cdot f'(x_i, y_i)$$

$$y_{i+1} = y_i + h \cdot (-2x_i y_i^2) + \frac{h^2}{2!} \cdot (-2y_i^2 + 8x_i^2 y_i^3)$$

$$y_1 = y_0 + h \cdot (-2x_0 y_0^2) + \frac{h^2}{2!} \cdot (-2y_0^2 + 8x_0^2 y_0^3)$$

$$= 1 + 0.2 \cdot [-2 \cdot (0) \cdot (1)^2] + \frac{(0.2)^2}{2!} [-2 \cdot (1)^2 + 8 \cdot (0)^2 \cdot (1)^3]$$

$$f'(x, y) = \frac{df}{dx} = \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} \cdot \frac{dy}{dx} = \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} \cdot f$$

$$= \frac{\partial}{\partial x} (-2xy^2) + \left[\frac{\partial}{\partial y} (-2xy^2) \right] \cdot (-2xy^2)$$

$$= -2y^2 + (-4xy) \cdot (-2xy^2) = -2y^2 + 8x^2 y^3$$

Taylor's Method

Example 07-01: Approximate the numerical solution of the following initial value problem using Taylor's method of order 1,2,4 for $0 \leq x \leq 1$ with step size $h = 0.2$.

$$y' = -2xy^2, y(0) = 1; \quad 0 \leq x \leq 1$$

$$\text{Exact Solution: } y(x) = \frac{1}{1+x^2}$$

$$f(x, y) = -2xy^2, N = \frac{1-0}{0.2} = 5$$

First Order:

($n = 1$)

$$y_{i+1} = y_i + h \cdot f(x_i, y_i)$$

$$y_{i+1} = y_i + h \cdot (-2x_i y_i^2)$$

$$y_1 = y_0 + h \cdot (-2x_0 y_0^2)$$

$$= 1 + 0.2 \cdot [-2 \cdot (0) \cdot (1)^2]$$

$$= 1$$

$$y_2 = y_1 + h \cdot (-2x_1 y_1^2)$$

$$= 1 + 0.2 \cdot [-2 \cdot (0.2) \cdot (1)^2]$$

$$= 0.92$$

Second Order:

($n = 2$)

$$y_{i+1} = y_i + h \cdot f(x_i, y_i) + \frac{h^2}{2!} \cdot f'(x_i, y_i)$$

$$y_{i+1} = y_i + h \cdot (-2x_i y_i^2) + \frac{h^2}{2!} \cdot (-2y_i^2 + 8x_i^2 y_i^3)$$

$$y_1 = y_0 + h \cdot (-2x_0 y_0^2) + \frac{h^2}{2!} \cdot (-2y_0^2 + 8x_0^2 y_0^3)$$

$$= 1 + 0.2 \cdot [-2 \cdot (0) \cdot (1)^2] + \frac{(0.2)^2}{2!} [-2 \cdot (1)^2 + 8 \cdot (0)^2 \cdot (1)^3]$$

$$= 0.96$$

$$f'(x, y) = \frac{df}{dx} = \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} \cdot \frac{dy}{dx} = \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} \cdot f$$

$$= \frac{\partial}{\partial x} (-2xy^2) + \left[\frac{\partial}{\partial y} (-2xy^2) \right] \cdot (-2xy^2)$$

$$= -2y^2 + (-4xy) \cdot (-2xy^2) = -2y^2 + 8x^2 y^3$$

Taylor's Method

Example 07-01: Approximate the numerical solution of the following initial value problem using Taylor's method of order 1,2,4 for $0 \leq x \leq 1$ with step size $h = 0.2$.

$$y' = -2xy^2, y(0) = 1; \quad 0 \leq x \leq 1$$

$$\text{Exact Solution: } y(x) = \frac{1}{1+x^2}$$

$$f(x, y) = -2xy^2, N = \frac{1-0}{0.2} = 5$$

First Order:

($n = 1$)

$$y_{i+1} = y_i + h \cdot f(x_i, y_i)$$

$$y_{i+1} = y_i + h \cdot (-2x_i y_i^2)$$

$$y_1 = y_0 + h \cdot (-2x_0 y_0^2)$$

$$= 1 + 0.2 \cdot [-2 \cdot (0) \cdot (1)^2]$$

$$= 1$$

$$y_2 = y_1 + h \cdot (-2x_1 y_1^2)$$

$$= 1 + 0.2 \cdot [-2 \cdot (0.2) \cdot (1)^2]$$

$$= 0.92$$

Second Order:

($n = 2$)

$$y_{i+1} = y_i + h \cdot f(x_i, y_i) + \frac{h^2}{2!} \cdot f'(x_i, y_i)$$

$$y_{i+1} = y_i + h \cdot (-2x_i y_i^2) + \frac{h^2}{2!} \cdot (-2y_i^2 + 8x_i^2 y_i^3)$$

$$f'(x, y) = \frac{df}{dx} = \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} \cdot \frac{dy}{dx} = \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} \cdot f$$

$$= \frac{\partial}{\partial x}(-2xy^2) + \left[\frac{\partial}{\partial y}(-2xy^2) \right] \cdot (-2xy^2)$$

$$= -2y^2 + (-4xy) \cdot (-2xy^2) = -2y^2 + 8x^2 y^3$$

Taylor's Method

Example 07-01: Approximate the numerical solution of the following initial value problem using Taylor's method of order 1,2,4 for $0 \leq x \leq 1$ with step size $h = 0.2$.

$$y' = -2xy^2, y(0) = 1; \quad 0 \leq x \leq 1$$

$$\text{Exact Solution: } y(x) = \frac{1}{1+x^2}$$

$$f(x, y) = -2xy^2, N = \frac{1 - 0}{0.2} = 5$$

First Order:

($n = 1$)

$$y_{i+1} = y_i + h \cdot f(x_i, y_i)$$

$$y_{i+1} = y_i + h \cdot (-2x_i y_i^2)$$

$$y_1 = y_0 + h \cdot (-2x_0 y_0^2)$$

$$= 1 + 0.2 \cdot [-2 \cdot (0) \cdot (1)^2]$$

$$= 1$$

$$y_2 = y_1 + h \cdot (-2x_1 y_1^2)$$

$$= 1 + 0.2 \cdot [-2 \cdot (0.2) \cdot (1)^2]$$

$$= 0.92$$

Second Order:

($n = 2$)

$$y_{i+1} = y_i + h \cdot f(x_i, y_i) + \frac{h^2}{2!} \cdot f'(x_i, y_i)$$

$$y_{i+1} = y_i + h \cdot (-2x_i y_i^2) + \frac{h^2}{2!} \cdot (-2y_i^2 + 8x_i^2 y_i^3)$$

$$y_2 = y_1 + h \cdot (-2x_1 y_1^2) + \frac{h^2}{2!} \cdot (-2y_1^2 + 8x_1^2 y_1^3)$$

$$f'(x, y) = \frac{df}{dx} = \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} \cdot \frac{dy}{dx} = \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} \cdot f$$

$$= \frac{\partial}{\partial x} (-2xy^2) + \left[\frac{\partial}{\partial y} (-2xy^2) \right] \cdot (-2xy^2)$$

$$= -2y^2 + (-4xy) \cdot (-2xy^2) = -2y^2 + 8x^2 y^3$$

Taylor's Method

Example 07-01: Approximate the numerical solution of the following initial value problem using Taylor's method of order 1,2,4 for $0 \leq x \leq 1$ with step size $h = 0.2$.

$$y' = -2xy^2, y(0) = 1; \quad 0 \leq x \leq 1$$

$$\text{Exact Solution: } y(x) = \frac{1}{1+x^2}$$

$$f(x, y) = -2xy^2, N = \frac{1-0}{0.2} = 5$$

First Order:

($n = 1$)

$$y_{i+1} = y_i + h \cdot f(x_i, y_i)$$

$$y_{i+1} = y_i + h \cdot (-2x_i y_i^2)$$

$$y_1 = y_0 + h \cdot (-2x_0 y_0^2)$$

$$= 1 + 0.2 \cdot [-2 \cdot (0) \cdot (1)^2]$$

$$= 1$$

$$y_2 = y_1 + h \cdot (-2x_1 y_1^2)$$

$$= 1 + 0.2 \cdot [-2 \cdot (0.2) \cdot (1)^2]$$

$$= 0.92$$

Second Order:

($n = 2$)

$$y_{i+1} = y_i + h \cdot f(x_i, y_i) + \frac{h^2}{2!} \cdot f'(x_i, y_i)$$

$$y_{i+1} = y_i + h \cdot (-2x_i y_i^2) + \frac{h^2}{2!} \cdot (-2y_i^2 + 8x_i^2 y_i^3)$$

$$y_2 = y_1 + h \cdot (-2x_1 y_1^2) + \frac{h^2}{2!} \cdot (-2y_1^2 + 8x_1^2 y_1^3)$$

$$= 0.96 + 0.2 \cdot [-2 \cdot (0.2) \cdot (0.96)^2] + \frac{(0.2)^2}{2!} [-2 \cdot (0.96)^2 + 8 \cdot (0.2)^2 \cdot (0.96)^3]$$

$$f'(x, y) = \frac{df}{dx} = \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} \cdot \frac{dy}{dx} = \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} \cdot f$$

$$= \frac{\partial}{\partial x} (-2xy^2) + \left[\frac{\partial}{\partial y} (-2xy^2) \right] \cdot (-2xy^2)$$

$$= -2y^2 + (-4xy) \cdot (-2xy^2) = -2y^2 + 8x^2 y^3$$

Taylor's Method

Example 07-01: Approximate the numerical solution of the following initial value problem using Taylor's method of order 1,2,4 for $0 \leq x \leq 1$ with step size $h = 0.2$.

$$y' = -2xy^2, y(0) = 1; \quad 0 \leq x \leq 1$$

$$\text{Exact Solution: } y(x) = \frac{1}{1+x^2}$$

$$f(x, y) = -2xy^2, N = \frac{1-0}{0.2} = 5$$

First Order:

($n = 1$)

$$y_{i+1} = y_i + h \cdot f(x_i, y_i)$$

$$y_{i+1} = y_i + h \cdot (-2x_i y_i^2)$$

$$y_1 = y_0 + h \cdot (-2x_0 y_0^2)$$

$$= 1 + 0.2 \cdot [-2 \cdot (0) \cdot (1)^2]$$

$$= 1$$

$$y_2 = y_1 + h \cdot (-2x_1 y_1^2)$$

$$= 1 + 0.2 \cdot [-2 \cdot (0.2) \cdot (1)^2]$$

$$= 0.92$$

Second Order:

($n = 2$)

$$y_{i+1} = y_i + h \cdot f(x_i, y_i) + \frac{h^2}{2!} \cdot f'(x_i, y_i)$$

$$y_{i+1} = y_i + h \cdot (-2x_i y_i^2) + \frac{h^2}{2!} \cdot (-2y_i^2 + 8x_i^2 y_i^3)$$

$$y_2 = y_1 + h \cdot (-2x_1 y_1^2) + \frac{h^2}{2!} \cdot (-2y_1^2 + 8x_1^2 y_1^3)$$

$$= 0.96 + 0.2 \cdot [-2 \cdot (0.2) \cdot (0.96)^2] + \frac{(0.2)^2}{2!} [-2 \cdot (0.96)^2 + 8 \cdot (0.2)^2 \cdot (0.96)^3]$$

$$= 0.855070$$

$$\begin{aligned} f'(x, y) &= \frac{df}{dx} = \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} \cdot \frac{dy}{dx} = \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} \cdot f \\ &= \frac{\partial}{\partial x} (-2xy^2) + \left[\frac{\partial}{\partial y} (-2xy^2) \right] \cdot (-2xy^2) \\ &= -2y^2 + (-4xy) \cdot (-2xy^2) = -2y^2 + 8x^2 y^3 \end{aligned}$$

