

Supporting Information for Using rapid temperature falls to estimate future strong cold front frequency in CMIP6 climate projections

Paola B Corrales^{1,2,3,4}, Martin Singh^{1,2,4}, Michael J. Reeder^{1,2}

¹School of Earth, Atmosphere and Environment, Monash University, Melbourne, Australia

²ARC Centre of Excellence for Climate Extremes, Melbourne, Australia

³Research School of Earth Sciences, Australian National University, Canberra, Australia

⁴ARC Centre of Excellence for the Weather of the 21st Century, Melbourne, Australia

Corresponding author: Paola B Corrales, paola.corrales@monash.edu




Figure S1: Multimodel mean change in the frequency of $\Delta T_{2.5}$ (days/season, SSP5-8.5 2018-2100 minus Historical 1980-2000) for a) DJF and b) JJA and ΔT values associated with the 2.5th percentile calculated for c) DJF and d) JJA. In a-b) dotted areas show where at least 90% of the models agree in the sign of change.




Figure S2: Mean change in the mean $\Delta T_{2.5}$ frequency and a-b) mean meridional temperature gradient $\partial \bar{T} / \partial y$ and c-d) the eddy kinetic energy of the storm track EKE for SSP5-8.5 (2080-2100) compared with the historical period (1980-2000) for each model and season a,c) DJF and b,d) JJA. Values are calculated for the northern (25° to 65°) and southern (-25° to -65°) hemispheres and over sea (circles) and land (triangles). Each solid line corresponds to a linear regression for each hemisphere and land or sea.