

A Simplified Model for Linear Mode Coupling in Multimode Fibers

This repo contains the code for the submission “A Simplified Model for Linear Mode Coupling in Multimode Fibers with Experimental Assessment” to the 2026 European Conference on Optical Communications.

Two multisectional models with the following transfer matrix for the i -th section are considered:

$$\mathbf{K}_i \mathbf{R}$$

where $\mathbf{R}$ is a block diagonal matrix with random unitary blocks uniformly distributed over the $U(N)$ group, whose blocks have the sizes of the mode groups, and

$$\mathbf{K}_i = e^{j\mathbf{B}_0 + \Delta\beta g(\text{XT}, G) (\mathbf{P}_i - \mathbf{P}_i^\dagger)}$$

where

$$\mathbf{B}_0$$

is the diagonal matrix containing the product between the section length L and the propagation constants $\beta_0^{(n)}$ (n is the mode index) for a graded-index multimode fiber (GIMMF) which are assumed to be constant within a mode group and to differ by $\Delta\beta$ from the neighbouring groups (the term $\Delta\beta L$ can also be regarded as an optimization parameter or kept fixed to a certain value, e.g., 10^4), and

$$\mathbf{P}_i$$

is a matrix whose elements are independent complex Gaussian random variables $P_{ab} \sim \mathcal{CN}(0, h_{A-B}^2)$.

In the paper, two choices for h_{A-B} are considered:

- 1) $h_{A-B} = 1$, i.e., same variance for all coupling coefficients. We call this the “uniform” choice. For $\text{XT} \leq -5$ dB, an approximate expression for g is

$$g(\text{XT}, G) \approx \sqrt{\frac{\text{XT}_t - 0.91 + 0.31G}{2M}}$$

where XT_t is the target crosstalk;

- 2) $h_{A-B} = e^{-2.25|A-B|}$. We call this the non-uniform choice. We don’t provide an approximate expression for g in this case, but rather a numerical one (see the first dataset listed in the Datasets section below) for $\text{XT} \in [-80, 0]$ dB and $G = 2, 3, \dots, 12$.

The models and the code are for a frequency-independent channel, the inclusion of frequency-dependence effects is immediate and explained in the paper.

Code

The repo is built around an abstract parent class `MultiSectionalModel` (see [localLibrary/](#)), which provides a common framework for simulating mode coupling in few-mode or multimode optical fibres using a transfer-matrix approach. The subclass `expmModel` implements the linear coupling model described above.

Other multisectional models can be implemented as subclasses of `MultiSectionalModel`, inheriting all fundamental functionalities such as Monte-Carlo simulations and computation of crosstalk metrics.

Example scripts

- [example1_single_section.m](#): Compute the transfer matrix and crosstalk statistics for a single section of the two models.
- [example2_concatenation.m](#): Generate the transfer matrix of a multisectional channel.
- [example3_tuningParameter.m](#): Obtain the numerical relation $g(XT, G)$ between the tuning parameter and the target crosstalk.

Datasets (in databases/)

- `single_segment_dB_vs_tuning_parameter_expm_exp_per_mode_a=-2.25_dBeta0=1e4_with_pol_with_igsc.mat`: $g(XT, G)$ for the non-uniform case, $XT \in [-80, 0]$ dB, $G = 2, \dots, 12$.
- `single_segment_dB_vs_tuning_parameter_expm_unif_dBeta0=1e4_with_pol_with_igsc.mat`: $g(XT, G)$ for the uniform case, with inter-group coupling.
- `single_segment_dB_vs_tuning_parameter_expm_unif_dBeta0=1e4_with_pol_no_igsc.mat`: $g(XT, G)$ for the uniform case, without inter-group coupling.

Requirements

Tested on MATLAB R2025b.

Citation

The paper has been accepted at the 2026 European Conference on Optical Communications (ECOC). A preprint is available on arXiv: [arXiv:2609.20617](#).

If you use this code, please cite:

P. Carniello, F. M. Ferreira, F. A. Barbosa, M.-J. Li, N. Hanik, “A Simplified Model for Linear Mode Coupling in Multimode Fibers with Experimental Assessment”, accepted at the 2026 European Conference on Optical Communications (ECOC), 2026. arXiv:2609.20617

```
@misc{carniello2026simplifiedmodellinearcode,  
  title={A Simplified Model for Linear Mode Coupling in  
  Multimode Fibers with Experimental Assessment},
```

```
author={Paolo Carniello and Filipe M. Ferreira and Fabio A.  
Barbosa and Ming-Jun Li and Norbert Hanik},  
year={2026},  
eprint={2609.20617},  
archivePrefix={arXiv},  
primaryClass={physics.optics},  
url={https://arxiv.org/abs/2609.20617},  
note={Accepted at the 2026 European Conference on Optical  
Communications (ECOC)},  
}
```