
A p -ADIC COHOMOLOGICAL APPROACH TO CONGRUENCES OF MEROMORPHIC MODULAR FORMS

by

Paolo Bordinon

Contents

1. Introduction.....	1
2. Geometric setting.....	2
3. Rigid cohomological description.....	4
4. Integral Frobenius structure.....	6
5. Arithmetic applications.....	8
References.....	11

1. Introduction

In this work, we provide a p -adic cohomological interpretation of certain congruences between Fourier coefficients of modular forms and Frobenius eigenvalues arising from the geometry of modular (and Shimura) curves, as studied in a recent work by Zhang [Zha25]. Zhang provided many numerical observations for congruences of the form

$$a_p\left(\frac{E_4}{j-j(C)}\right) \equiv a_p(C)^2 \pmod{p}$$

where $C/\mathbb{Q}$ is an elliptic curve with good reduction at p and $a_p(C) = p+1-|C(\mathbb{F}_p)|$. To study these relations, we introduce a cohomological framework and use the Gysin sequence to relate classes attached to modular forms to differential forms attached to the elliptic curves corresponding to the poles.

More precisely, let X be the elliptic or quaternionic modular curve of level N , defined over $\mathbb{Z}_p$ with $(p, N) = 1$. Consider ω , the line bundle obtained from the universal object whose global sections are modular forms. Let $\mathcal{H}$ be the vector bundle of rank 2 obtained from the relative de Rham cohomology. To relate meromorphic modular forms with prescribed structure at the poles, we use de Rham cohomology with coefficients and relate classes in $H_{\text{dR}}^1(X \setminus \alpha, \mathcal{H})$ with fibers $\mathcal{H}|_{\alpha}$ for α a $\mathbb{Z}_p$ -point. Using comparison theorems between rigid and de Rham cohomology, we relate meromorphic modular forms to overconvergent p -adic modular forms. In this way, we can naturally study the Frobenius action on the various cohomology groups. Enlarging the set of poles considered will allow us to study an explicit Frobenius action on the meromorphic modular forms representing the cohomology classes and to relate it with the action of the U_p -operator. Theorem A is the core result of this paper and shows that the Frobenius eigenvalues of the de Rham cohomology at the poles are related to the U_p action on meromorphic modular forms.

Theorem A. — *Let X be a modular curve (elliptic or quaternionic) of level N defined over $\mathbb{Z}_p$, with $N > 3$, $p > 3$ and $(N, p) = 1$. Fix a $\mathbb{Z}_p$ -point α on the modular curve. Let k be a positive*

integer with $p > k + 1$, let $P(X), Q(X) \in \mathbb{Q}_p[X]$ be respectively the characteristic polynomials of the Frobenius action on $\mathrm{Sym}^k(\alpha^*\mathcal{H})[1]$ and on $H_{\mathrm{dR}}^1(X^{\mathrm{rig}}, \mathcal{H}_k^{\mathrm{rig}})$

$$P(X) = \det(\mathrm{Frob} - X | \mathrm{Sym}^k(\alpha^*\mathcal{H})[1]) = \sum_{i=0}^{k+1} c_i X^i,$$

$$Q(X) = \det(\mathrm{Frob} - X | H_{\mathrm{dR}}^1(X^{\mathrm{rig}}, \mathcal{H}_k)),$$

$$P(X)Q(X) = \sum_{i=0}^{d+k+1} e_i X^i$$

with $d = \dim_{\mathbb{Q}_p} M_{k+2} + \dim_{\mathbb{Q}_p} S_{k+2}$. Let $M = d + k + 1$, then for every $f \in M_{k+2}^{\mathrm{mero}, \{\alpha\}}$ we have

$$\sum_{i=0}^M e_{M-i} p^{M-i} U_p^i(f) \in \theta^{k+1} \left(M_{-k}^{\mathrm{mero}, \{\alpha\}} \right)$$

where θ is the differential operator $q \frac{d}{dq}$ in the elliptic case.

The integrality provided by crystalline cohomology then allows us to deduce congruence relations. In the elliptic setting, we can derive an explicit congruence at the level of q -expansions as shown in Corollary 1.

Several technical constructions are only sketched in this article; a complete and systematic treatment will appear in the author's PhD thesis.

1.1. An important note on a recent result. — A recent preprint of Allen, Long and Saad [ALS25] came out in November 2025 during the writing of this article. The authors analyze the same congruences between meromorphic modular forms and elliptic curves and give a proof of many of Zhang's conjectures. Their methods extend the classical work of Scholl [Sch85] and a more recent work by Kazalicki-Scholl [KS13] to the case of meromorphic modular forms. The works are similar in many aspects, in particular the initial cohomological setting and the Gysin sequence are used in fundamentally the same way. Our approach differs from theirs primarily in the use of log-crystalline cohomology to identify a Frobenius stable lattice and the use of an explicit Frobenius lift enlarging the set of poles to the set of supersingular points. This allows us to avoid computing with local expansions at the cusp at infinity, enlarge our framework to the Shimura setting and derive relations between Fourier coefficients only at the very end. On the other hand, their precise work takes care of the order of the poles of the meromorphic modular forms allowing them to obtain very sharp congruences on the cohomological subgroup.

After a very kind and fruitful exchange of emails with the authors, where they have looked at a first draft of this work and at the techniques involved, they allowed us to publish our version.

2. Geometric setting

Let $p > 3$ be a prime, let $N > 3$ be an integer coprime with p . Consider one of the following settings

- $X = X_1(N)$, the modular curve of level $\Gamma_1(N)$ defined over $\mathbb{Z}_p$ classifying elliptic curves with level structure,
- $X = X_B$ a Shimura curve defined over $\mathbb{Z}_p$ for B indefinite non-split quaternion algebra over $\mathbb{Q}$ of discriminant δ coprime with p and of level $V_1(N)$ classifying false elliptic curves with level structure.

We will denote by $\overline{X} = X \times k$ its special fiber, let $C \subset X(\mathbb{Z}_p)$ be the finite collection (possibly empty) of cusps. Let ω be the ample line bundle over X obtained from the pushforward of the differential bundle on the universal object. Let $(\mathcal{H}, \nabla)$ be the rank 2 vector bundle arising from the relative de Rham cohomology over X with logarithmic connection

$$\nabla : \mathcal{H} \rightarrow \mathcal{H} \otimes_{\mathbb{Z}_p} \Omega_X^1(\log C)$$

coming from the Gauss-Manin connection. In the quaternionic case we are fixing a choice of a non-trivial idempotent element of $M_2(\mathbb{Z}_p)$ and projecting the four dimensional vector bundle coming

from the relative de Rham cohomology to obtain $\mathcal{H}_k$ and analogously with ω . See [Kas04]. Denote by $\mathcal{H}_k$ its k -th symmetric tensor power $\mathcal{H}_k = \text{Sym}^k \mathcal{H}$.

We recall the standard properties of $\mathcal{H}_k$. The vector bundles ω and $\mathcal{H}$ fit in a short exact sequence

$$(1) \quad 0 \rightarrow \omega \rightarrow \mathcal{H} \rightarrow \omega^{-1} \rightarrow 0$$

inducing a $(k+1)$ -step decreasing filtration on $\mathcal{H}_k$

$$(2) \quad \mathcal{H}_k = \text{Fil}^0 \mathcal{H}_k \supseteq \text{Fil}^1 \mathcal{H}_k \supseteq \cdots \text{Fil}^k \mathcal{H}_k \cong \omega^k \supseteq \text{Fil}^{k+1} \mathcal{H}_k = 0.$$

The graded pieces of the filtration are isomorphic to

$$\text{Gr}_i \mathcal{H}_k = \text{Fil}^{k-i} \mathcal{H}_k / \text{Fil}^{k+1-i} \mathcal{H}_k \cong \omega^{k-2i}.$$

The connection obeys *Griffith's transversality*

$$\nabla \text{Fil}^{k-i} \mathcal{H}_k \subseteq \text{Fil}^{k-i-1} \mathcal{H}_k \otimes_{\mathcal{O}_X} \Omega_X^1(\log C)$$

and that these maps induce a *Kodaira-Spencer* type of isomorphism of $\mathcal{O}_X$ -modules for $p > k+1$

$$(3) \quad \nabla : \text{Gr}_i \mathcal{H}_k \cong \text{Gr}_{i+1} \mathcal{H}_k \otimes_X \Omega_X^1(\log C).$$

Following the usual notation, we write

$$M_k(X) = \Gamma(X, \omega^k), \quad S_k(X) = \Gamma(X, \omega^{k-2} \otimes \Omega_X^1).$$

We denote the complex arising from the connection by

$$\mathcal{H}_k^\bullet := [0 \rightarrow \mathcal{H}_k \rightarrow \mathcal{H}_k \otimes_{\mathcal{O}_X} \Omega_X^1]$$

and denote its hypercohomology groups by H_{dR}^i

$$H_{\text{dR}}^i(X, \mathcal{H}_k) := \mathbb{H}^i(X, \mathcal{H}_k^\bullet).$$

Lemma 1. — *We have a short exact sequence*

$$(4) \quad 0 \rightarrow \omega_{-k} \xrightarrow{\theta^{k+1}} \omega_{k+2} \rightarrow \mathcal{H}_k \otimes_X \Omega_X^1(\log C) / \nabla_k(\mathcal{H}_k) \rightarrow 0$$

where the right map is compatible with inclusions and an Eichler-Shimura short exact sequence of $\mathcal{O}_K$ -modules

$$0 \rightarrow M_{k+2}(X) \rightarrow H_{\text{dR}}^1(X, \mathcal{H}_k) \rightarrow S_{k+2}(X)^\vee \rightarrow 0.$$

Proof. — Consider the filtration $(\text{Fil}^p \mathcal{H}_k^\bullet)_p$ of the complex $\mathcal{H}_k^\bullet$

$$\text{Fil}^p \mathcal{H}_k^\bullet := [0 \rightarrow \text{Fil}^p \mathcal{H}_k \rightarrow \text{Fil}^{p-1} \mathcal{H}_k \otimes_{\mathcal{O}_X} \Omega_X^1(\log C)]$$

and take the associated spectral sequence $E_0^{p,q} = \text{Fil}^p \mathcal{H}_k^{p+q} / \text{Fil}^{p+1} \mathcal{H}_k^{p+q}$. The only terms that survive in page 1 are

$$E_1^{0,0} \cong \mathcal{H}_k / \text{Fil}^1 \mathcal{H}_k, \quad E_1^{k+1,-k} \cong \text{Fil}^k \mathcal{H}_k \otimes_{\mathcal{X}} \Omega_{\mathcal{X}}^1(\log C).$$

The spectral sequence degenerates at page $k+2$ converging to $E_{k+2}^{k+1,-k} = H^1(\mathcal{H}_k^\bullet)$. This gives rise to the short exact sequence

$$0 \rightarrow \mathcal{H}_k / \text{Fil}^1 \mathcal{H}_k \xrightarrow{\theta^{k+1}} \text{Fil}^k \mathcal{H}_k \otimes_{\mathcal{X}} \Omega_{\mathcal{X}}^1(\log C) \rightarrow \mathcal{H}_k \otimes_{\mathcal{X}} \Omega_{\mathcal{X}}^1(\log C) / \nabla_k(\mathcal{H}_k) \rightarrow 0.$$

The second exact sequence is obtained applying the spectral sequence to the hypercohomology of the graded pieces obtaining

$$\begin{aligned} 0 \rightarrow H_{\text{dR}}^0(X, \mathcal{H}_k^\bullet) &\rightarrow H^0(X, \omega^{-k}) \rightarrow H^0(X, \omega^k \otimes \Omega_X^1(\log C)) \rightarrow H_{\text{dR}}^1(X, \mathcal{H}_k^\bullet) \rightarrow \\ &\rightarrow H^1(X, \omega^{-k}) \rightarrow H^1(X, \omega^k \otimes \Omega_X^1(\log C)) \rightarrow \cdots \end{aligned}$$

giving the desired exact sequence using ampleness of ω and Serre duality. $\square$

Fix an effective divisor $S = \sum_{i=1}^n \alpha_i$ supported on the open curve $X \setminus C$ and defined over $\mathbb{Z}_p$,

$$\alpha_i : \text{Spec}(\mathcal{O}_K) \rightarrow X \setminus C$$

for K a finite field extension over $\mathbb{Q}_p$ with distinct close points on the special fiber $\overline{X}$. We denote by $\overline{S}$ their reduction to the special fiber $\overline{X}$ and $S_{\mathbb{Q}_p}$ their restriction to the generic fiber. We have that the curve $U = X_{\mathbb{Q}_p} \setminus S_{\mathbb{Q}_p}$ is affine.

As a direct consequence of the lemma, the de Rham cohomology of $\mathcal{H}_k$ can be computed by

$$H_{\text{dR}}^1(U, C), \mathcal{H}_k) = M_{k+2}^{\text{mero}, S} / \theta^{k+1} (M_{-k, \lambda}^{\text{mero}, S})$$

where we have used the notation $M_{k+2}^{\text{mero}, S} = \Gamma(U, \omega^{k+2})$ and should be thought as *meromorphic modular forms with poles along S* .

3. Rigid cohomological description

Consider X^{rig} to be the rigid analytification of X and the reduction map

$$\text{red} : X^{\text{rig}} \rightarrow \overline{X}(\overline{\mathbb{F}_p})$$

where we fixed $\overline{k}$ algebraic closure of k . We then consider the wide open obtained from X^{rig} removing closed balls in the residue disks of the points S . Since X is smooth, $\text{red}^{-1}(\overline{\alpha})$ is conformal to an open ball B for every $\overline{\alpha} \in \overline{S}$. Fixing this parametrization and fixing $0 < \lambda < 1$, define

$$V_\lambda = X^{\text{rig}} \setminus \bigcup_{\alpha \in S} B(\alpha, \lambda)$$

with $B(\alpha, \lambda)$ closed ball inside $\text{red}^{-1}(\overline{\alpha})$ centered in α . After an application of rigid GAGA to the coherent sheaves, the acyclicity of wide open neighborhoods implies that the hypercohomology of the complex $\mathcal{H}_k^\bullet$ can be computed simply by

$$H_{\text{dR}}^1(V_\lambda, \mathcal{H}_k^{\text{rig}}) \cong \Gamma(V_\lambda, \mathcal{H}_k^{\text{rig}} \otimes \Omega_{X^{\text{rig}}}^1(\log C)) / \nabla_k(\Gamma(V_\lambda, \mathcal{H}_k^{\text{rig}}))$$

Borrowing once again the notation from the modular curve case, we will denote the sections of ω^{rig} over V_λ by

$$M_{k, \lambda}^{\dagger, S} = \Gamma(V_\lambda, (\omega^{\text{rig}})^k)$$

and they should be thought as *λ -overconvergent modular forms*. We will denote it simply as $M_{k, \lambda}^{\dagger}$ when the set of poles is clear from the context.

We can apply the acyclicity property of V_λ to the short exact sequence of Lemma 1.

$$(5) \quad M_{k+2, \lambda}^{\dagger} / \theta^{k+1} (M_{-k, \lambda}^{\dagger}) \cong \Gamma(V_\lambda, \mathcal{H}_k^{\text{rig}} \otimes \Omega_{X^{\text{rig}}}^1(\log C)) / \nabla_k(\Gamma(V_\lambda, \mathcal{H}_k^{\text{rig}})) \cong H_{\text{dR}}^1(X \setminus S, \mathcal{H}_k).$$

Since every vector bundle with connection on a smooth formal model X of X^{rig} is overconvergent, the work of Baldassarri-Chiarellotto [BC91] establishes a comparison isomorphism

$$H_{\text{dR}}^1((X_{\mathbb{Q}_p}, S_{\mathbb{Q}_p}), \mathcal{H}_k) \cong H_{\text{dR}}^1(V_\lambda, \mathcal{H}_k^{\text{rig}}).$$

The left-hand side in the previous isomorphism is the de Rham cohomology with logarithmic coefficients. In characteristic 0, by an analytic result of Deligne [Del70] and later extended to an algebraic version in [AKR07] we can relate it to the open curve case

$$H_{\text{dR}}^1(X_{\mathbb{Q}_p} \setminus S_{\mathbb{Q}_p}, \mathcal{H}_k) \cong H_{\text{dR}}^1(V_\lambda, \mathcal{H}_k^{\text{rig}}).$$

In particular, we have the following commutative diagram where all arrows are isomorphisms

$$(6) \quad \begin{array}{ccc} M_{k+2}^{\text{mero}, S} / \theta^{k+1} (M_{-k, \lambda}^{\text{mero}, S}) & \longrightarrow & H_{\text{dR}}^1(X_{\mathbb{Q}_p} \setminus S_{\mathbb{Q}_p}, \mathcal{H}_k) \\ \downarrow & & \downarrow \\ M_{k+2, \lambda}^{\dagger} / \theta^{k+1} (M_{-k, \lambda}^{\dagger}) & \longrightarrow & H_{\text{dR}}^1(V_\lambda, \mathcal{H}_k^{\text{rig}}). \end{array}$$

In order to study and relate the Frobenius action on global sections and on the fibers we will use the structure of *overconvergent log F -isocrystal* of $(\mathcal{H}_k, \nabla)$. The relative de Rham cohomology in the modular curve case has this property thanks to the classical work of Katz [Kat73] while the Shimura curve case has been studied by Kassaei in [Kas04].

We then apply the Frobenius equivariant residue sequence following Coleman [Col94] and Coleman-Iovita [CI10].

$$(7) \quad 0 \rightarrow H_{\mathrm{dR}}^1(X^{\mathrm{rig}}, \mathcal{H}_k^{\mathrm{rig}}) \rightarrow H_{\mathrm{dR}}^1(V_\lambda, \mathcal{H}_k^{\mathrm{rig}}) \xrightarrow{\mathrm{Res}} \bigoplus_{\alpha \in S_K} H_{\mathrm{dR}}^0(A_{\alpha, \lambda}, \mathcal{H}_k)[1] \rightarrow 0$$

Where $A_{\alpha, \lambda}$ are rigid subspaces of V_λ conformal to the open annuli $\mathrm{res}^{-1}(\bar{\alpha}) \setminus B(\alpha, \lambda)$ and $[1]$ refers to the shift in the Frobenius action. The previous exact sequence gives in fact a rigid analogue of the Gysin sequence. Since every residue disk admits a basis of horizontal section for an overconvergent isocrystal as proved by Katz in [Kat71], we can rewrite the exact sequence as

$$(8) \quad 0 \rightarrow H_{\mathrm{dR}}^1(X^{\mathrm{rig}}, \mathcal{H}_k^{\mathrm{rig}}) \rightarrow H_{\mathrm{dR}}^1(V_\lambda, \mathcal{H}_k^{\mathrm{rig}}) \xrightarrow{\mathrm{Res}} \bigoplus_{\alpha \in S_{\mathbb{Q}_p}} \mathrm{Sym}^k(\alpha^* \mathcal{H})[1] \rightarrow 0.$$

In particular we have that the cohomology groups have the following dimensions

- $\dim_{\mathbb{Q}_p} H_{\mathrm{dR}}^1(X^{\mathrm{rig}}, \mathcal{H}_k^{\mathrm{rig}}) = \dim_{\mathbb{Q}_p} M_{k+2}(X_{\mathbb{Q}_p}) + \dim_{\mathbb{Q}_p} S_{k+2}(X_{\mathbb{Q}_p})$ by Lemma 1,
- $\dim_{\mathbb{Q}_p} \mathrm{Sym}^k(\alpha^* \mathcal{H}) = k + 1$ for every $\alpha \in S_{\mathbb{Q}_p}$,
- $\dim_{\mathbb{Q}_p} H_{\mathrm{dR}}^1(V_\lambda, \mathcal{H}_k^{\mathrm{rig}}) = \dim_{\mathbb{Q}_p} H_{\mathrm{dR}}^1(X^{\mathrm{rig}}, \mathcal{H}_k^{\mathrm{rig}}) + |S_{\mathbb{Q}_p}|(k + 1).$

In the elliptic setting we have

$$\mathrm{Sym}^k(\alpha^* \mathcal{H}) \cong \mathrm{Sym}^k(H_{\mathrm{dR}}^1(\mathcal{E}_\alpha))$$

for $\mathcal{E}_\alpha$ elliptic curve corresponding to the point α . In the quaternionic setting we have

$$\mathrm{Sym}^k(\alpha^* \mathcal{H}) \cong \mathrm{Sym}^k(e_1 \cdot H_{\mathrm{dR}}^1(\mathcal{A}_\alpha))$$

for $\mathcal{A}_\alpha$ abelian surface corresponding to the point α and e_1 a fixed choice of idempotent in $M_2(\mathbb{Z}_p)$.

Remark 1 (Parallel transport of eigenbasis). — *The purpose of this remark is to explain how Frobenius eigenbases behave when the poles are moved inside the same residue disk. The rigid Gysin sequence (7) does not depend on the choice of poles as long as they are lifts of a set of point modulo p . Consider $S = \{\alpha_1, \dots, \alpha_n\}$ a set of smooth points on the modular curve X . Consider $S' = \{\alpha'_1, \dots, \alpha'_n\}$ a second set for which α_i and α'_i lie in the same residue disk for $i = 1, \dots, n$. Recall that the 0-th hypercohomology group is given by*

$$H_{\mathrm{dR}}^0(A_{\alpha, \lambda}, \mathcal{H}_k) \cong \Gamma(A_{\alpha, \lambda}, \mathcal{H}_k^{\nabla=0}).$$

Then we have the following commutative diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & H_{\mathrm{dR}}^1(X_{\mathbb{Q}_p}, \mathcal{H}_k) & \longrightarrow & H_{\mathrm{dR}}^1(X_{\mathbb{Q}_p} \setminus S_{\mathbb{Q}_p}, \mathcal{H}_k) & \longrightarrow & \bigoplus_{\alpha \in S} H_{\mathrm{dR}}^1(\mathcal{E}_\alpha)[1] \longrightarrow 0 \\ & & \uparrow & & \uparrow & & \uparrow \\ 0 & \longrightarrow & H_{\mathrm{dR}}^1(X^{\mathrm{rig}}, \mathcal{H}_k^{\mathrm{rig}}) & \longrightarrow & H_{\mathrm{dR}}^1(V_\lambda, \mathcal{H}_k^{\mathrm{rig}})[1] & \longrightarrow & \bigoplus_{\alpha \in S} \Gamma(A_{\alpha, \lambda}, \mathcal{H}_k^{\nabla=0})[1] \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & H_{\mathrm{dR}}^1(X_{\mathbb{Q}_p}, \mathcal{H}_k) & \longrightarrow & H_{\mathrm{dR}}^1(X_{\mathbb{Q}_p} \setminus S'_{\mathbb{Q}_p}, \mathcal{H}_k) & \longrightarrow & \bigoplus_{\alpha' \in S'} H_{\mathrm{dR}}^1(\mathcal{E}_{\alpha'})[1] \longrightarrow 0 \end{array}$$

where the vertical maps are isomorphisms. Fixing a basis for one set of poles S , we can move the basis to S' using parallel transport computing an horizontal basis

$$\chi(\alpha, \alpha') : H_{\mathrm{dR}}^1(\mathcal{E}_\alpha) \rightarrow H_{\mathrm{dR}}^1(\mathcal{E}_{\alpha'}).$$

For example, we could try to compute the eigenbasis on CM points and then move the basis to any other point in the same residue disk.

4. Integral Frobenius structure

We now consider the log-crystalline cohomology group $H_{\log-\text{crys}}^1((\overline{X}, \overline{S})/\mathbb{Z}_p, \mathcal{H}_k)$. Since $\mathcal{H}_k$ has the structure of a log F -crystal as studied by Ogus [Ogu94], we obtain a $\mathbb{Z}_p$ module with a Frobenius action. By the classical work of Kato [Kat89] we have a comparison theorem with de Rham cohomology

$$H_{\log-\text{crys}}^1((\overline{X}, \overline{S})/\mathbb{Z}_p, \mathcal{H}_k) \otimes \mathbb{Q}_p \cong H_{\text{dR}}^1((X_{\mathbb{Q}_p}, S_{\mathbb{Q}_p}), \mathcal{H}_k)$$

and a comparison with rigid cohomology

$$H_{\log-\text{crys}}^1((\overline{X}, \overline{S})/\mathbb{Z}_p, \mathcal{H}_k) \otimes \mathbb{Q}_p \cong H_{\text{dR}}^1(V_\lambda, \mathcal{H}_k^{\text{rig}})$$

compatible with the action of Frobenius. We then find a Frobenius stable $\mathbb{Z}_p$ -lattice Λ in the rigid cohomology

$$\Lambda := \text{Im}(H_{\text{crys}}^1((\overline{X}, \overline{C} \cup \overline{S})/\mathbb{Z}_p, \mathcal{H}_k) \rightarrow H_{\text{crys}}^1((\overline{X}, \overline{C} \cup \overline{S})/\mathbb{Z}_p, \mathcal{H}_k) \otimes_{\mathbb{Z}_p} \mathbb{Q}_p \cong H_{\text{dR}}^1(V_\lambda, \mathcal{H}_k^{\text{rig}})).$$

By equation (5) we can see Λ contained in

$$\Lambda \hookrightarrow M_{k+2, \lambda}^\dagger / \theta^{k+1} (M_{-k, \lambda}^\dagger).$$

4.1. Explicit Frobenius lift. — Consider now the classical case of S consisting of the set of supersingular points and $\lambda < 1/p + 1$. As explained by [Kat73] and [Kas04], there exist a section of the projection map $X(p) \rightarrow X$ induced by the *canonical subgroup*. This morphism gives a lift of Frobenius V on the modular curve X given by

$$(9) \quad V : V_\lambda \rightarrow V_{\lambda^p}$$

acting at the level of the moduli space sending the elliptic curve or abelian surface to the corresponding quotient by the canonical subgroup. In the case of the elliptic modular curve, the V operator acts on q -expansion by

$$(Vf)(q^{1/N}) = p^k f(q^{p/N}).$$

The morphism V is flat and we can then define a U_p operator as the trace of Frobenius acting on the de Rham cohomology $H_{\text{dR}}^1(V_{\lambda^p}, \mathcal{H}_k)$. Coleman proved the following theorem in the case of the elliptic modular curve.

Theorem 1 ([Col96] Theorem 5.4). — *The space $H_{\text{dR}}^1(V_\lambda, \mathcal{H}_k^{\text{rig}})$ is finite dimensional and the operators V and U_p acting on the modular curve by correspondence induce endomorphisms Frob and Ver of $H_{\text{dR}}^1(V_\lambda, \mathcal{H}_k^{\text{rig}})$ such that*

$$\text{Frob} \circ \text{Ver} = \text{Ver} \circ \text{Frob} = p^{k+1}.$$

As further explained, the action of Verschiebung on the cohomology group $H_{\text{dR}}^1(V_{\lambda^p}, \mathcal{H}_k^{\text{rig}})$ is given by

$$\text{Ver} : [f] \rightarrow [U_p f]$$

for $f \in M_{k+2, \lambda}^\dagger$ under the isomorphism (5)

$$M_{k+2, \lambda}^\dagger / \theta^{k+1} (M_{-k, \lambda}^\dagger) \cong H_{\text{dR}}^1(V_\lambda, \mathcal{H}_k^{\text{rig}}).$$

This relation extends formally from the results of Kassaei for the Shimura curve over $\mathbb{Q}$. A complete proof of this statement will appear in forthcoming work by the author.

Enlarging the set S and removing disks from the ordinary locus, we can restrict the action of Frobenius and U_p to a smaller domain. The key easy lemma for Frobenius neighborhoods is the following

Lemma 2. — *Let B the residue disk of an ordinary point α defined over $\mathbb{F}_p$, let t be a Serre-Tate parameter. Then $B \cong \text{Spf}(\mathbb{Z}_p[[t]])_\eta$ and the Frobenius lift V restricted to B is a strict contraction.*

Fix α to be an ordinary $\mathbb{Z}_p$ point on X and $B \cong \text{Spf}(\mathbb{Z}_p[[t]])_\eta$ its residue disk, where we have fixed a parametrization. Consider V_λ the rigid variety obtained as before by removing small disks in the supersingular locus and V the Frobenius lift obtained from the canonical subgroup. Then by the previous lemma,

$$V|_B : B \rightarrow B$$

acts as a strict contraction. By the previous Lemma we can then find a small λ' for which V acts on the annulus contained in B as

$$V : A_{\alpha, \lambda} \rightarrow A_{\alpha, \lambda'^p}.$$

This means that we can remove from V_λ a small ball in B and several annuli around on the supersingular locus to obtain a wide open $V_{\lambda'}$ where the Frobenius lift V acts as in (9).

Combining the results of this section we obtain that there exists $f_1, \dots, f_r \in M_{k+2, \lambda}^\dagger$ λ -overconvergent modular forms with $r = \dim_{\mathbb{Q}_p} H_{\text{dR}}^1(V_{\lambda^p}, \mathcal{H}_k^{\text{rig}})$ such that

$$\Lambda = \langle [f_i] : i = 1, \dots, r \rangle_{\mathbb{Z}_p}$$

on which the V operator acts as an endomorphism and for which we have the relation $V \circ U_p = U_p \circ V = p^{k+1}$. Under the chain of isomorphisms (6) we can pick $f_1, \dots, f_r$ to be meromorphic modular forms over $\mathbb{Q}_p$ with poles along S .

Theorem A. — *Let X be a modular curve (elliptic or quaternionic) of level N defined over $\mathbb{Z}_p$, with $N > 3$, $p > 3$ and $(N, p) = 1$. Fix α be a $\mathbb{Z}_p$ point on the modular curve. Let k be a positive integer with $p > k + 1$, let $P(X), Q(X) \in \mathbb{Q}_p[X]$ be respectively the characteristic polynomial for the Frobenius action on $\text{Sym}^k(\alpha^* \mathcal{H})[1]$ and on $H_{\text{dR}}^1(X^{\text{rig}}, \mathcal{H}_k^{\text{rig}})$*

$$P(X) = \det(\text{Frob} - X | \text{Sym}^k(\alpha^* \mathcal{H})[1]) = \sum_{i=0}^{k+1} c_i X^i,$$

$$Q(X) = \det(\text{Frob} - X | H_{\text{dR}}^1(X^{\text{rig}}, \mathcal{H}_k)) = \sum_{i=0}^d d_i X^i,$$

$$P(X)Q(X) = \sum_{i=0}^{d+k+1} e_i X^i$$

with $d = \dim_{\mathbb{Q}_p} M_{k+2} + \dim_{\mathbb{Q}_p} S_{k+2}$. Let $M = d + k + 1$, then for every $f \in M_{k+2}^{\text{mero}, \{\alpha\}}$ we have

$$\sum_{i=0}^M e_{M-i} p^{M-i} U_p^i(f) \in \theta^{k+1} \left(M_{-k}^{\text{mero}, \{\alpha\}} \right)$$

If we assume $P(X), Q(X)$ coprime, then there exist $f_1, \dots, f_{k+1} \in M_{k+2}^{\text{mero}, \{\alpha\}}$ such that $[f_i] \in \Lambda$ for $i = 1, \dots, k+1$ and

$$\sum_{i=0}^{k+1} c_i p^{k+1-i} U_p^i(f) \in \theta^{k+1} \left(M_{-k}^{\text{mero}, \{\alpha\}} \right)$$

for every $f \in \langle f_1, \dots, f_{k+1} \rangle_{\mathbb{Z}_p}$.

Proof. — Consider the Gysin exact sequence (8)

$$0 \rightarrow H_{\text{dR}}^1(X^{\text{rig}}, \mathcal{H}_k^{\text{rig}}) \rightarrow H_{\text{dR}}^1(V_\lambda, \mathcal{H}_k^{\text{rig}}) \xrightarrow{\text{Res}} \text{Sym}^k(\alpha^* \mathcal{H})[1] \rightarrow 0.$$

The assumption that the characteristic polynomials of Frobenius are coprime imply that the extension group of $\mathbb{Q}_p$ -vector spaces with Frobenius action is zero and then the sequence split. In general, the product of the characteristic polynomials of Frobenius annihilates $H_{\text{dR}}^1(V_\lambda, \mathcal{H}_k^{\text{rig}})$. For every class $[f] \in H_{\text{dR}}^1(V_\lambda, \mathcal{H}_k^{\text{rig}})$ with $f \in M_{k+2}^{\text{mero}, \{\alpha\}}$ representative obtained under the chain of isomorphisms (6) we then have

$$(10) \quad \sum_{i=0}^M e_i \text{Frob}^i([f]) = 0.$$

We can now use the trick of enlarging the set of removed points to allow an explicit lift of Frobenius. Consider $S = \{\alpha\} \cup SS_p$ where SS_p is the set of supersingular points on X . Consider the following

commutative diagram obtained by functoriality (8)

$$\begin{array}{ccccccc}
0 & \longrightarrow & H_{\mathrm{dR}}^1(X^{\mathrm{rig}}, \mathcal{H}_k^{\mathrm{rig}}) & \longrightarrow & M_{k+2, \lambda}^{\dagger, S} / \theta^{k+1} \left(M_{-k, \lambda}^{\dagger, S} \right) & \longrightarrow & \bigoplus_{\alpha \in SS_{p, \mathbb{Q}_p}} \mathrm{Sym}^k H_{\mathrm{dR}}^1(\mathcal{E}_\alpha) \oplus \mathrm{Sym}^k H_{\mathrm{dR}}^1(\alpha^* \mathcal{E})[1] \longrightarrow 0 \\
& & \uparrow & & \uparrow & & \uparrow \\
0 & \longrightarrow & H_{\mathrm{dR}}^1(X^{\mathrm{rig}}, \mathcal{H}_k^{\mathrm{rig}}) & \longrightarrow & M_{k+2, \lambda}^{\dagger, \{\alpha\}} / \theta^{k+1} \left(M_{-k, \lambda}^{\dagger, \{\alpha\}} \right) & \longrightarrow & \mathrm{Sym}^k H_{\mathrm{dR}}^1(\alpha^* \mathcal{E})[1] \longrightarrow 0
\end{array}$$

By Theorem 1 we have that Frobenius and Verschiebung act on $M_{k+2, \lambda}^{\dagger, S} / \theta^{k+1} \left(M_{-k, \lambda}^{\dagger, S} \right)$ by V and U_p respectively. Composing (10) with Ver^{d+k+1} we obtain

$$(11) \quad \sum_{i=0}^M e_i(\mathrm{Ver}^M \circ \mathrm{Frob}^i)[f] = 0,$$

$$(12) \quad \sum_{j=0}^M e_{M-j} p^{M-j} U_p^j(f) \in \theta^{k+1} \left(M_{-k}^{\mathrm{mero}, \{\alpha\}} \right).$$

In the case where the exact sequence splits, we consider a basis of elements coming from Λ and we obtain the previous relation with the characteristic polynomial $P(X) = \det(\mathrm{Frob} - X | \mathrm{Sym}^k(\alpha^* \mathcal{H})[1])$. $\square$

We conclude the section with two remarks. The first on a possible improvement using Hecke operators, the latter on known results about the slope of p -newforms.

Remark 2 (Hecke operators). — *We could strengthen the previous theorem using the theory of Hecke operators. The geometric correspondences inducing Hecke operators act on the cohomology groups. By taking a linear combination $T = \sum \lambda_n T_n$, we can define an operator that annihilates the first classical term in the Gysin sequence (8). In this way, the Frobenius submodule obtained from the cohomology group $H_{\mathrm{dR}}^1(V_\lambda, \mathcal{H}_k^{\mathrm{rig}})$ under the action of T becomes isomorphic to the piece coming from the de Rham cohomology of the fibers.*

Remark 3 (Slope of p -newforms). — *Consider once again the case of S consisting of all the supersingular points of X . Then [Col96] in the case of elliptic modular curves identified the parabolic part of $H_{\mathrm{dR}}^1(V_{p/p+1}, \mathcal{H}_k)$ with classical modular forms of level $\Gamma_1(N) \cap \Gamma_0(p)$ and weight $k+2$. This has been later generalized in [Kas09] including the case of quaternionic modular forms. Theorem 1 allows us to relate the slopes of eigenvalues of the Frobenius with the slopes of eigenvalues of U_p operator. In the elliptic case we obtain that the Frobenius module*

$$\bigoplus_{\alpha \in S_{\mathbb{Q}_p}} \mathrm{Sym}^k(\alpha^* \mathcal{H}) \cong \bigoplus_{\alpha \in S_{\mathbb{Q}_p}} \mathrm{Sym}^k H_{\mathrm{dR}}^1(\mathcal{E}_\alpha)$$

is isoclinic of slope $\frac{k}{2}$ where $\mathcal{E}_\alpha$ is the elliptic curve corresponding to the point α . From the exact sequence (8) we deduce that the newspace of the classical modular forms of level $\Gamma_1(N) \cap \Gamma_0(p)$ is isoclinic of slope $\frac{k+2}{2}$ due to the 1-shift and then slope $k/2$ due to the relation $V \circ U_p = p^{k+1}$.

5. Arithmetic applications

5.1. Congruences on q -expansion. — In this section, we restrict to the case where $X = X_1(N)$ is an elliptic modular curve and we consider the expansion at the cusp $[\infty]$. We summarize here the classical geometric description of the q -expansion, referring to [Kat73] for further details. Consider the infinitesimal neighborhood $\mathrm{Spf}(\mathbb{Z}_p[[q^{1/N}]])$ of the cusp $[\infty]$ of the formal model $\mathfrak{X}$. The fiber product with the universal elliptic curve gives rise to the Tate curve

$$\begin{array}{ccc}
\mathrm{Tate}(q^{1/N}) & \longrightarrow & \mathcal{E} \\
\downarrow & & \downarrow \\
\mathrm{Spf}(\mathbb{Z}_p[[q^{1/N}]]) & \xrightarrow{\iota} & \mathfrak{X}.
\end{array}$$

The global sections of the line bundle are given by

$$\Gamma(\mathrm{Spf}(\mathbb{Z}_p[[q^{1/N}]]), \iota^* \omega) = \mathbb{Z}_p[[q^{1/N}]] \omega_{\mathrm{can}}$$

where ω_{can} is the canonical differential dt/t on the Tate curve $\mathbb{G}_m/q^{\mathbb{Z}}$. The global sections of the sheaf of differential 1-forms on X with logarithmic poles at the cusps are given by

$$\Gamma(\mathrm{Spf}(\mathbb{Z}_p[[q^{1/N}]]), \iota^* \Omega_{\mathcal{X}}^1(\log C)) = \mathbb{Z}_p[[q^{1/N}]] \frac{dq}{q}.$$

The Kodaira-Spencer isomorphism (3) gives an identification

$$KS(\omega_{\mathrm{can}}^2) = \frac{dq}{q}$$

Given a modular form $f \in \Gamma(U, \omega^k \otimes \Omega_X^1(\log C))$ with $U \subset X$ we can then define its q -expansion as

$$\iota^* f = f(q^{1/N}) \omega_{\mathrm{can}}^k \otimes \frac{dq}{q}, \quad f(q^{1/N}) = \sum_{n=0}^{\infty} a_n(f) q^{\frac{n}{N}}.$$

The pull back of the relative de Rham cohomology at infinity is given by

$$\Gamma(\mathrm{Spf}(\mathbb{Z}_p[[q^{1/N}]]), \iota^* \mathcal{H}) = \mathbb{Z}_p[[q^{1/N}]] \omega_{\mathrm{can}} \oplus \mathbb{Z}_p[[q^{1/N}]] \eta_{\mathrm{can}}$$

where

$$\nabla(\omega_{\mathrm{can}}) =: \eta_{\mathrm{can}} \otimes \frac{dq}{q} \in \iota^*(\mathcal{H} \otimes \Omega_X^1), \quad \nabla(\eta_{\mathrm{can}}) = 0.$$

In general, we will have

$$\Gamma(\mathrm{Spf}(\mathbb{Z}_p[[q^{1/N}]]), \iota^* \mathcal{H}_k) = \bigoplus_{a+b=k} \mathbb{Z}_p[[q^{1/N}]] \omega_{\mathrm{can}}^a \eta_{\mathrm{can}}^b$$

By the acyclicity of $\mathrm{Spf}(\mathbb{Z}_p[[q^{1/N}]])$, we obtain

$$H_{\mathrm{dR}}^1(\mathrm{Spf}(\mathbb{Z}_p[[q^{1/N}]]), \iota^* \mathcal{H}_k) = \Gamma(\mathrm{Spf}(\mathbb{Z}_p[[q^{1/N}]]), \iota^*(\mathcal{H}_k \otimes_{\mathcal{X}} \Omega_X^1)) / \nabla(\Gamma(\mathrm{Spf}(\mathbb{Z}_p[[q^{1/N}]]), \iota^*(\mathcal{H}_k))).$$

Using the exact sequence in Lemma 1 we then obtain

$$\begin{aligned} H_{\mathrm{dR}}^1(\mathrm{Spf}(\mathbb{Z}_p[[q^{1/N}]]), \iota^* \mathcal{H}_k) &\cong \Gamma(\mathrm{Spf}(\mathbb{Z}_p[[q^{1/N}]]), \iota^* \omega^{k+2} / \theta^{k+1} (\Gamma(\mathrm{Spf}(\mathbb{Z}_p[[q^{1/N}]]), \iota^* \omega^{-k}))) \\ &\cong \mathbb{Z}_p[[q^{1/N}]] / \theta^{k+1} (\mathbb{Z}_p[[q^{1/N}]]). \end{aligned}$$

Combining the constructions, we can then form a q -expansion map on the cohomology classes

Lemma 3. — *The morphism $\iota : \mathrm{Spf}(\mathbb{Z}_p[[q^{1/N}]])) \rightarrow \mathfrak{X}$ between formal schemes induces the following morphism*

$$\begin{aligned} \Lambda &\rightarrow \mathbb{Z}_p[[q^{1/N}]] / \theta^{k+1} (\mathbb{Z}_p[[q^{1/N}]])) \\ [f] &\mapsto [f(q)]. \end{aligned}$$

Proof. — By an integral comparison with the de Rham cohomology [Kat89] (Theorem 6.4) we can compute the crystalline cohomology by

$$H_{\mathrm{crys}}^1(\overline{X}, \overline{C} \cup \overline{S}) / \mathbb{Z}_p, \mathcal{H}_k) \cong H_{\mathrm{dR}}^1((\mathfrak{X}, S), \mathcal{H}_k)$$

The map is then induced by the functoriality along the morphism ι

$$H_{\mathrm{dR}}^1((\mathfrak{X}, S), \mathcal{H}_k) \rightarrow H_{\mathrm{dR}}^1(\mathbb{Z}_p[[q^{1/N}]]), \iota^* \mathcal{H}_k).$$

By the previous remark, we have

$$H_{\mathrm{dR}}^1(\mathbb{Z}_p[[q^{1/N}]]), \iota^* \mathcal{H}_k) \cong \mathbb{Z}_p[[q^{1/N}]] / \theta^{k+1} (\mathbb{Z}_p[[q^{1/N}]]).$$

□

Corollary 1. — Let X be an elliptic modular curve of level N defined over $\mathbb{Z}_p$, with $N > 3$, $p > 3$ and $(N, p) = 1$. Fix a $\mathbb{Z}_p$ -point α on the modular curve. Let k be a positive integer with $p > k + 1$, let $P(X), Q(X) \in \mathbb{Q}_p[X]$ be respectively the characteristic polynomials of the Frobenius action on $\mathrm{Sym}^k H_{\mathrm{dR}}^1(\alpha^* \mathcal{E})[1]$, with $\mathcal{E} \rightarrow X$ the universal elliptic curve, and on $H_{\mathrm{dR}}^1(X^{\mathrm{rig}}, \mathcal{H}_k^{\mathrm{rig}})$

$$P(X) = \det(\mathrm{Frob} - X | \mathrm{Sym}^k(\alpha^* \mathcal{H})[1]) = \sum_{i=0}^{k+1} c_i X^i,$$

$$Q(X) = \det(\mathrm{Frob} - X | H_{\mathrm{dR}}^1(X^{\mathrm{rig}}, \mathcal{H}_k)) = \sum_{i=0}^d d_i X^i,$$

$$P(X)Q(X) = \sum_{i=0}^{d+k+1} e_i X^i$$

with $d = \dim_{\mathbb{Q}_p} M_{k+2} + \dim_{\mathbb{Q}_p} S_{k+2}$. Let $M = d + k + 1$, then for every $f \in \Lambda$ we have

$$\sum_{i=0}^M e_{M-i} p^{M-i} a_{np^{l+i}}(f) \equiv 0 \pmod{p^{l(k+1)}}$$

If we assume that $P(X), Q(X)$ are coprime, then there exist $f_1, \dots, f_{k+1} \in M_{k+2}^{\mathrm{mero}, \{\alpha\}}$ such that $[f_i] \in \Lambda$ for $i = 1, \dots, k+1$ and

$$\sum_{i=0}^{k+1} c_{k+1-i} p^{k+1-i} a_{np^{l+i}}(f) \equiv 0 \pmod{p^{l(k+1)}}$$

for every $f \in \langle f_1, \dots, f_{k+1} \rangle_{\mathbb{Z}_p}$.

Proof. — By Theorem A we have the following relation

$$\sum_{i=0}^M e_{M-i} p^{M-i} U_p^i(f) \in \theta^{k+1} \left(M_{-k}^{\mathrm{mero}, \{\alpha\}} \right).$$

The action of U_p , which is the trace of the Frobenius lift V on the q -expansion of weight $k+2$ modular forms, is classically given by

$$f(q^{1/N}) = \sum_{n \geq 0} a_n(f) q^{n/N} \mapsto (U_p f)(q^{1/N}) = \sum_{n \geq 0} a_{pn}(f) q^{n/N}.$$

In particular, using the previous Lemma, we obtain

$$\sum_{i=0}^M e_{M-i} p^{M-i} U_p^i(f(q^{1/N})) \in \theta^{k+1} \left(\mathbb{Z}_p[[q^{1/N}]] \right)$$

from which we deduce for $l > 1$

$$\sum_{i=0}^M e_{M-i} p^{M-i} a_{np^{l+i}}(f) \equiv 0 \pmod{p^{l(k+1)}}.$$

□

5.2. An explicit example - Zhang's congruences. — In [Zha25], Zhang provided numerous numerical observation about meromorphic modular forms and their relation with the Frobenius eigenvalues of the poles. In this short section we want to show how the previous constructions and results provide a framework to understand his congruences.

We will work directly with the coarse moduli space $X(1) \cong \mathbb{P}_j^1$. The descent of the results from level N to level 1 can be carried over by classical techniques. We consider Example 4.6 of *loc. cit.*. Fix $k = 2$ and $p > k + 1 = 3$, $C/\mathbb{Q}$ an elliptic curve given by the equation

$$y^2 + xy = x^3 - x^2 - 2x - 1.$$

We have C is a CM elliptic curve with respect to the order $\mathbb{Z}[\frac{1+\sqrt{-7}}{2}]$ of discriminant -7 and $j(C) = -3375$. Consider $S = \{j(C)\} \cup SS_p$ where SS_p consists in a set of lift of supersingular points on $X(1)$. The Gysin sequence (8) gives us the following commutative diagram

$$\begin{array}{ccccccc}
0 & \longrightarrow & H_{\mathrm{dR}}^1(X^{\mathrm{rig}}, \mathcal{H}_2^{\mathrm{rig}}) & \longrightarrow & M_{4,p^{1/p+1}}^{\dagger, SS_p} / \theta^3(M_{-2,p^{1/p+1}}^{\dagger, SS_p}) & \longrightarrow & \bigoplus_{\alpha \in SS_{p, \mathbb{Q}_p}} \mathrm{Sym}^2 H_{\mathrm{dR}}^1(\mathcal{E}_\alpha)[1] \longrightarrow 0 \\
& & \downarrow = & & \downarrow & & \downarrow \\
0 & \longrightarrow & H_{\mathrm{dR}}^1(X^{\mathrm{rig}}, \mathcal{H}_2^{\mathrm{rig}}) & \longrightarrow & M_{4,\lambda}^{\dagger, S} / \theta^3(M_{-k,\lambda}^{\dagger, S}) & \longrightarrow & \bigoplus_{\alpha \in SS_{p, \mathbb{Q}_p}} \mathrm{Sym}^2 H_{\mathrm{dR}}^1(\mathcal{E}_\alpha) \oplus \mathrm{Sym}^2 H_{\mathrm{dR}}^1(C)[1] \longrightarrow 0 \\
& & \uparrow = & & \uparrow & & \uparrow \\
0 & \longrightarrow & H_{\mathrm{dR}}^1(X^{\mathrm{rig}}, \mathcal{H}_2^{\mathrm{rig}}) & \longrightarrow & M_{4,p^{1/p+1}}^{\dagger, \{j(C)\}} / \theta^3(M_{-2,p^{1/p+1}}^{\dagger, \{j(C)\}}) & \longrightarrow & \mathrm{Sym}^2 H_{\mathrm{dR}}^1(C)[1] \longrightarrow 0
\end{array}$$

The radius λ can be chosen accordingly to Lemma 2 to have an explicit lift of Frobenius corresponding with the quotient of the canonical subgroup. We have an isomorphism of $\mathbb{Q}_p$ -vector spaces

$$\bigoplus_{\alpha \in SS_{p, \mathbb{Q}_p}} \mathrm{Sym}^2 H_{\mathrm{dR}}^1(\mathcal{E}_\alpha) \cong S_4(\Gamma_0(p))^{p-\mathrm{new}}$$

this identification is at the core of the proof of classicality of Coleman.

The bottom row can be identified with the equivalent Gysin exact sequence in de Rham cohomology, in particular we have the isomorphism

$$M_4^{\mathrm{mero}, \{j(C)\}} / \theta^3 \left(M_{-2,\lambda}^{\mathrm{mero}, \{j(C)\}} \right) \cong M_{4,p^{1/p+1}}^{\dagger, \{j(C)\}} / \theta^{k+1} \left(M_{-2,p^{1/p+1}}^{\dagger, \{j(C)\}} \right)$$

The three dimensional space given by $\mathrm{Sym}^2 H_{\mathrm{dR}}^1(C)$ can be represented in the cohomology group by meromorphic modular forms with poles along $j(C) = -3375$. Zhang identifies the following basis

$$\begin{aligned}
f_1 &= \frac{E_4}{j + 3375}, \\
f_2 &= 19 \cdot \frac{E_4}{j + 3375} - 91125 \cdot \frac{E_4}{(j + 3375)^2}, \\
f_3 &= 1399 \cdot \frac{E_4}{j + 3375} - 19008675 \cdot \frac{E_4}{(j + 3375)^2} + 54251268750 \cdot \frac{E_4}{(j + 3375)^3}.
\end{aligned}$$

Then for every ordinary prime $p > 3$ we have

$$\begin{aligned}
a_{np^{l+1}}(f_1) &\equiv u_p(C)^2 a_{np^l}(f_1) \pmod{p^{3l}}, \\
a_{np^{l+1}}(f_2) &\equiv p a_{np^l}(f_1) \pmod{p^{3l}}, \\
a_{np^{l+1}}(f_3) &\equiv p^2 u_p(C)^{-2} a_{np^l}(f_1) \pmod{p^{3l}}
\end{aligned}$$

where $u_p(C)$ is the p -adic root of the polynomial $X^2 - a_p(C)X + p$. According to Corollary 1, the meromorphic modular forms f_1, f_2, f_3 are the representatives of the classes in $H_{\mathrm{dR}}^1(V_\lambda, \mathcal{H}_k^{\mathrm{rig}})$ corresponding to the Frobenius eigenspace isomorphic to $\mathrm{Sym}^2 H_{\mathrm{dR}}^1(C)$.

References

- [AKR07] Timothy G. Abbott, Kiran S. Kedlaya, and David Roe. Bounding picard numbers of surfaces using p -adic cohomology. *Experimental Mathematics*, 16(3):313–326, 2007. Also appeared as “Bounding Picard numbers of surfaces using p -adic cohomology” (2006) arXiv preprint.
- [ALS25] Michael Allen, Ling Long, and Hasan Saad. Atkin and swinnerton-dyer congruences for meromorphic modular forms, 2025.
- [BC91] F. Baldassarri and B. Chiarellotto. Algebraic versus rigid cohomology with logarithmic coefficients. In *Barsotti Symposium in Algebraic Geometry*, volume 15 of *Perspectives in Mathematics*, pages 163–193. Academic Press, 1991.
- [CI10] Robert F. Coleman and Adrian Iovita. Hidden structures on semistable curves. In *Représentations p -adiques de groupes p -adiques III : méthodes globales et géométriques*, number 331 in *Astérisque*, pages 179–254. Société mathématique de France, 2010. MR2667889.
- [Col94] Robert F. Coleman. A p -adic shimura isomorphism and p -adic periods of modular forms. In *p -adic monodromy and the Birch and Swinnerton-Dyer conjecture*, volume 165 of *Contemporary Mathematics*, pages 21–51. American Mathematical Society, 1994. Exposé, originally from 1991.
- [Col96] Robert F. Coleman. Classical and overconvergent modular forms. *Inventiones Mathematicae*, 124(1-3):215–241, 1996.

- [Del70] Pierre Deligne. *Équations différentielles à points singuliers réguliers*, volume 163 of *Lecture Notes in Mathematics*. Springer, Berlin, 1970.
- [Kas04] Payman L. Kassaei. P-adic modular forms over shimura curves over totally real fields. *Compositio Mathematica*, 140(2):359–395, 2004.
- [Kas09] Payman L. Kassaei. Overconvergence and classicality: the case of curves. *Journal für die reine und angewandte Mathematik*, 631:109–139, 2009.
- [Kat71] Nicholas M. Katz. Travaux de dwork. In *Séminaire Bourbaki, 23e année (1970/71), Exp. No. 409*, volume 244 of *Lecture Notes in Mathematics*, pages 167–200. Springer, 1971.
- [Kat73] Nicholas M. Katz. p -adic properties of modular schemes and modular forms. In A. Antoine and F. Oort, editors, *Modular Functions of One Variable III*, volume 350 of *Lecture Notes in Mathematics*, pages 69–190. Springer, 1973.
- [Kat89] Kazuya Kato. Logarithmic structures of fontaine–illusie. In *Algebraic Analysis, Geometry, and Number Theory (Baltimore, MD, 1988)*, pages 191–224. Johns Hopkins University Press, 1989. Also available as Kato’s 1988 manuscript, received November 8 1988; see the PDF at math.brown.edu.
- [KS13] Matija Kazalicki and Anthony J. Scholl. Modular forms, de rham cohomology and congruences. *Transactions of the American Mathematical Society*, 365(9):4729–4762, 2013.
- [Ogu94] Arthur Ogus. *F-crystals, Griffiths transversality, and the Hodge decomposition*. Number 221 in *Astérisque*. Société mathématique de France, 1994.
- [Sch85] Anthony J. Scholl. Modular forms and de rham cohomology; atkin–swinnerton-dyer congruences. *Inventiones Mathematicae*, 79(1):49–77, 1985.
- [Zha25] Pengcheng Zhang. Elliptic curves and fourier coefficients of meromorphic modular forms, 2025.

December 2025

PAOLO BORDIGNON • *E-mail* : p.bordignon@math.leidenuniv.nl, Mathematical Institute, Gorlaeus Building, Einsteinweg 55, 2333 CC, Leiden (Netherlands)