
ON A COHOMOLOGICAL DESCRIPTION OF CONGRUENCES BETWEEN MODULAR FORMS AND ELLIPTIC CURVES

by

Paolo Bordinon

Contents

1. Introduction.....	1
2. Geometric setting.....	2
3. Rigid cohomological description.....	3
4. Integral Frobenius structure.....	5
5. Arithmetic applications.....	6
References.....	9

1. Introduction

The aim of this paper is to give a cohomological interpretation of certain congruences between Fourier coefficients of modular forms and Frobenius eigenvalues arising from the geometry of modular (and Shimura) curves. More precisely, we study congruences for overconvergent and meromorphic modular forms through the de Rham, rigid, and crystalline cohomology of natural coefficient systems attached to the universal abelian variety.

Let $p > 3$ be a prime and $N > 3$ an integer coprime to p . We work either with the modular curve $X_1(N)$ or, more generally, with a Shimura curve of quaternionic type, both defined over $\mathbb{Z}_p$. Denote by ω the line bundle of invariant differentials and by $(\mathcal{H}, \nabla)$ the rank-two vector bundle given by relative de Rham cohomology of the universal object, equipped with its Gauss–Manin connection. For an integer $k \geq 0$, we consider the symmetric power $\mathcal{H}_k = \mathrm{Sym}^k \mathcal{H}$, which carries a natural filtration and a logarithmic connection satisfying Griffiths transversality.

A key point is that the de Rham cohomology of $(\mathcal{H}_k, \nabla)$ on suitable open curves admits an explicit description in terms of modular forms modulo iterates of the θ -operator. This provides a bridge between cohomology classes and q -expansions, allowing arithmetic information to be extracted from Frobenius actions on cohomology. In particular, by comparing de Rham, rigid, and logarithmic crystalline realizations, one obtains canonical Frobenius-stable lattices whose images under the q -expansion map yield congruences for Fourier coefficients.

The guiding philosophy is that congruences for modular forms should be understood as shadows of Frobenius structures on geometric cohomology groups. From this perspective, classical results of Coleman on overconvergent modular forms and the slope theory of the U_p -operator arise naturally from the geometry of the supersingular locus, while new congruences for meromorphic modular forms can be interpreted via the Frobenius action on the de Rham cohomology of the poles.

2. Geometric setting

Let $p > 3$ be a prime, let $N > 3$ be an integer coprime with p . Consider one of the following setting

- $X = X_1(N)$ modular curve of level $\Gamma_1(N)$ defined over $\mathbb{Z}_p$ classifying elliptic curves with level structure,
- $X = X_B$ a Shimura curve of level $V_1(N)$ defined over $\mathbb{Z}_p$ classifying false elliptic curves with level structure.

We will denote $\bar{X} = X \times k$ its special fiber, $C \subset X(\mathbb{Z}_p)$ the finite collection (possibly empty) of cusps. Let ω be the ample line bundle over X obtained from the pushforward of the differential bundle on the universal object. Let $(\mathcal{H}, \nabla)$ be the rank 2 vector bundle arising from the relative de Rham cohomology over X with logarithmic connection

$$\nabla : \mathcal{H} \rightarrow \mathcal{H} \otimes_{\mathbb{Z}_p} \Omega_X^1(\log C)$$

coming from the Gauss-Manin connection. In the quaternionic case we are fixing a choice of a non-trivial idempotent element of $M_2(\mathbb{Z}_p)$ and projecting the four dimensional vector bundle coming from the relative de Rham cohomology to obtain $\mathcal{H}_k$ and analogously with ω . See [Kas04]. Denote by $\mathcal{H}_k$ its k -th symmetric tensor power $\mathcal{H}_k = \text{Sym}^k \mathcal{H}$.

We recall the standard properties of $\mathcal{H}_k$. The vector bundles ω and $\mathcal{H}$ fit in a short exact sequence

$$(1) \quad 0 \rightarrow \omega \rightarrow \mathcal{H} \rightarrow \omega^{-1} \rightarrow 0$$

inducing a $(k+1)$ -step decreasing filtration on H_k

$$(2) \quad \mathcal{H}_k = \text{Fil}^0 \mathcal{H}_k \supseteq \text{Fil}^1 \mathcal{H}_k \supseteq \cdots \supseteq \text{Fil}^k \mathcal{H}_k \cong \omega^k \supseteq \text{Fil}^{k+1} \mathcal{H}_k = 0.$$

The graded pieces of the filtration are isomorphic to

$$\text{Gr}_i \mathcal{H}_k = F^{k-i} \mathcal{H}_k / F^{k+1-i} \mathcal{H}_k \cong \omega^{k-2i}.$$

The connection obeys *Griffith's transversality*

$$\nabla \text{Fil}^{k-i} \mathcal{H}_k \subseteq \text{Fil}^{k-i-1} \mathcal{H}_k \otimes_{\mathcal{O}_X} \Omega_X^1(\log C)$$

and that these maps induce a *Kodaira-Spencer* type of isomorphism of $\mathcal{O}_X$ -modules for $p > k+1$

$$(3) \quad \nabla : \text{Gr}_i \mathcal{H}_k \cong \text{Gr}_{i+1} \mathcal{H}_k \otimes_X \Omega_X^1(\log C).$$

Following the usual notation, we write

$$M_k(X) = \Gamma(X, \omega^k), \quad S_k(X) = \Gamma(X, \omega^{k-2} \otimes \Omega_X^1).$$

We denote the complex arising from the connection by

$$\mathcal{H}_k^\bullet := [0 \rightarrow \mathcal{H}_k \rightarrow \mathcal{H}_k \otimes_{\mathcal{O}_X} \Omega_X^1]$$

and denote its hypercohomology groups by H_{dR}^i

$$H_{\text{dR}}^i(X, \mathcal{H}_k) := \mathbb{H}^i(X, \mathcal{H}_k^\bullet).$$

Lemma 1. — *We have a short exact sequence*

$$(4) \quad 0 \rightarrow \omega_{-k} \xrightarrow{\theta^{k+1}} \omega_{k+2} \rightarrow \mathcal{H}_k \otimes_X \Omega_X^1(\log C) / \nabla_k(\mathcal{H}_k) \rightarrow 0$$

where the right map is compatible with inclusions and an Eichler-Shimura short exact sequence of $\mathcal{O}_K$ -modules

$$0 \rightarrow M_{k+2}(X) \rightarrow H_{\text{dR}}^1(X, \mathcal{H}_k) \rightarrow S_{k+2}(X)^\vee \rightarrow 0.$$

Proof. — Consider the filtration $(\mathrm{Fil}^p \mathcal{H}_k^\bullet)_p$ of the complex $\mathcal{H}_k^\bullet$

$$\mathrm{Fil}^p \mathcal{H}_k^\bullet := [0 \rightarrow \mathrm{Fil}^p \mathcal{H}_k \rightarrow \mathrm{Fil}^{p-1} \mathcal{H}_k \otimes_{\mathcal{O}_X} \Omega_X^1(\log C)] .$$

and take the associated spectral sequence $E_0^{p,q} = \mathrm{Fil}^p \mathcal{H}_k^{p+q} / \mathrm{Fil}^{p+1} \mathcal{H}_k^{p+q}$. The only terms that survive in page 1 are

$$E_1^{0,0} \cong \mathcal{H}_k / \mathrm{Fil}^1 \mathcal{H}_k, \quad E_1^{k+1,-k} \cong \mathrm{Fil}^k \mathcal{H}_k \otimes_{\mathcal{X}} \Omega_{\mathcal{X}}^1(\log C).$$

The spectral sequence degenerates at page $k+2$ converging to $E_{k+2}^{k+1,-k} = H^1(\mathcal{H}_k^\bullet)$. This gives rise to the short exact sequence

$$0 \rightarrow \mathcal{H}_k / \mathrm{Fil}^1 \mathcal{H}_k \xrightarrow{\theta^{k+1}} \mathrm{Fil}^k \mathcal{H}_k \otimes_{\mathcal{X}} \Omega_{\mathcal{X}}^1(\log C) \rightarrow \mathcal{H}_k \otimes_{\mathcal{X}} \Omega_{\mathcal{X}}^1(\log C) / \nabla_k(\mathcal{H}_k) \rightarrow 0.$$

The second exact sequence is obtained applying the spectral sequence to the hypercohomology of the graded pieces obtaining

$$\begin{aligned} 0 \rightarrow H_{\mathrm{dR}}^0(X, \mathcal{H}_k^\bullet) \rightarrow H^0(X, \omega^{-k}) \rightarrow H^0(X, \omega^k \otimes \Omega_X^1(\log C)) \rightarrow H_{\mathrm{dR}}^1(X, \mathcal{H}_k^\bullet) \rightarrow \\ \rightarrow H^1(X, \omega^{-k}) \rightarrow H^1(X, \omega^k \otimes \Omega_X^1(\log C)) \rightarrow \dots \end{aligned}$$

giving the desired exact sequence using ampleness of ω and Serre duality. $\square$

Fix an effective divisor $S = \sum_{i=1}^n \alpha_i$ supported on the open curve $X \setminus C$ and defined over $\mathbb{Z}_p$,

$$\alpha_i : \mathrm{Spec}(\mathcal{O}_K) \rightarrow X \setminus C$$

for K a finite field extension over $\mathbb{Q}_p$ with distinct close points on the special fiber $\overline{X}$. We denote by $\overline{S}$ their reduction to the special fiber $\overline{X}$ and $S_{\mathbb{Q}_p}$ their restriction to the generic fiber. We have that the curve $U = X_{\mathbb{Q}_p} \setminus S_{\mathbb{Q}_p}$ is affine.

As a direct consequence of the lemma, the de Rham cohomology of $\mathcal{H}_k$ can be computed by

$$H_{\mathrm{dR}}^1(U, C), \mathcal{H}_k) = M_{k+2}^{\mathrm{mero}, S} / \theta^{k+1} (M_{-k, \lambda}^{\mathrm{mero}, S})$$

where we have used the notation $M_{k+2}^{\mathrm{mero}, S} = \Gamma(U, \omega^{k+2})$ and should be thought as *meromorphic modular forms with poles along S*.

3. Rigid cohomological description

Consider X^{rig} to be the rigid analytification of X and the reduction map

$$\mathrm{red} : X^{\mathrm{rig}} \rightarrow \overline{X}(\overline{\mathbb{F}}_p)$$

where we fixed $\overline{k}$ algebraic closure of k . We then consider the wide open obtained from X^{rig} removing closed balls in the residue disks of the points S . Since X is smooth, $\mathrm{red}^{-1}(\overline{\alpha})$ is conformal to an open ball B for every $\overline{\alpha} \in \overline{S}$. Fixing this parametrization and fixing $0 < \lambda < 1$, define

$$V_\lambda = X^{\mathrm{rig}} \setminus \bigcup_{\alpha \in S} B(\alpha, \lambda)$$

with $B(\alpha, \lambda)$ closed ball inside $\mathrm{red}^{-1}(\overline{\alpha})$ centered in α . After an application of rigid GAGA to the coherent sheaves, the aciclycity of wide open neighborhoods implies that the hypercohomology of the complex $\mathcal{H}_k^\bullet$ can be computed simply by

$$H_{\mathrm{dR}}^1(V_\lambda, \mathcal{H}_k^{\mathrm{rig}}) \cong \Gamma(V_\lambda, \mathcal{H}_k^{\mathrm{rig}} \otimes \Omega_{X^{\mathrm{rig}}}^1(\log C)) / \nabla_k(\Gamma(V_\lambda, \mathcal{H}_k^{\mathrm{rig}}))$$

Borrowing once again the notation from the modular curve case, we will denote the sections of ω^{rig} over V_λ by

$$M_{k, \lambda}^{\dagger, S} = \Gamma(V_\lambda, (\omega^{\mathrm{rig}})^k)$$

and they should be think as λ -overconvergent modular forms. We will denote it simply as $M_{k, \lambda}^{\dagger}$ when the set of poles is clear from the context.

We can apply the acyclicity property of V_λ to the short exact sequence of Lemma 1.

$$(5) \quad M_{k+2,\lambda}^\dagger / \theta^{k+1}(M_{-k,\lambda}^\dagger) \cong \Gamma(V_\lambda, \mathcal{H}_k^{\text{rig}} \otimes \Omega_{X^{\text{rig}}}^1(\log C)) / \nabla_k(\Gamma(V_\lambda, \mathcal{H}_k^{\text{rig}})) \cong H_{\text{dR}}^1(X \setminus S, \mathcal{H}_k).$$

Since every vector bundle with connection on a smooth formal model X of X^{rig} is overconvergent, the work of Baldassarri-Chiarellotto [BC91] establishes a comparison isomorphism

$$H_{\text{dR}}^1((X_{\mathbb{Q}_p}, S_{\mathbb{Q}_p}), \mathcal{H}_k) \cong H_{\text{dR}}^1(V_\lambda, \mathcal{H}_k^{\text{rig}}).$$

The left-hand side in the previous isomorphism is the de Rham cohomology with logarithmic coefficients. In characteristic 0, by an analytic result of Deligne [Del70] and later extended to an algebraic version in [AKR07] we can relate it to the open curve case

$$H_{\text{dR}}^1(X_{\mathbb{Q}_p} \setminus S_{\mathbb{Q}_p}, \mathcal{H}_k) \cong H_{\text{dR}}^1(V_\lambda, \mathcal{H}_k^{\text{rig}}).$$

In particular, we have the following commutative diagram where all arrows are isomorphisms

$$\begin{array}{ccc} M_{k+2}^{\text{mero},S} / \theta^{k+1}(M_{-k,\lambda}^{\text{mero},S}) & \longrightarrow & H_{\text{dR}}^1(X_{\mathbb{Q}_p} \setminus S_{\mathbb{Q}_p}, \mathcal{H}_k) \\ \downarrow & & \downarrow \\ M_{k+2,\lambda}^\dagger / \theta^{k+1}(M_{-k,\lambda}^\dagger) & \longrightarrow & H_{\text{dR}}^1(V_\lambda, \mathcal{H}_k^{\text{rig}}). \end{array}$$

In order to study and relate the Frobenius action on global sections and on the fibers we will use the structure of *overconvergent log F -isocrystal* of $(\mathcal{H}_k, \nabla)$. The relative de Rham cohomology in the modular curve case has this property thanks to the classical work of Katz [Kat73] while the Shimura curve case has been studied by Kassaei in [Kas04].

We then apply the Frobenius equivariant residue sequence following Coleman [Col94] and Coleman-Iovita [CI10].

$$(6) \quad 0 \rightarrow H_{\text{dR}}^1(X^{\text{rig}}, \mathcal{H}_k^{\text{rig}}) \rightarrow H_{\text{dR}}^1(V_\lambda, \mathcal{H}_k^{\text{rig}}) \xrightarrow{\text{Res}} \bigoplus_{\alpha \in S_K} H_{\text{dR}}^0(A_{\alpha,\lambda}, \mathcal{H}_k)[1] \rightarrow 0$$

Where $A_{\alpha,\lambda}$ are rigid subspaces of V_λ conformal to the open annuli $\text{res}^{-1}(\bar{\alpha}) \setminus B(\alpha, \lambda)$ and [1] refers to the shift in the Frobenius action. The previous exact sequence gives in fact a rigid analogue of the Gysin sequence. Since every residue disk admits a basis of horizontal section for an overconvergent isocrystal as proved by Katz in [Kat71], we can rewrite the exact sequence as

$$(7) \quad 0 \rightarrow H_{\text{dR}}^1(X^{\text{rig}}, \mathcal{H}_k^{\text{rig}}) \rightarrow H_{\text{dR}}^1(V_\lambda, \mathcal{H}_k^{\text{rig}}) \xrightarrow{\text{Res}} \bigoplus_{\alpha \in S_{\mathbb{Q}_p}} \text{Sym}^k(\alpha^* \mathcal{H})[1] \rightarrow 0.$$

In particular we have that the cohomology groups have the following dimensions

- $\dim_{\mathbb{Q}_p} H_{\text{dR}}^1(X^{\text{rig}}, \mathcal{H}_k^{\text{rig}}) = \dim_{\mathbb{Q}_p} M_{k+2}(X_{\mathbb{Q}_p}) + \dim_{\mathbb{Q}_p} S_{k+2}(X_{\mathbb{Q}_p})$ by Lemma 1,
- $\dim_{\mathbb{Q}_p} \text{Sym}^k(\alpha^* \mathcal{H}) = k + 1$ for every $\alpha \in S_{\mathbb{Q}_p}$,
- $\dim_{\mathbb{Q}_p} H_{\text{dR}}^1(V_\lambda, \mathcal{H}_k^{\text{rig}}) = \dim_{\mathbb{Q}_p} H_{\text{dR}}^1(X^{\text{rig}}, \mathcal{H}_k^{\text{rig}}) + |S_{\mathbb{Q}_p}|(k + 1)$.

In the elliptic setting we have

$$\text{Sym}^k(\alpha^* \mathcal{H}) \cong \text{Sym}^k(H_{\text{dR}}^1(\mathcal{E}_\alpha))$$

for $\mathcal{E}_\alpha$ elliptic curve corresponding to the point α . In the quaternionic setting we have

$$\text{Sym}^k(\alpha^* \mathcal{H}) \cong \text{Sym}^k(e_1 \cdot H_{\text{dR}}^1(\mathcal{A}_\alpha))$$

for $\mathcal{A}_\alpha$ abelian surface corresponding to the point α and e_1 a fixed choice of idempotent in $M_2(\mathbb{Z}_p)$.

4. Integral Frobenius structure

We now consider the log-crystalline cohomology group $H_{\log-\text{crys}}^1((\overline{X}, \overline{S})/\mathbb{Z}_p, \mathcal{H}_k)$. Since $\mathcal{H}_k$ has the structure of a log F -crystal as studied by Ogus [Ogu94], we obtain a $\mathbb{Z}_p$ module with a Frobenius action. By the classical work of Kato [Kat89] we have a comparison theorem with de Rham cohomology

$$H_{\log-\text{crys}}^1((\overline{X}, \overline{S})/\mathbb{Z}_p, \mathcal{H}_k) \otimes \mathbb{Q}_p \cong H_{\text{dR}}^1((X_{\mathbb{Q}_p}, S_{\mathbb{Q}_p}), \mathcal{H}_k)$$

and a comparison with rigid cohomology

$$H_{\log-\text{crys}}^1((\overline{X}, \overline{S})/\mathbb{Z}_p, \mathcal{H}_k) \otimes \mathbb{Q}_p \cong H_{\text{dR}}^1(V_\lambda, \mathcal{H}_k^{\text{rig}})$$

compatible with the action of Frobenius. We then find a Frobenius stable $\mathbb{Z}_p$ -lattice Λ in the rigid cohomology

$$\Lambda := \text{Im}(H_{\text{crys}}^1((\overline{X}, \overline{C} \cup \overline{S})/\mathbb{Z}_p, \mathcal{H}_k) \rightarrow H_{\text{crys}}^1((\overline{X}, \overline{C} \cup \overline{S})/\mathbb{Z}_p, \mathcal{H}_k) \otimes_{\mathbb{Z}_p} \mathbb{Q}_p \cong H_{\text{dR}}^1(V_\lambda, \mathcal{H}_k^{\text{rig}})).$$

By equation (5) we can see Λ contained in

$$\Lambda \hookrightarrow M_{k+2, \lambda}^\dagger / \theta^{k+1} (M_{-k, \lambda}^\dagger).$$

4.1. Explicit Frobenius lift. — Consider now the classical case of S consisting of the set supersingular points and $\lambda < 1/p + 1$. As explained by [Kat73] and [Kas04], there exist a section of the map $X(p) \rightarrow X$ that forgets the level p structure induced by the *canonical subgroup*. This morphism given a lift of Frobenius V on the modular curve X given by

$$(8) \quad V : V_\lambda \rightarrow V_{\lambda^p}$$

acting at the level of the moduli space sending the elliptic curve or abelian surface to the corresponding quotient by the canonical subgroup. In the case of the elliptic modular curve, the V operator acts on q -expansion by

$$(Vf)(q^{1/N}) = p^k f(q^{p/N}).$$

The morphism V is flat and we can then define a U_p operator as the trace of Frobenius acting on the de Rham cohomology $H_{\text{dR}}^1(V_{\lambda^p}, \mathcal{H}_k)$. Coleman proved the following theorem in the case of the elliptic modular curve.

Theorem 1 ([Col96] Theorem 5.4). — *The space $H_{\text{dR}}^1(V_\lambda, \mathcal{H}_k^{\text{rig}})$ is finite dimensional and the operators V and U_p acting on the modular curve by correspondence induce endomorphisms Frob and Ver of $H_{\text{dR}}^1(V_\lambda, \mathcal{H}_k^{\text{rig}})$ such that*

$$\text{Frob} \circ \text{Ver} = \text{Ver} \circ \text{Frob} = p^{k+1}.$$

As further explained, the action of Verschiebung on the cohomology group $H_{\text{dR}}^1(V_{\lambda^p}, \mathcal{H}_k^{\text{rig}})$ is given by

$$\text{Ver} : [f] \rightarrow [U_p f]$$

for $f \in M_{k+2, \lambda}^\dagger$ under the isomorphism (5)

$$M_{k+2, \lambda}^\dagger / \theta^{k+1} (M_{-k, \lambda}^\dagger) \cong H_{\text{dR}}^1(V_{\lambda^p}, \mathcal{H}_k^{\text{rig}}).$$

This relation extends formally from the results of Kassaei for the Shimura curve over $\mathbb{Q}$.

Enlarging the set S and removing disks from the ordinary locus, we can restrict the action of Frobenius and U_p to a smaller domain. The key easy lemma for Frobenius neighborhoods is the following

Lemma 2. — *Let B the residue disk of an ordinary point α defined over $\mathbb{F}_p$, let t be a Serre-Tate parameter. Then $B \cong \text{Spf}(\mathbb{Z}_p[[t]])_\eta$ and the Frobenius lift V restricted to B is a strict contraction.*

We can then find a small enough λ for which the Frobenius lift V acts as in (8). Combining the results of this section we obtain that there exists $f_1, \dots, f_r \in M_{k+2, \lambda}^\dagger$ λ -overconvergent modular forms with $r = \dim_{\mathbb{Q}_p} H_{\text{dR}}^1(V_{\lambda^p}, \mathcal{H}_k^{\text{rig}})$ such that

$$\Lambda = \langle [f_i] : i = 1, \dots, r \rangle_{\mathbb{Z}_p}$$

On which the V operator acts as an endomorphism and for which we have the relation $V \circ U_p = U_p \circ V = p^{k+1}$.

We conclude the section with a remark on known results about the slope of p -newforms.

Remark 1. — Consider once again the case of S consisting of all the supersingular points of X . Then [Col96] in the case of elliptic modular curves identified the parabolic part of $H_{\text{dR}}^1(V_{p/p+1}, \mathcal{H}_k)$ with classical modular forms of level $\Gamma_1(N) \cap \Gamma_0(p)$. This has been later generalized in [Kas09] including the case of quaternionic modular forms. Theorem 1 allows us to relate the slopes of eigenvalues of the Frobenius with the slopes of eigenvalues of U_p operator. In the elliptic case we obtain that the Frobenius eigenvalues are $\pm p^{\frac{k+1}{2}}$ on the space

$$\bigoplus_{\alpha \in S_{\mathbb{Q}_p}} \text{Sym}^k(\alpha^* \mathcal{H}) \cong \bigoplus_{\alpha \in S_{\mathbb{Q}_p}} \text{Sym}^k H_{\text{dR}}^1(\mathcal{E}_\alpha)$$

where $\mathcal{E}_\alpha$ is the elliptic curve corresponding to the point α . From the exact sequence (7) we deduce that the newspace of the classical modular forms of level $\Gamma_1(N) \cap \Gamma_0(p)$ have slope $\frac{k-1}{2}$ due to the 1-shift.

5. Arithmetic applications

5.1. Congruences on q -expansion. — In this section we restrict to the case $X = X_1(N)$ elliptic modular curve and we consider the expansion at the cusp $[\infty]$. We summarize here the classical geometric description of q -expansion, referring to [Kat73] for further details. Consider the infinitesimal neighborhood $\text{Spf}(\mathbb{Z}_p[[q]])$ of the cusp $[\infty]$ of the formal model $\mathfrak{X}$. The fiber product with the universal elliptic curve gives rise to the Tate curve

$$\begin{array}{ccc} \text{Tate}(q^{1/N}) & \longrightarrow & \mathcal{E} \\ \downarrow & & \downarrow \\ \text{Spf}(\mathbb{Z}_p[[q^{1/N}]]) & \xrightarrow{\iota} & \mathfrak{X}. \end{array}$$

The global sections of the line bundle are given by

$$\Gamma(\text{Spf}(\mathbb{Z}_p[[q^{1/N}]]), \iota^* \omega) = \mathbb{Z}_p[[q^{1/N}]] \omega_{\text{can}}$$

where ω_{can} is the canonical differential dt/t on the Tate curve $\mathbb{G}_m/q^{\mathbb{Z}}$. The global sections of the sheaf of differential 1-forms on X with logarithmic poles at the cusps are given by

$$\Gamma(\text{Spf}(\mathbb{Z}_p[[q^{1/N}]]), \iota^* \Omega_{\mathfrak{X}}^1(\log \mathcal{C})) = \mathbb{Z}_p[[q^{1/N}]] \frac{dq}{q}.$$

The Kodaira-Spencer isomorphism (3) gives an identification

$$KS(\omega_{\text{can}}^2) = \frac{dq}{q}$$

Given a modular form $f \in \Gamma(U, \omega^k \otimes \Omega_X^1(\log C))$ with $U \subset X$ we can then define its q -expansion as

$$\iota^* f = f(q^{1/N}) \omega_{\text{can}}^k \otimes \frac{dq}{q}, \quad f(q^{1/N}) = \sum_{n=0}^{\infty} a_n(f) q^{\frac{n}{N}}.$$

The pull back of the relative de Rham cohomology at infinity is given by

$$\Gamma(\mathrm{Spf}(\mathbb{Z}_p[[q^{1/N}]]), \iota^* \mathcal{H}) = \mathbb{Z}_p[[q^{1/N}]] \omega_{\mathrm{can}} \oplus \mathbb{Z}_p[[q^{1/N}]] \eta_{\mathrm{can}}$$

where

$$\nabla(\omega_{\mathrm{can}}) =: \eta_{\mathrm{can}} \otimes \frac{dq}{q} \in \iota^*(\mathcal{H} \otimes \Omega_X^1), \quad \nabla(\eta_{\mathrm{can}}) = 0.$$

In general we will have,

$$\Gamma(\mathrm{Spf}(\mathbb{Z}_p[[q^{1/N}]]), \iota^* \mathcal{H}_k) = \bigoplus_{a+b=k} \mathbb{Z}_p[[q^{1/N}]] \omega_{\mathrm{can}}^a \eta_{\mathrm{can}}^b$$

By the acyclicity of $\mathrm{Spf}(\mathbb{Z}_p[[q^{1/N}]])$, we have that

$$H_{\mathrm{dR}}^1(\mathrm{Spf}(\mathbb{Z}_p[[q^{1/N}]]), \iota^* \mathcal{H}_k) = \Gamma(\mathrm{Spf}(\mathbb{Z}_p[[q^{1/N}]]), \iota^*(\mathcal{H}_k \otimes_{\mathcal{X}} \Omega_{\mathcal{X}}^1)) / \nabla(\Gamma(\mathrm{Spf}(\mathbb{Z}_p[[q^{1/N}]]), \iota^*(\mathcal{H}_k))).$$

Using the exact sequence in Lemma 1 we then obtain

$$\begin{aligned} H_{\mathrm{dR}}^1(\mathrm{Spf}(\mathbb{Z}_p[[q^{1/N}]]), \iota^* \mathcal{H}_k) &\cong \Gamma(\mathrm{Spf}(\mathbb{Z}_p[[q^{1/N}]]), \iota^* \omega^{k+2} / \theta^{k+1} (\Gamma(\mathrm{Spf}(\mathbb{Z}_p[[q^{1/N}]]), \iota^* \omega^{-k}))) \\ &\cong \mathbb{Z}_p[[q^{1/N}]] / \theta^{k+1} (\mathbb{Z}_p[[q^{1/N}]]). \end{aligned}$$

Combining the constructions we can form a q -expansion map on the cohomology classes

Theorem 2. — *The morphism $\iota : \mathrm{Spf}(\mathbb{Z}_p[[q^{1/N}]])) \rightarrow \mathfrak{X}$ between formal schemes induces the following morphism*

$$\begin{aligned} \Lambda &\rightarrow \mathbb{Z}_p[[q^{1/N}]] / \theta^{k+1} (\mathbb{Z}_p[[q^{1/N}]])) \\ [f] &\mapsto [f(q)]. \end{aligned}$$

Proof. — By an integral comparison with the de Rham cohomology [Kat89] (Theorem 6.4) we can compute the crystalline cohomology by

$$H_{\mathrm{crys}}^1((\overline{X}, \overline{C} \cup \overline{S}) / \mathbb{Z}_p, \mathcal{H}_k) \cong H_{\mathrm{dR}}^1((\mathfrak{X}, S), \mathcal{H}_k)$$

The map is then induced by the functoriality along the morphism ι

$$H_{\mathrm{dR}}^1((\mathfrak{X}, S), \mathcal{H}_k) \rightarrow H_{\mathrm{dR}}^1(\mathbb{Z}_p[[q^{1/N}]]), \iota^* \mathcal{H}_k).$$

and by the previous remark we have

$$H_{\mathrm{dR}}^1(\mathbb{Z}_p[[q^{1/N}]]), \iota^* \mathcal{H}_k) \cong \mathbb{Z}_p[[q^{1/N}]] / \theta^{k+1} (\mathbb{Z}_p[[q^{1/N}]]).$$

□

Corollary 1. — *Consider $K/\mathbb{Q}_p$ a finite field extension, suppose $[f] \in \Lambda \otimes_{\mathbb{Z}_p} \mathcal{O}_k$ is an eigenvector for the action of Frobenius with eigenvalue μ , then for every $n, l > 1$*

$$a_{np^{l+1}}(f) \equiv \frac{p^{k+1}}{\mu} a_{np^l}(f) \pmod{p^{l(k+1)}}.$$

Proof. — The action of U_p , trace of the Frobenius lift V on the q -expansion of weight $k+2$ modular forms, is classically given by

$$f(q^{1/N}) = \sum_{n \geq 0} a_n(f) q^{n/N} \mapsto (U_p f)(q^{1/N}) = \sum_{n \geq 0} a_{pn}(f) q^{n/N}.$$

Taking $[f]$ to be an eigenclass in Λ corresponding to an eigenvalue μ for the action of Frobenius we then have $[f]$ is an eigenclass of the U_p operator with eigenvalue p^{k+1}/μ by Theorem 1

$$\left(\frac{p^{k+1}}{\mu} f - U_p f \right) (q^{1/N}) \in \theta^{k+1} \mathbb{Z}_p[[q^{1/N}]]$$

from which we deduce for $l > 1$

$$\frac{p^{k+1}}{\mu} a_{np^l}(f) - a_{np^{l+1}}(f) \equiv 0 \pmod{p^{l(k+1)}}.$$

□

5.2. An explicit example - P. Zhang congruences. — In [Zha25], Zhang provided numerous numerical observation about meromorphic modular forms and their relation with the Frobenius eigenvalues of the poles. In this short section we want to show how the previous constructions and results provide a framework to understand his congruences.

We will work directly with the coarse moduli space $X(1) \cong \mathbb{P}_j^1$. The descent of the results from level N to level 1 can be carried over by classical techniques. We consider Example 4.6 of *loc. cit.*. Fix $k = 2$ and $p > k + 1 = 3$, $C/\mathbb{Q}$ an elliptic curve given by the equation

$$y^2 + xy = x^3 - x^2 - 2x - 1.$$

We have C is a CM elliptic curve with respect to the order $\mathbb{Z}[\frac{1+\sqrt{-7}}{2}]$ of discriminant -7 and $j(C) = -3375$. Consider $S = \{j(C)\} \cup SS_p$ where SS_p consists in a set of lift of supersingular points on $X(1)$. The Gysin sequence (7) gives us the following commutative diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & H_{\text{dR}}^1(X^{\text{rig}}, \mathcal{H}_k^{\text{rig}}) & \longrightarrow & M_{k+2, p^{1/p+1}}^{\dagger, SS_p} / \theta^{k+1} (M_{-k, p^{1/p+1}}^{\dagger, SS_p}) & \longrightarrow & \bigoplus_{\alpha \in SS_p, \mathbb{Q}_p} \text{Sym}^4 H_{\text{dR}}^1(\mathcal{E}_\alpha)[1] \longrightarrow 0 \\ & & \downarrow = & & \downarrow & & \downarrow \\ 0 & \longrightarrow & H_{\text{dR}}^1(X^{\text{rig}}, \mathcal{H}_k^{\text{rig}}) & \longrightarrow & M_{k+2, \lambda}^{\dagger, S} / \theta^{k+1} (M_{-k, \lambda}^{\dagger, S}) & \longrightarrow & \bigoplus_{\alpha \in SS_p, \mathbb{Q}_p} \text{Sym}^4 H_{\text{dR}}^1(\mathcal{E}_\alpha) \oplus \text{Sym}^4 H_{\text{dR}}^1(C)[1] \longrightarrow 0 \\ & & \uparrow = & & \uparrow & & \uparrow \\ 0 & \longrightarrow & H_{\text{dR}}^1(X^{\text{rig}}, \mathcal{H}_k^{\text{rig}}) & \longrightarrow & M_{k+2, p^{1/p+1}}^{\dagger, \{j(C)\}} / \theta^{k+1} (M_{-k, p^{1/p+1}}^{\dagger, \{j(C)\}}) & \longrightarrow & \text{Sym}^4 H_{\text{dR}}^1(C)[1] \longrightarrow 0 \end{array}$$

The radius λ can be chosen accordingly to Lemma 2 to have an explicit lift of Frobenius corresponding with the quotient of the canonical subgroup. We have an isomorphism of $\mathbb{Q}_p$ -vector spaces

$$\bigoplus_{\alpha \in SS_p, \mathbb{Q}_p} \text{Sym}^4 H_{\text{dR}}^1(\mathcal{E}_\alpha) \cong S_4(\Gamma_0(p))^{p-\text{new}}$$

this identification is at the core of the proof of classicality of Coleman.

The bottom row can be identified with the equivalent Gysin exact sequence in de Rham cohomology, in particular we have the isomorphism

$$M_{k+2}^{\text{mero}, \{j(C)\}} / \theta^{k+1} \left(M_{-k, \lambda}^{\text{mero}, \{j(C)\}} \right) \cong M_{k+2, p^{1/p+1}}^{\dagger, \{j(C)\}} / \theta^{k+1} \left(M_{-k, p^{1/p+1}}^{\dagger, \{j(C)\}} \right)$$

The three dimensional space given by $\text{Sym}^4 H_{\text{dR}}^1(C)$ can be represented in the cohomology group by meromorphic modular forms with poles along $j(C) = -3375$. Zhang identifies the following basis

$$\begin{aligned} f_1 &= \frac{E_4}{j + 3375}, \\ f_2 &= 19 \cdot \frac{E_4}{j + 3375} - 91125 \cdot \frac{E_4}{(j + 3375)^2}, \\ f_3 &= 1399 \cdot \frac{E_4}{j + 3375} - 19008675 \cdot \frac{E_4}{(j + 3375)^2} + 54251268750 \cdot \frac{E_4}{(j + 3375)^3}. \end{aligned}$$

Then for every ordinary prime $p > 3$ we have

$$\begin{aligned} a_{np^{l+1}}(f_1) &\equiv u_p(C)^2 a_{np^l}(f_1) \pmod{p^{3l}}, \\ a_{np^{l+1}}(f_2) &\equiv p a_{np^l}(f_1) \pmod{p^{3l}}, \\ a_{np^{l+1}}(f_3) &\equiv p^2 u_p(C)^{-2} a_{np^l}(f_1) \pmod{p^{3l}}. \end{aligned}$$

References

- [AKR07] Timothy G. Abbott, Kiran S. Kedlaya, and David Roe. Bounding picard numbers of surfaces using p -adic cohomology. *Experimental Mathematics*, 16(3):313–326, 2007. Also appeared as “Bounding Picard numbers of surfaces using p -adic cohomology” (2006) arXiv preprint.
- [BC91] F. Baldassarri and B. Chiarellotto. Algebraic versus rigid cohomology with logarithmic coefficients. In *Barsotti Symposium in Algebraic Geometry*, volume 15 of *Perspectives in Mathematics*, pages 163–193. Academic Press, 1991.
- [CI10] Robert F. Coleman and Adrian Iovita. Hidden structures on semistable curves. In *Représentations p -adiques de groupes p -adiques III : méthodes globales et géométriques*, number 331 in *Astérisque*, pages 179–254. Société mathématique de France, 2010. MR2667889.
- [Col94] Robert F. Coleman. A p -adic shimura isomorphism and p -adic periods of modular forms. In *p -adic monodromy and the Birch and Swinnerton-Dyer conjecture*, volume 165 of *Contemporary Mathematics*, pages 21–51. American Mathematical Society, 1994. Exposé, originally from 1991.
- [Col96] Robert F. Coleman. Classical and overconvergent modular forms. *Inventiones Mathematicae*, 124(1-3):215–241, 1996.
- [Del70] Pierre Deligne. *Équations différentielles à points singuliers réguliers*, volume 163 of *Lecture Notes in Mathematics*. Springer, Berlin, 1970.
- [Kas04] Payman L. Kassaei. P -adic modular forms over shimura curves over totally real fields. *Compositio Mathematica*, 140(2):359–395, 2004.
- [Kas09] Payman L. Kassaei. Overconvergence and classicality: the case of curves. *Journal für die reine und angewandte Mathematik*, 631:109–139, 2009.
- [Kat71] Nicholas M. Katz. Travaux de dwork. In *Séminaire Bourbaki, 23e année (1970/71), Exp. No. 409*, volume 244 of *Lecture Notes in Mathematics*, pages 167–200. Springer, 1971.
- [Kat73] Nicholas M. Katz. p -adic properties of modular schemes and modular forms. In A. Antoinne and F. Oort, editors, *Modular Functions of One Variable III*, volume 350 of *Lecture Notes in Mathematics*, pages 69–190. Springer, 1973.
- [Kat89] Kazuya Kato. Logarithmic structures of fontaine–illusie. In *Algebraic Analysis, Geometry, and Number Theory (Baltimore, MD, 1988)*, pages 191–224. Johns Hopkins University Press, 1989. Also available as Kato’s 1988 manuscript, received November 8 1988; see the PDF at math.brown.edu.
- [Ogu94] Arthur Ogus. *F-crystals, Griffiths transversality, and the Hodge decomposition*. Number 221 in *Astérisque*. Société mathématique de France, 1994.
- [Zha25] Pengcheng Zhang. Elliptic curves and fourier coefficients of meromorphic modular forms, 2025.