

Profunctorial Semantics I

Fosco Loregian 

March 21, 2020

Algebraic structures

A **group** is a set equipped with operations

- $m : G \times G \rightarrow G$
- $i : G \rightarrow G$
- $e : 1 \rightarrow G$

...

you know the drill

Algebraic structures

Theorem (Higman-Neumann 1953)

A **group** is a set equipped with a single binary operation $/ : G \times G \rightarrow G$ subject to the single equation

$$x / (((x/x)/y)/z) / (((x/x)/x)/z) = y$$

for every $x, y, z \in X$.

Algebraic structures

Theorem (Higman-Neumann 1953)

A **group** is a set equipped with a single binary operation $/ : G \times G \rightarrow G$ subject to the single equation

$$x / (((x/x)/y)/z) / (((x/x)/x)/z) = y$$

for every $x, y, z \in X$.

Well.

This is awkward.

The theory of equationally definable classes of algebras, initiated by Birkhoff in the early thirties, is [...] hampered in its usefulness by two defects. [...] the second is the awkwardness inherent in the presentation of an equationally definable class in terms of operations and equations.

The theory of equationally definable classes of algebras, initiated by Birkhoff in the early thirties, is [...] hampered in its usefulness by two defects. [...] the second is the awkwardness inherent in the presentation of an equationally definable class in terms of operations and equations.

Quite recently, Lawvere, by introducing the notion - closely akin to the clones P. Hall - of an algebraic theory, rectified the second defect.

Definition

An **operator domain** is a sequence $\underline{\Omega} = (\Omega_n \mid n \in \mathbb{N})$; the elements of Ω_n are called **operations** of **arity** n .

Definition

An **interpretation** \underline{E} of an operator domain $\underline{\Omega}$ consists of a pair $(E, (f_\omega \mid \omega \in \Omega_n, n \in \mathbb{N}))$ where $f_\omega : E^n \rightarrow E$ is an n -ary operation on the set E called the *carrier* of \underline{E} .


Definition

An **operator domain** is a sequence $\underline{\Omega} = (\Omega_n \mid n \in \mathbb{N})$; the elements of Ω_n are called **operations** of **arity** n .

Definition

An **interpretation** \underline{E} of an operator domain $\underline{\Omega}$ consists of a pair $(E, (f_\omega \mid \omega \in \Omega_n, n \in \mathbb{N}))$ where $f_\omega : E^n \rightarrow E$ is an n -ary operation on the set E called the *carrier* of \underline{E} .

An operator domain can be represented as a (rooted) graph: for example, for groups



Way better to use functors.

A **Lawvere theory** is an identity-on-objects functor $p : \mathbf{Fin}^{\mathbf{o}} \rightarrow \mathcal{L}$ that commutes with finite products.

Unwinding the definition:

- \mathcal{L} is a category with the same objects as \mathbf{Fin} , the category of finite sets and functions;

A **Lawvere theory** is an identity-on-objects functor $p : \mathbf{Fin}^{\circ} \rightarrow \mathcal{L}$ that commutes with finite products.

Unwinding the definition:

- \mathcal{L} is a category with the same objects as \mathbf{Fin} , the category of finite sets and functions;
- p is a functor that acts trivially on objects

A **Lawvere theory** is an identity-on-objects functor $p : \mathbf{Fin}^{\mathbf{o}} \rightarrow \mathcal{L}$ that commutes with finite products.

Unwinding the definition:

- \mathcal{L} is a category with the same objects as \mathbf{Fin} , the category of finite sets and functions;
- p is a functor that acts trivially on objects
- The only thing that can change between \mathbf{Fin} and \mathcal{L} is the number of morphisms $[n] \rightarrow [m]$.

A **Lawvere theory** is an identity-on-objects functor $p : \mathbf{Fin}^{\mathbf{o}} \rightarrow \mathcal{L}$ that commutes with finite products.

Unwinding the definition:

- \mathcal{L} is a category with the same objects as \mathbf{Fin} , the category of finite sets and functions;
- p is a functor that acts trivially on objects
- The only thing that can change between \mathbf{Fin} and \mathcal{L} is the number of morphisms $[n] \rightarrow [m]$.

Equivalently: p is a **promonad** on the opposite of \mathbf{Fin} , regarded as an object of the bicategory of profunctors, that preserves the monoidal structure. \mathcal{L} is the Kleisli object of p .

$$\left\{ \begin{array}{l} \text{identity on obj} \\ \text{left adjoints} \\ p: [\mathcal{L}, \mathbf{Set}] \rightarrow [\mathbf{Fin}^{\mathbf{o}}, \mathbf{Set}] \end{array} \right\} \rightleftharpoons \left\{ \begin{array}{l} \text{monads in Prof} \\ p: \mathbf{Fin}^{\mathbf{o}} \rightsquigarrow \mathbf{Fin}^{\mathbf{o}} \end{array} \right\}$$

- The trivial theory is the identity functor $1_{\text{Fin}} : \text{Fin}^{\circ} \rightarrow \text{Fin}^{\circ}$

- The trivial theory is the identity functor $1_{\mathbf{Fin}} : \mathbf{Fin}^{\circ} \rightarrow \mathbf{Fin}^{\circ}$
- Since p preserves products, it is **uniquely determined by its value on $[1]$** . This means that if $p : \mathbf{Fin}^{\circ} \rightarrow \mathcal{L}$ is a Lawvere theory, then every object of \mathcal{L} is X^n if $p[1] = X$.

- The trivial theory is the identity functor $1_{\mathbf{Fin}} : \mathbf{Fin}^{\mathbf{o}} \rightarrow \mathbf{Fin}^{\mathbf{o}}$
- Since p preserves products, it is **uniquely determined by its value on $[1]$** . This means that if $p : \mathbf{Fin}^{\mathbf{o}} \rightarrow \mathcal{L}$ is a Lawvere theory, then every object of \mathcal{L} is X^n if $p[1] = X$.
- The only difference between \mathbf{Fin} and \mathcal{L} is thus the set of morphisms $[n] \rightarrow [m]$.

- The trivial theory is the identity functor $1_{\mathbf{Fin}} : \mathbf{Fin}^{\circ} \rightarrow \mathbf{Fin}^{\circ}$
- Since p preserves products, it is **uniquely determined by its value on $[1]$** . This means that if $p : \mathbf{Fin}^{\circ} \rightarrow \mathcal{L}$ is a Lawvere theory, then every object of \mathcal{L} is X^n if $p[1] = X$.
- The only difference between \mathbf{Fin} and \mathcal{L} is thus the set of morphisms $[n] \rightarrow [m]$.

The theory of groups is generated by

$$\mathcal{L}_{\text{Grp}} = \begin{array}{ccccc} & & i & & \\ & & \circlearrowleft & & \\ [0] & \xrightarrow{e} & [1] & \xleftarrow{m} & [2] \end{array}$$

and their compositions/products.

A **model** for a Lawvere theory p is a product-preserving functor $\ell : \mathcal{L} \rightarrow \mathbf{Set}$.

A **model** for a Lawvere theory p is a product-preserving functor $\ell : \mathcal{L} \rightarrow \mathbf{Set}$.

The category $\mathbf{Mod}(p)$ for a Lawvere theory is a full, **reflective** subcategory of the category $[\mathcal{L}, \mathbf{Set}]$ of all functors $\mathcal{L} \rightarrow \mathbf{Set}$.

Theorem

The following conditions are equivalent:

A **model** for a Lawvere theory p is a product-preserving functor $\ell : \mathcal{L} \rightarrow \mathbf{Set}$.

The category $\mathbf{Mod}(p)$ for a Lawvere theory is a full, **reflective** subcategory of the category $[\mathcal{L}, \mathbf{Set}]$ of all functors $\mathcal{L} \rightarrow \mathbf{Set}$.

Theorem

The following conditions are equivalent:

- *ℓ is a model for a Lawvere theory \mathcal{L} ;*

A **model** for a Lawvere theory p is a product-preserving functor $\ell : \mathcal{L} \rightarrow \mathbf{Set}$.

The category $\mathbf{Mod}(p)$ for a Lawvere theory is a full, **reflective** subcategory of the category $[\mathcal{L}, \mathbf{Set}]$ of all functors $\mathcal{L} \rightarrow \mathbf{Set}$.

Theorem

The following conditions are equivalent:

- *ℓ is a model for a Lawvere theory \mathcal{L} ;*
- *The composition $\ell \circ p$ preserves finite products;*

A **model** for a Lawvere theory p is a product-preserving functor $\ell : \mathcal{L} \rightarrow \mathbf{Set}$.

The category $\mathbf{Mod}(p)$ for a Lawvere theory is a full, **reflective** subcategory of the category $[\mathcal{L}, \mathbf{Set}]$ of all functors $\mathcal{L} \rightarrow \mathbf{Set}$.

Theorem

The following conditions are equivalent:

- *ℓ is a model for a Lawvere theory \mathcal{L} ;*
- *The composition $\ell \circ p$ preserves finite products;*
- *The composition $\ell \circ p$ is representable (with respect to the inclusion $J : \mathbf{Fin} \rightarrow \mathbf{Set}$), i.e.*

$$\ell(X[n]) \cong \mathbf{Set}(J[n], A)$$

for some $A \in \mathbf{Set}$.

Proof of reflectiveness

The category $\text{Mod}(p)$ is reflective:

Proof of reflectiveness

The category $\text{Mod}(p)$ is reflective:

A functor $F : \mathcal{L} \rightarrow \text{Set}$ preserves products if and only if it is orthogonal with respect to all σ_{AB} in

$$\begin{array}{ccc} yA \amalg yB & \xrightarrow{\forall t} & F \\ \forall \sigma_{AB} \downarrow & \nearrow & \\ y(A \times B) & & \end{array}$$

Theorem

Let \mathcal{E} be a locally presentable category and $\Sigma \subset \text{hom}(\mathcal{E})$ a set of morphism with (finitely) presentable domain; then the subcategory of Σ -orthogonal object is always reflective and (finitely) accessibly embedded.

As a consequence of the previous theorem, the square

$$\begin{array}{ccc} \text{Mod}(p) & \xrightarrow{r} & [\mathcal{L}, \text{Set}] \\ u \downarrow & & \downarrow _ \circ X \\ \text{Set} & \xrightarrow{[J, 1]} & [\text{Fin}^0, \text{Set}] \end{array}$$

is a pullback.

As a consequence of the previous theorem, the square

$$\begin{array}{ccc} \mathrm{Mod}(p) & \xrightarrow{r} & [\mathcal{L}, \mathrm{Set}] \\ u \downarrow & & \downarrow_{-\circ X} \\ \mathrm{Set} & \xrightarrow{[J, 1]} & [\mathrm{Fin}^0, \mathrm{Set}] \end{array}$$

is a pullback.

As a consequence of the previous theorem, the square

$$\begin{array}{ccc} \mathbf{Mod}(p) & \xrightarrow{r} & [\mathcal{L}, \mathbf{Set}] \\ u \downarrow & & \downarrow - \circ X \\ \mathbf{Set} & \xrightarrow{[J, 1]} & [\mathbf{Fin}^0, \mathbf{Set}] \end{array}$$

is a pullback.

- $\mathbf{Mod}(p)$ is a **reflective** subcategory of $[\mathcal{L}, \mathbf{Set}]$. We write $r_! \dashv r$ for the resulting adjunction.

As a consequence of the previous theorem, the square

$$\begin{array}{ccc} \text{Mod}(p) & \xrightarrow{r} & [\mathcal{L}, \text{Set}] \\ u \downarrow & & \downarrow - \circ X \\ \text{Set} & \xrightarrow{[J, 1]} & [\text{Fin}^0, \text{Set}] \end{array}$$

is a pullback.

- $\text{Mod}(p)$ is a **reflective** subcategory of $[\mathcal{L}, \text{Set}]$. We write $r_! \dashv r$ for the resulting adjunction.
- The functor u is **monadic**, with left adjoint f .

As a consequence of the previous theorem, the square

$$\begin{array}{ccc}
 \text{Mod}(p) & \xrightarrow{r} & [\mathcal{L}, \text{Set}] \\
 u \downarrow & & \downarrow - \circ X \\
 \text{Set} & \xrightarrow{[J, 1]} & [\text{Fin}^0, \text{Set}]
 \end{array}$$

is a pullback.

- $\text{Mod}(p)$ is a **reflective** subcategory of $[\mathcal{L}, \text{Set}]$. We write $r_! \dashv r$ for the resulting adjunction.
- The functor u is **monadic**, with left adjoint f .
- This sets up a functor

$$\mathfrak{M} : \text{Th}_L(\text{Fin}) \rightarrow \text{Mnd}_{<\omega}(\text{Set})$$

because the monad uf above is finitary.

Proof of monadicity

- **Monadicity of u :** a monadic functor has a left adjoint, reflects isomorphisms, and creates u -split coequalizers (those parallel pairs that u sends to split coequalizers, have a coequalizer, that u preserves).

Apart from the existence of f , all properties are stable under pullback.

Proof of monadicity

- **Monadicity of u :** a monadic functor has a left adjoint, reflects isomorphisms, and creates u -split coequalizers (those parallel pairs that u sends to split coequalizers, have a coequalizer, that u preserves).
Apart from the existence of f , all properties are stable under pullback.
- **u commutes with filtered colimits:** it is representable by a finitely presentable object.

$$u(\ell) = \ell[1] \cong [\mathcal{L}, \mathbf{Set}](y[1], \ell)$$

Proof of monadicity

- Being conservative is stable under pullback: conservative functors are a right orthogonal class, precisely ι^\perp where $\iota : \{0 \rightarrow 1\} \rightarrow \{0 \cong 1\}$.

Proof of monadicity

- Being conservative is stable under pullback: conservative functors are a right orthogonal class, precisely ι^\perp where $\iota : \{0 \rightarrow 1\} \rightarrow \{0 \cong 1\}$.
- Creating coequalizers of u -split pairs is stable under pullback:

$$\begin{array}{ccc} \mathcal{A} & \xrightarrow{s} & \mathcal{B} \\ u \downarrow & & \downarrow p^* \\ \mathcal{C} & \xrightarrow[t]{} & \mathcal{L} \end{array}$$

if p^* creates them, so does u .

Proof of monadicity

- Being conservative is stable under pullback: conservative functors are a right orthogonal class, precisely ι^\perp where $\iota : \{0 \rightarrow 1\} \rightarrow \{0 \cong 1\}$.
- Creating coequalizers of u -split pairs is stable under pullback:

$$\begin{array}{ccc} \mathcal{A} & \xrightarrow{s} & \mathcal{B} \\ u \downarrow & & \downarrow p^* \\ \mathcal{C} & \xrightarrow[t]{} & \mathcal{L} \end{array}$$

if p^* creates them, so does u .

- Every inverse image is monadic.

$$\mathrm{Th}_L(\mathbf{Fin}) \cong \mathrm{Mnd}_{<\omega}(\mathbf{Set})$$

Construct a functor in the opposite direction,

$$\exists : \mathrm{Mnd}_{<\omega}(\mathbf{Set}) \rightarrow \mathrm{Th}_L(\mathbf{Fin});$$

$$\mathrm{Th}_L(\mathbf{Fin}) \cong \mathrm{Mnd}_{<\omega}(\mathbf{Set})$$

Construct a functor in the opposite direction,

$$\exists : \mathrm{Mnd}_{<\omega}(\mathbf{Set}) \rightarrow \mathrm{Th}_L(\mathbf{Fin});$$

given T , we consider the composition $\mathbf{Fin} \hookrightarrow \mathbf{Set} \xrightarrow{F^T} \mathbf{Set}^T$
and its bo-ff factorization,

$$\begin{array}{ccc} \mathcal{L}^o & \xrightarrow{ff} & \mathbf{Set}^T \\ \uparrow b & & \uparrow F^T \\ \mathbf{Fin} & \xrightarrow{J} & \mathbf{Set} \end{array}$$

$$\mathrm{Th}_L(\mathbf{Fin}) \cong \mathrm{Mnd}_{<\omega}(\mathbf{Set})$$

Construct a functor in the opposite direction,

$$\exists : \mathrm{Mnd}_{<\omega}(\mathbf{Set}) \rightarrow \mathrm{Th}_L(\mathbf{Fin});$$

given T , we consider the composition $\mathbf{Fin} \hookrightarrow \mathbf{Set} \xrightarrow{F^T} \mathbf{Set}^T$
and its bo-ff factorization,

$$\begin{array}{ccc} \mathcal{L}^o & \xrightarrow{ff} & \mathbf{Set}^T \\ \uparrow b & & \uparrow F^T \\ \mathbf{Fin} & \xrightarrow{J} & \mathbf{Set} \end{array}$$

- the left vertical arrow is a Lawvere theory.

$$\mathrm{Th}_L(\mathbf{Fin}) \cong \mathrm{Mnd}_{<\omega}(\mathbf{Set})$$

Construct a functor in the opposite direction,

$$\exists : \mathrm{Mnd}_{<\omega}(\mathbf{Set}) \rightarrow \mathrm{Th}_L(\mathbf{Fin});$$

given T , we consider the composition $\mathbf{Fin} \hookrightarrow \mathbf{Set} \xrightarrow{F^T} \mathbf{Set}^T$
and its bo-ff factorization,

$$\begin{array}{ccc} \mathcal{L}^0 & \xrightarrow{ff} & \mathbf{Set}^T \\ \uparrow b & & \uparrow F^T \\ \mathbf{Fin} & \xrightarrow{J} & \mathbf{Set} \end{array}$$

- the left vertical arrow is a Lawvere theory.

- $\mathbf{Set}^T \cong \mathcal{L}\text{-models}$:

$$\begin{array}{ccc} \mathbf{Set}^T & \xrightarrow{\quad} & [\mathcal{L}, \mathbf{Set}] \\ \downarrow & \lrcorner & \downarrow [b^0, \mathbf{Set}] \\ \mathbf{Set} & \xrightarrow{[J, 1]} & [\mathbf{Fin}^0, \mathbf{Set}] \end{array}$$

Theories as promonads

There is a 2-monad $\tilde{S} : \mathbf{Prof} \rightarrow \mathbf{Prof}$ whose algebras are exactly promonoidal categories.

Theories as promonads

There is a 2-monad $\tilde{S} : \mathbf{Prof} \rightarrow \mathbf{Prof}$ whose algebras are exactly promonoidal categories.

Given a profunctor $p : \mathcal{A} \rightsquigarrow \mathcal{B}$ between promonoidal categories $(\mathcal{A}, \mathfrak{P}, J_A), (\mathcal{B}, \mathfrak{Q}, J_B)$:

Theories as promonads

There is a 2-monad $\tilde{S} : \mathbf{Prof} \rightarrow \mathbf{Prof}$ whose algebras are exactly promonoidal categories.

Given a profunctor $p : \mathcal{A} \rightsquigarrow \mathcal{B}$ between promonoidal categories $(\mathcal{A}, \mathfrak{P}, J_A), (\mathcal{B}, \mathfrak{Q}, J_B)$:

- p is a **pseudo- \tilde{S} -algebra** morphism;

Theories as promonads

There is a 2-monad $\tilde{S} : \mathbf{Prof} \rightarrow \mathbf{Prof}$ whose algebras are exactly promonoidal categories.

Given a profunctor $p : \mathcal{A} \rightsquigarrow \mathcal{B}$ between promonoidal categories $(\mathcal{A}, \mathfrak{P}, J_A), (\mathcal{B}, \mathfrak{Q}, J_B)$:

- p is a **pseudo- \tilde{S} -algebra** morphism;
- The cocontinuous left adjoint \hat{p} associated to p is **strong monoidal** with respect to the convolution monoidal product on presheaf categories;

Theories as promonads

There is a 2-monad $\tilde{S} : \mathbf{Prof} \rightarrow \mathbf{Prof}$ whose algebras are exactly promonoidal categories.

Given a profunctor $p : \mathcal{A} \rightsquigarrow \mathcal{B}$ between promonoidal categories $(\mathcal{A}, \mathfrak{P}, J_A), (\mathcal{B}, \mathfrak{Q}, J_B)$:

- p is a **pseudo- \tilde{S} -algebra** morphism;
- The cocontinuous left adjoint \hat{p} associated to p is **strong monoidal** with respect to the convolution monoidal product on presheaf categories;

If $\mathfrak{P}, \mathfrak{Q}$ on \mathcal{A}, \mathcal{B} are representable then

Theories as promonads

There is a 2-monad $\tilde{S} : \mathbf{Prof} \rightarrow \mathbf{Prof}$ whose algebras are exactly promonoidal categories.

Given a profunctor $p : \mathcal{A} \rightsquigarrow \mathcal{B}$ between promonoidal categories $(\mathcal{A}, \mathfrak{P}, J_A), (\mathcal{B}, \mathfrak{Q}, J_B)$:

- p is a **pseudo- \tilde{S} -algebra** morphism;
- The cocontinuous left adjoint \hat{p} associated to p is **strong monoidal** with respect to the convolution monoidal product on presheaf categories;

If $\mathfrak{P}, \mathfrak{Q}$ on \mathcal{A}, \mathcal{B} are representable then

- Both mates $p^{\triangleleft} : \mathcal{A} \rightarrow P\mathcal{B}$ and $p^{\triangle} : \mathcal{B} \rightarrow P^*\mathcal{A}$ are **strong monoidal** wrt convolution on their codomains.

Theories as $[\mathbf{Fin}, \mathbf{Set}]$ -categories

Theorem

$$[\mathbf{Fin}, \mathbf{Set}] \cong \mathbf{End}_{<\omega}(\mathbf{Set})$$

Proof: use Yoneda lemma.

Theories as $[\mathbf{Fin}, \mathbf{Set}]$ -categories

Theorem

$$[\mathbf{Fin}, \mathbf{Set}] \cong \mathbf{End}_{<\omega}(\mathbf{Set})$$

Proof: use Yoneda lemma.

Equivalence is monoidal; the \circ -transported structure is called the **substitution** monoidal product of functors

$F, G : \mathbf{Fin} \rightarrow \mathbf{Set}$:

$$F \ominus G : m \mapsto \int^n Fn \times (Gm)^n$$

Theories as $[\mathbf{Fin}, \mathbf{Set}]$ -categories

Theorem

$$[\mathbf{Fin}, \mathbf{Set}] \cong \mathbf{End}_{<\omega}(\mathbf{Set})$$

Proof: use Yoneda lemma.

Equivalence is monoidal; the \circ -transported structure is called the **substitution** monoidal product of functors

$F, G : \mathbf{Fin} \rightarrow \mathbf{Set}$:

$$F \ominus G : m \mapsto \int^n F n \times (G m)^n$$

Substitution is (highly!) non-symmetric, right closed monoidal structure (not left closed).

Theories as $[\mathbf{Fin}, \mathbf{Set}]$ -categories

Theorem

$$[\mathbf{Fin}, \mathbf{Set}] \cong \mathbf{End}_{<\omega}(\mathbf{Set})$$

Proof: use Yoneda lemma.

Equivalence is monoidal; the \circ -transported structure is called the **substitution** monoidal product of functors

$F, G : \mathbf{Fin} \rightarrow \mathbf{Set}$:

$$F \ominus G : m \mapsto \int^n F n \times (G m)^n$$

Substitution is (highly!) non-symmetric, right closed monoidal structure (not left closed).

The category $[\mathbf{Fin}, \mathbf{Set}]$ works as base of enrichment.

Theories as $[\mathbf{Fin}, \mathbf{Set}]$ -categories

From [Garner]

From now on we blur the distinction between the categories $[\mathbf{Fin}, \mathbf{Set}] \cong \mathbf{End}_{<\omega}(\mathbf{Set}) = \mathcal{W}$:

Theories as $[\mathbf{Fin}, \mathbf{Set}]$ -categories

From [Garner]

From now on we blur the distinction between the categories $[\mathbf{Fin}, \mathbf{Set}] \cong \mathbf{End}_{<\omega}(\mathbf{Set}) = \mathcal{W}$:

- A **finitary monad** is a monoid in \mathcal{W} , i.e. a \mathcal{W} -category with a single object;

Theories as $[\mathbf{Fin}, \mathbf{Set}]$ -categories

From [Garner]

From now on we blur the distinction between the categories $[\mathbf{Fin}, \mathbf{Set}] \cong \mathbf{End}_{<\omega}(\mathbf{Set}) = \mathcal{W}$:

- A **finitary monad** is a monoid in \mathcal{W} , i.e. a \mathcal{W} -category with a single object;
- A Lawvere theory is a \mathcal{W} -category that is **absolute** (=Cauchy-, =Karoubi-)**complete** as an enriched category and generated by a single object.

Theories as $[\mathbf{Fin}, \mathbf{Set}]$ -categories

From [Garner]

From now on we blur the distinction between the categories $[\mathbf{Fin}, \mathbf{Set}] \cong \mathbf{End}_{<\omega}(\mathbf{Set}) = \mathcal{W}$:

- A **finitary monad** is a monoid in \mathcal{W} , i.e. a \mathcal{W} -category with a single object;
- A Lawvere theory is a \mathcal{W} -category that is **absolute** (=Cauchy-, =Karoubi-)**complete** as an enriched category and generated by a single object.

Lawvere theories form a reflective subcategory in finitary monads; reflection is the enriched **Cauchy completion** functor.

Theories as \mathcal{W} -categories

In this perspective there is no difference between a Lawvere theory and its associated monad: they are the very same thing, up to a Cauchy-completion operation.

Theories as \mathcal{W} -categories

In this perspective there is no difference between a Lawvere theory and its associated monad: they are the very same thing, up to a Cauchy-completion operation.

(The Cauchy completion of a monoid in \mathbf{Cat} is rarely a monoid: take the “generic idempotent” $M = \{1, e\}$ and split $e : * \rightarrow *$ as $r : 0 \hookrightarrow * : s$).

Theories as \mathcal{W} -categories

In this perspective there is no difference between a Lawvere theory and its associated monad: they are the very same thing, up to a Cauchy-completion operation.

(The Cauchy completion of a monoid in \mathbf{Cat} is rarely a monoid: take the “generic idempotent” $M = \{1, e\}$ and split $e : * \rightarrow *$ as $r : 0 \hookrightarrow * : s$).

In order to add all \mathcal{W} -absolute colimits, at least all tensors $y[n] \odot X$ must be added to the single object X .

Theories as \mathcal{W} -categories

Equivalently,

- A Lawvere \mathcal{W} -category is an enriched category where every object A is the tensor $y[n] \odot X$ for a distinguished object $X \cong y[1] \odot X$. All such categories are \mathcal{W} -absolute complete.
- A \mathcal{W} -category is a special kind of **cartesian multicategory**: one where a multimorphism $f : X_1 \dots X_n \rightarrow Y$ is such that $X_1 = X_2 = \dots = X_n$.

Generalisations/extensions:

Generalisations/extensions:

- let \mathbb{N} be the discrete category over natural numbers;

Generalisations/extensions:

- let \mathbb{N} be the **discrete** category over natural numbers;
- let \mathbb{P} be the **groupoid** of natural numbers;

Generalisations/extensions:

- let \mathbb{N} be the **discrete** category over natural numbers;
- let \mathbf{P} be the **groupoid** of natural numbers;

The categories $[\mathbb{N}, \mathbf{Set}]$ and $[\mathbf{P}, \mathbf{Set}]$ become monoidal with respect to substitution products \ominus_N, \ominus_P :

$$F \ominus_N G : n \mapsto \coprod_{k \in \mathbb{N}} G_k \times \coprod_{\vec{n} : \sum n_i = n} X_{n_1} \times \cdots \times X_{n_k}$$

$$F \ominus_P G : n \mapsto \int^{k, \vec{n}} Y_k \times X_{n_1} \times \cdots \times X_{n_k} \times \mathbf{P}(\sum n_i, n)$$

PRO(P)S

\ominus_N and \ominus_P -monoids are respectively non-symmetric and symmetric operads.

PRO(P)S

\ominus_N and \ominus_P -monoids are respectively non-symmetric and symmetric operads.

- A **PRO** is an identity-on-objects strong monoidal functor $p : \mathbb{N}^0 \rightarrow \mathcal{P}$. \mathcal{P} is possibly non-cartesian.

PRO(P)S

\ominus_N and \ominus_P -monoids are respectively non-symmetric and symmetric operads.

- A **PRO** is an identity-on-objects strong monoidal functor $p : \mathbb{N}^0 \rightarrow \mathcal{P}$. \mathcal{P} is possibly non-cartesian.
- A **PROP** is an identity-on-objects strong monoidal functor $p : \mathbb{N}^0 \rightarrow \mathcal{P}$. \mathcal{P} is symmetric monoidal.

PRO(P)S

\ominus_N and \ominus_P -monoids are respectively non-symmetric and symmetric operads.

- A **PRO** is an identity-on-objects strong monoidal functor $p : \mathbb{N}^0 \rightarrow \mathcal{P}$. \mathcal{P} is possibly non-cartesian.
- A **PROP** is an identity-on-objects strong monoidal functor $p : \mathbb{N}^0 \rightarrow \mathcal{P}$. \mathcal{P} is symmetric monoidal.

Still examples of promonoidal promonads and symmetric promonoidal promonads.

PRO(P)s and operads

Every PRO $p : \mathbb{N}^0 \rightarrow \mathcal{T}$ gives rise to the operad $O(\mathcal{T}) = (\mathcal{T}(n, 1) \mid n \in \mathbb{N})$.

PRO(P)s and operads

Every PRO $p : \mathbb{N}^0 \rightarrow \mathcal{T}$ gives rise to the operad $\mathcal{O}(\mathcal{T}) = (\mathcal{T}(n, 1) \mid n \in \mathbb{N})$.

Conversely, any operad $(\mathcal{O}(n) \mid n \in \mathbb{N})$ gives rise to a pro $T(\mathcal{O})$, where

$$T(\mathcal{O})(n, m) = \coprod_{k_1 + \dots + k_m = n} \mathcal{O}(k_1) \times \dots \times \mathcal{O}(k_m).$$

PRO(P)s and operads

Every PRO $p : \mathbb{N}^0 \rightarrow \mathcal{T}$ gives rise to the operad $\mathcal{O}(\mathcal{T}) = (\mathcal{T}(n, 1) \mid n \in \mathbb{N})$.

Conversely, any operad $(\mathcal{O}(n) \mid n \in \mathbb{N})$ gives rise to a pro $T(\mathcal{O})$, where

$$T(\mathcal{O})(n, m) = \coprod_{k_1 + \dots + k_m = n} \mathcal{O}(k_1) \times \dots \times \mathcal{O}(k_m).$$

(It would be helpful to imagine a picture of m trees stacked vertically.)

PRO(P)s and operads

Every PRO $p : \mathbb{N}^0 \rightarrow \mathcal{T}$ gives rise to the operad $O(\mathcal{T}) = (\mathcal{T}(n, 1) \mid n \in \mathbb{N})$.

Conversely, any operad $(\mathcal{O}(n) \mid n \in \mathbb{N})$ gives rise to a pro $T(\mathcal{O})$, where

$$T(\mathcal{O})(n, m) = \coprod_{k_1 + \dots + k_m = n} \mathcal{O}(k_1) \times \dots \times \mathcal{O}(k_m).$$

(It would be helpful to imagine a picture of m trees stacked vertically.)

If we begin with an operad \mathcal{O} , we have $\mathcal{O} = O(T(\mathcal{O}))$. (This is because $T(\mathcal{O})(n, 1) = \mathcal{O}(n)$, according to the above formula.)

On the other hand, if we start with a PRO \mathcal{T} , then there exists a canonical map of PROs $T(O(\mathcal{T})) \rightarrow \mathcal{T}$, given by, for each n and m , a canonical function

$$\coprod_{k_1 + \dots + k_m = n} \mathcal{T}(k_1, 1) \times \dots \times \mathcal{T}(k_m, 1) \rightarrow \mathcal{T}(n, m) \quad (\star)$$

induced from the monoidal product on \mathcal{T} .

On the other hand, if we start with a PRO \mathcal{T} , then there exists a canonical map of PROs $T(O(\mathcal{T})) \rightarrow \mathcal{T}$, given by, for each n and m , a canonical function

$$\coprod_{k_1 + \dots + k_m = n} \mathcal{T}(k_1, 1) \times \dots \times \mathcal{T}(k_m, 1) \rightarrow \mathcal{T}(n, m) \quad (\star)$$

induced from the monoidal product on \mathcal{T} .

This sets up an adjunction

$$T : \mathbf{Opd}[\mathbf{S}] \rightleftarrows \mathbf{PRO}[\mathbf{P}] : O$$

with fully faithful left adjoint, so that [symmetric] operads can be regarded as a PRO[P]s \mathcal{T} such that each function (\star) is bijective.

The evil plan

Re-enact [Garner] away from Set.

The evil plan

Re-enact [Garner] away from \mathbf{Set} .

Let \mathcal{V} be a locally presentable base of enrichment; let $\mathfrak{F}(\mathcal{V})$ be the subcategory of finitely presentable objects:

The evil plan

Re-enact [Garner] away from \mathbf{Set} .

Let \mathcal{V} be a locally presentable base of enrichment; let $\mathfrak{F}(\mathcal{V})$ be the subcategory of finitely presentable objects:

- $\mathfrak{F}(\mathcal{V})$ is the free **finite weighted** cocompletion of the point;

The evil plan

Re-enact [Garner] away from \mathbf{Set} .

Let \mathcal{V} be a locally presentable base of enrichment; let $\mathfrak{F}(\mathcal{V})$ be the subcategory of finitely presentable objects:

- $\mathfrak{F}(\mathcal{V})$ is the free **finite weighted** cocompletion of the point;
- There is a strong monoidal equivalence of categories

$$[\mathfrak{F}(\mathcal{V}), \mathcal{V}] \cong [\mathcal{V}, \mathcal{V}]_{<\omega}$$

between functors $\mathfrak{F}(\mathcal{V}) \rightarrow \mathcal{V}$ and finitary endo- \mathcal{V} -functors;

The evil plan

- \mathcal{V} -substitution is

$$F * G = A \mapsto \int^B FB \otimes_{\mathcal{V}} (GA)^B \leftarrow \mathcal{V}\text{-power}$$

The evil plan

- \mathcal{V} -substitution is

$$F * G = A \mapsto \int^B FB \otimes_{\mathcal{V}} (GA)^B \leftarrow \mathcal{V}\text{-power}$$

- Equivalence between **finitary \mathcal{V} -monads** and **enriched-Cauchy-complete categories generated by a single object** under iterated finite powers.

The evil plan

- \mathcal{V} -substitution is

$$F * G = A \mapsto \int^B FB \otimes_{\mathcal{V}} (GA)^B \leftarrow \mathcal{V}\text{-power}$$

- Equivalence between **finitary \mathcal{V} -monads** and **enriched-Cauchy-complete categories generated by a single object** under iterated finite powers.
- **Models** for a Lawvere theory correspond to **algebras** for the associated finitary monad; free models are free algebras are representables in

$$\begin{aligned}\mathrm{Alg}(T, \mathcal{C}) &= [\mathfrak{F}(\mathcal{V}), \mathcal{V}]\text{-Cat}(T, \mathcal{C}) \\ (\text{Cauchy compl.}) &\cong [\mathfrak{F}(\mathcal{V}), \mathcal{V}]\text{-Cat}(\hat{T}, \mathcal{C}) \\ &= \mathrm{Mod}(\hat{T}, \mathcal{C})\end{aligned}$$

The evil plan

class of lims	finite \times	\mathbb{D} -lims	finite powers	weighted \mathbb{D} -limits	bicat \times
basic theory	\mathbf{Fin}^0	completion of $\{*\}$	completion of $\{*\}$	completion of $\{*\}$	completion of $\{*\}$
semantics in	Set	Set	\mathcal{V}	\mathcal{V}	Prof
eq. with _ monads	finitary	\mathbb{D} -accessible	$[\mathfrak{F}(V), V]$ -monoids	$[?, V]$ -monoids	???

Profunctorial semantics

- Characterise the free carbicat $\mathbb{CB}(\ast)$ on a singleton: see [link here](#);

Profunctorial semantics

- Characterise the free carbicat $\mathbb{CB}(\ast)$ on a singleton: see [link here](#));
- Check if the univ property of \mathbf{Fin} remains true for $\mathbb{CB}(\ast)$;

Profunctorial semantics

- Characterise the free carbicat $\mathbb{CB}(\ast)$ on a singleton: see [link here](#));
- Check if the univ property of \mathbf{Fin} remains true for $\mathbb{CB}(\ast)$;
- Take $\mathbb{CB}(\ast) = F$, and consider its **free cocompletion** in the bicolimit sense

Profunctorial semantics

- Characterise the free carbicat $\mathbb{CB}(\ast)$ on a singleton: see [link here](#));
- Check if the univ property of \mathbf{Fin} remains true for $\mathbb{CB}(\ast)$;
- Take $\mathbb{CB}(\ast) = F$, and consider its **free cocompletion** in the bicolimit sense
- Prove that

$$\begin{aligned}[PF, PF] &\cong [\mathbb{CB}(\ast), PF] \\ &\cong PF\end{aligned}$$

monoidally; \odot -**monoids** := monoids in PF wrt composition in $[PF, PF]$.

Profunctorial semantics

- Prove that there is a **syntax-VS-semantics** adjunction here: theories are promonoidal promonads T on (a 1-skeleton of) $\mathbb{CB}(\ast)$, and models are carbicat homomorphisms $\mathrm{Kl}(T) \rightarrow \mathrm{Prof}$. There is an equivalence

$$\{\text{theories}\} \cong \{\text{??? monads}\}$$

Profunctorial semantics

- Prove that there is a **syntax-VS-semantics** adjunction here: theories are promonoidal promonads T on (a 1-skeleton of) $\mathbb{CB}(\ast)$, and models are carbicat homomorphisms $\mathbf{Kl}(T) \rightarrow \mathbf{Prof}$. There is an equivalence

$$\{\text{theories}\} \cong \{\text{??? monads}\}$$

- Let PROs come into play: analogue of the adjunction between PROs and operads.

Bibliography

- Lawvere, F. William. "Functorial semantics of algebraic theories." Proceedings of the National Academy of Sciences of the United States of America 50.5 (1963): 869.
- Linton, Fred E.J. "Some aspects of equational categories." Proceedings of the Conference on Categorical Algebra. Springer, Berlin, Heidelberg, 1966.
- Garner, Richard. "Lawvere theories, finitary monads and Cauchy-completion." Journal of Pure and Applied Algebra 218.11 (2014): 1973-1988.
- Nishizawa, Koki, and John Power. "Lawvere theories enriched over a general base." Journal of Pure and Applied Algebra 213.3 (2009): 377-386.
- Hyland, Martin, and John Power. "The category theoretic understanding of universal algebra: Lawvere theories and monads." Electronic Notes in Theoretical Computer Science 172 (2007): 437-458.

