Minimal and intrinsic topologies on monoids of elementary embeddings

Paolo Marimon
joint work with J. de la Nuez Gonzalez, Zaniar Ghadernezhad, and Michael Pinsker

University of Oxford

University of East Anglia, 3 November 2026



Outline

1. Preliminaries
2. The Zariski topology
3. Proving minimality

$\mathrm{Aut}(\mathbb{A})$ and $\mathrm{EEmb}(\mathbb{A})$.

$\mathbb{A}$:= a (first-order) structure.

We are interested in two natural spaces of symmetries:

- $\mathrm{Aut}(\mathbb{A})$:= automorphism group of $\mathbb{A}$;
- $\mathrm{EEmb}(\mathbb{A})$:= monoid of elementary embeddings of $\mathbb{A}$ ($f : \mathbb{A} \to \mathbb{A}$ preserving the truth of first-order formulas).

$\mathrm{Aut}(\mathbb{A})$ and $\mathrm{EEmb}(\mathbb{A})$.

$\mathbb{A}$:= a (first-order) structure.

We are interested in two natural spaces of symmetries:

- $\mathrm{Aut}(\mathbb{A})$:= automorphism group of $\mathbb{A}$;
- $\mathrm{EEmb}(\mathbb{A})$:= monoid of elementary embeddings of $\mathbb{A}$ ($f : \mathbb{A} \to \mathbb{A}$ preserving the truth of first-order formulas).

$\mathrm{Aut}(\mathbb{A})$ and $\mathrm{EEmb}(\mathbb{A})$.

$\mathbb{A}$:= a (first-order) structure.

We are interested in two natural spaces of symmetries:

- $\mathrm{Aut}(\mathbb{A})$:= automorphism group of $\mathbb{A}$;
- $\mathrm{EEmb}(\mathbb{A})$:= monoid of elementary embeddings of $\mathbb{A}$ ($f : \mathbb{A} \to \mathbb{A}$ preserving the truth of first-order formulas).

$\mathrm{Aut}(\mathbb{A})$ and $\mathrm{EEmb}(\mathbb{A})$.

$\mathbb{A}$:= a (first-order) structure.
We are interested in two natural spaces of symmetries:

- $\mathrm{Aut}(\mathbb{A})$:= automorphism group of $\mathbb{A}$;
- $\mathrm{EEmb}(\mathbb{A})$:= monoid of elementary embeddings of $\mathbb{A}$ ($f : \mathbb{A} \to \mathbb{A}$ preserving the truth of first-order formulas).

Examples

- $\mathbb{A} := (\mathbb{N}; =)$.
  - $\mathrm{Aut}(\mathbb{A}) := \mathrm{Sym}(\mathbb{N})$;
  - $\mathrm{EEmb}(\mathbb{A}) := \mathrm{Inj}(\mathbb{N})$.
- $\mathbb{A} := (\mathbb{F}_q^{\aleph_0}; 0; +; (\lambda_r)_{r\in\mathbb{F}_q})$ ($\aleph_0$-dimensional vector space over $\mathbb{F}_q$).
  - $\mathrm{Aut}(\mathbb{A}) := \mathrm{GL}(\aleph_0, \mathbb{F}_q)$;
  - $\mathrm{EEmb}(\mathbb{A}) :=$ injective linear transformations $T : \mathbb{F}_q^{\aleph_0} \to \mathbb{F}_q^{\aleph_0}$.

$\mathrm{Aut}(\mathbb{A})$ and $\mathrm{EEmb}(\mathbb{A})$.

$\mathbb{A}$:= a (first-order) structure.
We are interested in two natural spaces of symmetries:

- $\mathrm{Aut}(\mathbb{A})$:= automorphism group of $\mathbb{A}$;
- $\mathrm{EEmb}(\mathbb{A})$:= monoid of elementary embeddings of $\mathbb{A}$ ($f : \mathbb{A} \to \mathbb{A}$ preserving the truth of first-order formulas).

Examples

- $\mathbb{A} := (\mathbb{N}; =)$.
  - $\mathrm{Aut}(\mathbb{A}) := \mathrm{Sym}(\mathbb{N})$;
  - $\mathrm{EEmb}(\mathbb{A}) := \mathrm{Inj}(\mathbb{N})$.
- $\mathbb{A} := (\mathbb{F}_q^{\aleph_0}; 0; +; (\lambda_r)_{r\in\mathbb{F}_q})$ ($\aleph_0$-dimensional vector space over $\mathbb{F}_q$).
  - $\mathrm{Aut}(\mathbb{A}) := \mathrm{GL}(\aleph_0, \mathbb{F}_q)$;
  - $\mathrm{EEmb}(\mathbb{A}) :=$ injective linear transformations $T : \mathbb{F}_q^{\aleph_0} \to \mathbb{F}_q^{\aleph_0}$.

Topologies on spaces of symmetries

We can endow $S \in \{\mathrm{Aut}(\mathbb{A}), \mathrm{EEmb}(\mathbb{A})\}$ with the **topology of pointwise convergence $\boldsymbol{\tau_{\mathrm{pw}}}$**:

$$\text{for } a, b \in \mathbb{A},\ \mathcal{U}_{a,b} := \{f \in S |\ f(a) = b\}\ ,$$

form a sub-basis of open sets for τ_{pw}.

Topologies on spaces of symmetries

We can endow $S \in \{\mathrm{Aut}(\mathbb{A}), \mathrm{EEmb}(\mathbb{A})\}$ with the **topology of pointwise convergence $\tau_{\mathbf{pw}}$**:

$$\text{for } a, b \in \mathbb{A},\ \mathcal{U}_{a,b} := \{f \in S |\ f(a) = b\}\ ,$$

form a sub-basis of open sets for τ_{pw}.
When $\mathbb{A}$ is countable, with τ_{pw}:

- $\mathrm{Aut}(\mathbb{A})$ is a Polish* topological group;
- $\mathrm{EEmb}(\mathbb{A})$ is a Polish topological monoid.

*__Polish:__ separable and completely metrizable.

ω-categorical structures

Definition (ω-categoricity)

$|\Omega| = \aleph_0$. $G \curvearrowright \Omega$ is **oligomorphic** if for all $n \in \mathbb{N}$, $G \curvearrowright \Omega^n$ has finitely many orbits in its diagonal action
$g \circ (a_1, \ldots, a_n) = (ga_1, \ldots, ga_n)$.
$\mathbb{A}$ is **ω-categorical** if $\mathrm{Aut}(\mathbb{A})$ is oligomorphic.

Examples

- $(\mathbb{N}; =)$;
- $(\mathbb{Q}; <)$;
- the countable random graph $\mathcal{R}$;
- countably infinite vector spaces over finite fields.

- In this context, $\mathrm{EEmb}(\mathbb{A})$ is the τ_{pw}-closure of $\mathrm{Aut}(\mathbb{A})$ in $\mathbb{A}^{\mathbb{A}}$.
- So, from now on, write $G := \mathrm{Aut}(\mathbb{A})$ and $\overline{G} := \mathrm{EEmb}(\mathbb{A})$.

ω-categorical structures

Definition (ω-categoricity)

$|\Omega| = \aleph_0$. $G \curvearrowright \Omega$ is **oligomorphic** if for all $n \in \mathbb{N}$, $G \curvearrowright \Omega^n$ has finitely many orbits in its diagonal action
$g \circ (a_1, \ldots, a_n) = (ga_1, \ldots, ga_n)$.
$\mathbb{A}$ is **ω-categorical** if $\mathrm{Aut}(\mathbb{A})$ is oligomorphic.

Examples

- $(\mathbb{N}; =)$;
- $(\mathbb{Q}; <)$;
- the countable random graph $\mathcal{R}$;
- countably infinite vector spaces over finite fields.

- In this context, $\mathrm{EEmb}(\mathbb{A})$ is the τ_{pw}-closure of $\mathrm{Aut}(\mathbb{A})$ in $\mathbb{A}^{\mathbb{A}}$.
- So, from now on, write $G := \mathrm{Aut}(\mathbb{A})$ and $\overline{G} := \mathrm{EEmb}(\mathbb{A})$.

ω-categorical structures

Definition (ω-categoricity)

$|\Omega| = \aleph_0$. $G \curvearrowright \Omega$ is **oligomorphic** if for all $n \in \mathbb{N}$, $G \curvearrowright \Omega^n$ has finitely many orbits in its diagonal action
$g \circ (a_1, \ldots, a_n) = (ga_1, \ldots, ga_n)$.
$\mathbb{A}$ is **ω-categorical** if $\mathrm{Aut}(\mathbb{A})$ is oligomorphic.

Examples

- $(\mathbb{N}; =)$;
- $(\mathbb{Q}; <)$;
- the countable random graph $\mathcal{R}$;
- countably infinite vector spaces over finite fields.

In this context, $\mathrm{EEmb}(\mathbb{A})$ is the τ_{pw}-closure of $\mathrm{Aut}(\mathbb{A})$ in $\mathbb{A}^{\mathbb{A}}$.

So, from now on, write $G := \mathrm{Aut}(\mathbb{A})$ and $\overline{G} := \mathrm{EEmb}(\mathbb{A})$.

ω-categorical structures

Definition (ω-categoricity)

$|\Omega| = \aleph_0$. $G \curvearrowright \Omega$ is **oligomorphic** if for all $n \in \mathbb{N}$, $G \curvearrowright \Omega^n$ has finitely many orbits in its diagonal action
$g \circ (a_1, \ldots, a_n) = (ga_1, \ldots, ga_n)$.
$\mathbb{A}$ is **ω-categorical** if $\mathrm{Aut}(\mathbb{A})$ is oligomorphic.

Examples

- $(\mathbb{N}; =)$;
- $(\mathbb{Q}; <)$;
- the countable random graph $\mathcal{R}$;
- countably infinite vector spaces over finite fields.

- In this context, $\mathrm{EEmb}(\mathbb{A})$ is the τ_{pw}-closure of $\mathrm{Aut}(\mathbb{A})$ in $\mathbb{A}^{\mathbb{A}}$.
- So, from now on, write $G := \mathrm{Aut}(\mathbb{A})$ and $\overline{G} := \mathrm{EEmb}(\mathbb{A})$.

Why care about topology?

Let $S \in \{\mathrm{Aut}(\mathbb{A}), \mathrm{EEmb}(\mathbb{A})\}$ for $\mathbb{A}$ ω-categorical.

- S with τ_{pw} encodes $\mathbb{A}$ up to bi-interpretability (Ahlbrandt and Ziegler 1986; Lascar 1991); ▸ More on bi-interpretations
- If we can distinguish τ_{pw} amongst the topologies compatible with the algebraic structure of S, S encodes $\mathbb{A}$ as a group/monoid!
- So we are interested in understanding features that τ_{pw} may have in the partial order of Polish/Hausdorff topologies compatible with the algebraic structure of S.

Why care about topology?

Let $S \in \{\mathrm{Aut}(\mathbb{A}), \mathrm{EEmb}(\mathbb{A})\}$ for $\mathbb{A}$ ω-categorical.

- S with τ_{pw} encodes $\mathbb{A}$ up to bi-interpretability (Ahlbrandt and Ziegler 1986; Lascar 1991); ▸ More on bi-interpretations
- If we can distinguish τ_{pw} amongst the topologies compatible with the algebraic structure of S, S encodes $\mathbb{A}$ as a group/monoid!
- So we are interested in understanding features that τ_{pw} may have in the partial order of Polish/Hausdorff topologies compatible with the algebraic structure of S.

Why care about topology?

Let $S \in \{\mathrm{Aut}(\mathbb{A}), \mathrm{EEmb}(\mathbb{A})\}$ for $\mathbb{A}$ ω-categorical.

- S with τ_{pw} encodes $\mathbb{A}$ up to bi-interpretability (Ahlbrandt and Ziegler 1986; Lascar 1991); ▸ More on bi-interpretations
- If we can distinguish τ_{pw} amongst the topologies compatible with the algebraic structure of S, S encodes $\mathbb{A}$ as a group/monoid!
- So we are interested in understanding features that τ_{pw} may have in the partial order of Polish/Hausdorff topologies compatible with the algebraic structure of S.

Minimality

Definition (Minimality)

A group topology τ on G is **minimal** if there are no strictly coarser Hausdorff group topologies.

Minimality

Definition (Minimality)

A group topology τ on G is **minimal** if there are no strictly coarser Hausdorff group topologies.

Question (Dikranjan and Megrelishvili 2013)

For which oligomorphic permutation groups is τ_{pw} minimal on G?

Minimality

Definition (Minimality)

A group topology τ on G is **minimal** if there are no strictly coarser Hausdorff group topologies.

Question (Dikranjan and Megrelishvili 2013)

For which oligomorphic permutation groups is τ_{pw} minimal on G?

Examples (for which τ_{pw} is minimal on $\mathrm{Aut}(\mathbb{A})$)

- $(\mathbb{N}; =)$ (Gaughan 1967);
- **stable** ω-categorical structures (Ben Yaacov and Tsankov 2016);
- the random graph, or countably infinite vector spaces over a finite field with a symplectic bilinear form (Ghadernezhad and Nuez González 2024).

Minimality

Definition (Minimality)

A group topology τ on G is **minimal** if there are no strictly coarser Hausdorff group topologies.

Question (Dikranjan and Megrelishvili 2013)

For which oligomorphic permutation groups is τ_{pw} minimal on G?

Examples (for which τ_{pw} is minimal on $\mathrm{Aut}(\mathbb{A})$)

- $(\mathbb{N}; =)$ (Gaughan 1967);
- **stable** ω-categorical structures (Ben Yaacov and Tsankov 2016);
- the random graph, or countably infinite vector spaces over a finite field with a symplectic bilinear form (Ghadernezhad and Nuez González 2024).

Minimality

Definition (Minimality)

A group topology τ on G is **minimal** if there are no strictly coarser Hausdorff group topologies.

Question (Dikranjan and Megrelishvili 2013)

For which oligomorphic permutation groups is τ_{pw} minimal on G?

Examples (for which τ_{pw} is minimal on $\mathrm{Aut}(\mathbb{A})$)

- $(\mathbb{N}; =)$ (Gaughan 1967);
- **stable** ω-categorical structures (Ben Yaacov and Tsankov 2016);
- the random graph, or countably infinite vector spaces over a finite field with a symplectic bilinear form (Ghadernezhad and Nuez González 2024).

Minimality

Definition (Minimality)

A group topology τ on G is **minimal** if there are no strictly coarser Hausdorff group topologies.

Question (Dikranjan and Megrelishvili 2013)

For which oligomorphic permutation groups is τ_{pw} minimal on G?

Examples (for which τ_{pw} is NOT minimal on $\mathrm{Aut}(\mathbb{A})$)

- $(\mathbb{Q};<)$ (Chang and Gartside 2017): we can define a coarser **interval topology** τ_{int} with, for $a, b < c \in \mathbb{Q}$:

$$\mathcal{U}_{a,(b,c)} := \{f \in \mathrm{Aut}(\mathbb{Q}) |\ f(a) \in (b,c)\}\ ,$$

forming a sub-basis of open sets.

Minimality

Definition (Minimality)

A group topology τ on G is **minimal** if there are no strictly coarser Hausdorff group topologies.

Question (Dikranjan and Megrelishvili 2013)

For which oligomorphic permutation groups is τ_{pw} minimal on G?

Examples (for which τ_{pw} is NOT minimal on $\mathrm{Aut}(\mathbb{A})$)

- the **random bipartite graph** $\mathcal{B}$: we can define a coarser **localised topology** τ_{pw}^{X} (where X is one of the $\mathrm{Aut}(\mathcal{B})$-orbits) with for $a, b \in X$:

$$\mathcal{U}_{a,b} := \{f \in \mathrm{Aut}(\mathcal{B}) |\ f(a) = b\}\ ,$$

forming a sub-basis of open sets.

Minimality

Definition (Minimality)

A semigroup topology τ on S is **minimal** if there are no strictly coarser Hausdorff semigroup topologies.

Minimality

Definition (Minimality)

A semigroup topology τ on S is **minimal** if there are no strictly coarser Hausdorff semigroup topologies.

Question

Are there any oligomorphic permutation groups for which τ_{pw} is not minimal on $\overline{G}$?†

†cf. Marimon and Pinsker 2025; Pinsker and Schindler 2023; Elliott, Jonušas, Mesyan, Mitchell, Morayne, and Peresse 2023.

Algebraic closure

Definition (Algebraic closure)


$G \curvearrowright \Omega$. For $B \subseteq \Omega$ finite,

$$\mathrm{acl}(B) := \{b \in \Omega \mid \text{the } G_B\text{-orbit of } b \text{ is finite}\}.$$

For $C \subseteq \Omega$,

$$\mathrm{acl}(C) = \bigcup_{B \subseteq C \text{ finite}} \mathrm{acl}(B)\,.$$

$\mathbb{A}$ has **no algebraicity** if $\mathrm{acl}(B) = B$ always (for $G = \mathrm{Aut}(\mathbb{A})$).



Algebraic closure

Definition (Algebraic closure)


$G \curvearrowright \Omega$. For $B \subseteq \Omega$ finite,

$$\mathrm{acl}(B) := \{b \in \Omega \mid \text{the } G_B\text{-orbit of } b \text{ is finite}\}.$$

For $C \subseteq \Omega$,

$$\mathrm{acl}(C) = \bigcup_{B \subseteq C \text{ finite}} \mathrm{acl}(B)\ .$$

$\mathbb{A}$ has **no algebraicity** if $\mathrm{acl}(B) = B$ always (for $G = \mathrm{Aut}(\mathbb{A})$).



Algebraic closure

Definition (Algebraic closure)


$G \curvearrowright \Omega$. For $B \subseteq \Omega$ finite,

$$\mathrm{acl}(B) := \{b \in \Omega \mid \text{the } G_B\text{-orbit of } b \text{ is finite}\}.$$

For $C \subseteq \Omega$,

$$\mathrm{acl}(C) = \bigcup_{B \subseteq C \text{ finite}} \mathrm{acl}(B)\ .$$

$\mathbb{A}$ has **no algebraicity** if $\mathrm{acl}(B) = B$ always (for $G = \mathrm{Aut}(\mathbb{A})$).



Algebraic closure

Definition (Algebraic closure)

$G \curvearrowright \Omega$. For $B \subseteq \Omega$ finite,

$$\mathrm{acl}(B) := \{b \in \Omega \mid \text{the } G_B\text{-orbit of } b \text{ is finite}\}.$$

For $C \subseteq \Omega$,

$$\mathrm{acl}(C) = \bigcup_{B \subseteq C \text{ finite}} \mathrm{acl}(B)\ .$$

$\mathbb{A}$ has **no algebraicity** if $\mathrm{acl}(B) = B$ always (for $G = \mathrm{Aut}(\mathbb{A})$).

Examples

- with no algebraicity: $(\mathbb{N}; =)$, $(\mathbb{Q}; <)$, $\mathcal{R}$, and $\mathcal{B}$;
- with algebraicity: vector spaces, and $\aleph_0$ disjoint copies of K_2.

Algebraic closure

Definition (Algebraic closure)

$G \curvearrowright \Omega$. For $B \subseteq \Omega$ finite,

$$\mathrm{acl}(B) := \{b \in \Omega \mid \text{the } G_B\text{-orbit of } b \text{ is finite}\}.$$

For $C \subseteq \Omega$,

$$\mathrm{acl}(C) = \bigcup_{B \subseteq C \text{ finite}} \mathrm{acl}(B)\ .$$

$\mathbb{A}$ has **no algebraicity** if $\mathrm{acl}(B) = B$ always (for $G = \mathrm{Aut}(\mathbb{A})$).

For $\mathbb{A}$ ω-categorical acl is **locally finite**: for finite B, $\mathrm{acl}(B)$ is finite.

Algebraic closure

Definition (Algebraic closure)

$G \curvearrowright \Omega$. For $B \subseteq \Omega$ finite,

$$\mathrm{acl}(B) := \{b \in \Omega \mid \text{the } G_B\text{-orbit of } b \text{ is finite}\}.$$

For $C \subseteq \Omega$,

$$\mathrm{acl}(C) = \bigcup_{B \subseteq C \text{ finite}} \mathrm{acl}(B)\ .$$

$\mathbb{A}$ has **no algebraicity** if $\mathrm{acl}(B) = B$ always (for $G = \mathrm{Aut}(\mathbb{A})$).

Theorem (Pinsker and Schindler 2023[‡])

Let $\mathbb{A}$ be ω-categorical and with no algebraicity.
Then, τ_{pw} is minimal on $\mathrm{EEmb}(\mathbb{A})$.

There are predecessors in Elliott, Jonušas, Mesyan, Mitchell, Morayne, and Peresse 2023 and Elliott, Jonušas, Mitchell, Peresse, and Pinsker 2023.

Algebraic closure

Definition (Algebraic closure)

$G \curvearrowright \Omega$. For $B \subseteq \Omega$ finite,

$$\mathrm{acl}(B) := \{b \in \Omega \mid \text{the } G_B\text{-orbit of } b \text{ is finite}\}.$$

For $C \subseteq \Omega$,

$$\mathrm{acl}(C) = \bigcup_{B \subseteq C \text{ finite}} \mathrm{acl}(B)\ .$$

$\mathbb{A}$ has **no algebraicity** if $\mathrm{acl}(B) = B$ always (for $G = \mathrm{Aut}(\mathbb{A})$).

Theorem (Pinsker and Schindler 2023)

Let $\mathbb{A}$ be ω-categorical and with no algebraicity.
Then, τ_{pw} is minimal on $\mathrm{EEmb}(\mathbb{A})$.

- This is very different from the case of $\mathrm{Aut}(\mathbb{A})$!

Algebraic closure

Definition (Algebraic closure)

$G \curvearrowright \Omega$. For $B \subseteq \Omega$ finite,

$$\mathrm{acl}(B) := \{b \in \Omega \mid \text{the } G_B\text{-orbit of } b \text{ is finite}\}.$$

For $C \subseteq \Omega$,

$$\mathrm{acl}(C) = \bigcup_{B \subseteq C \text{ finite}} \mathrm{acl}(B)\ .$$

$\mathbb{A}$ has **no algebraicity** if $\mathrm{acl}(B) = B$ always (for $G = \mathrm{Aut}(\mathbb{A})$).

Theorem (Pinsker and Schindler 2023)

Let $\mathbb{A}$ be ω-categorical and with no algebraicity.
*Then, τ_{pw} is minimal on $\mathrm{EEmb}(\mathbb{A})$ because it coincides with the **semigroup Zariski topology** τ_{Z}.*

The Zariski topology

Definition (the Zariski topology)

$\mathcal{S} :=$ a semigroup. The (semigroup) **Zariski** topology τ_{Z} has a sub-basis of open sets given by solutions to semigroup disequalities:

$$\{s \in \mathcal{S} \,|t_k s t_{k-1} s \dots t_1 s t_0 \neq q_l s q_{l-1} s \dots q_1 s q_0\} \,,$$

for $k, l \geq 1$ and $t_0, \dots, t_k, q_0, \dots q_l \in \mathcal{S}$.

τ_{Z} is contained in any Hausdorff semigroup topology on $\mathcal{S}$.
(Elliott, Jonušas, Mesyan, Mitchell, Morayne, and Peresse 2023)

The Zariski topology

Definition (the Zariski topology)

$\mathcal{S} :=$ a semigroup. The (semigroup) **Zariski** topology τ_{Z} has a sub-basis of open sets given by solutions to semigroup disequalities:

$$\{s \in \mathcal{S} \mid t_k s t_{k-1} s \dots t_1 s t_0 \neq q_l s q_{l-1} s \dots q_1 s q_0\},$$

for $k, l \geq 1$ and $t_0, \dots, t_k, q_0, \dots q_l \in \mathcal{S}$.

τ_{Z} is contained in any Hausdorff semigroup topology on $\mathcal{S}$.
(Elliott, Jonušas, Mesyan, Mitchell, Morayne, and Peresse 2023)

When is $\tau_{\mathrm{Z}} \subsetneq \tau_{\mathrm{pw}}$?

Pinsker and Schindler 2023 show that for $\mathbb{A} := \aleph_0$-many copies of K_2, $\tau_{\mathrm{Z}} \subsetneq \tau_{\mathrm{pw}}$ on $\mathrm{EEmb}(\mathbb{A})$ answering a question of Elliott, Jonušas, Mesyan, Mitchell, Morayne, and Peresse 2023.

We prove the following more general statement:

When is $\tau_{\mathrm{Z}} \subsetneq \tau_{\mathrm{pw}}$?

Pinsker and Schindler 2023 show that for $\mathbb{A} := \aleph_0$-many copies of K_2, $\tau_{\mathrm{Z}} \subsetneq \tau_{\mathrm{pw}}$ on $\mathrm{EEmb}(\mathbb{A})$ answering a question of Elliott, Jonušas, Mesyan, Mitchell, Morayne, and Peresse 2023.
We prove the following more general statement:

Theorem (de la Nuez Gonzalez, Ghadernezhad, M., Pinsker 2026)

Let $G \curvearrowright \Omega$ have locally finite algebraic closure and non-trivial centre. Then, the semigroup Zariski topology τ_{Z} on $\overline{G}$ is not Hausdorff.

When is $\tau_{\mathrm{Z}} \subsetneq \tau_{\mathrm{pw}}$?

Pinsker and Schindler 2023 show that for $\mathbb{A} := \aleph_0$-many copies of K_2, $\tau_{\mathrm{Z}} \subsetneq \tau_{\mathrm{pw}}$ on $\mathrm{EEmb}(\mathbb{A})$ answering a question of Elliott, Jonušas, Mesyan, Mitchell, Morayne, and Peresse 2023.
We prove the following more general statement:

Theorem (de la Nuez Gonzalez, Ghadernezhad, M., Pinsker 2026)

Let $G \curvearrowright \Omega$ have locally finite algebraic closure and non-trivial centre. Then, the semigroup Zariski topology τ_{Z} on $\overline{G}$ is not Hausdorff.

When is $\tau_{\mathrm{Z}} \subsetneq \tau_{\mathrm{pw}}$?

Pinsker and Schindler 2023 show that for $\mathbb{A} := \aleph_0$-many copies of K_2, $\tau_{\mathrm{Z}} \subsetneq \tau_{\mathrm{pw}}$ on $\mathrm{EEmb}(\mathbb{A})$ answering a question of Elliott, Jonušas, Mesyan, Mitchell, Morayne, and Peresse 2023.
We prove the following more general statement:

Theorem (de la Nuez Gonzalez, Ghadernezhad, M., Pinsker 2026)

Let $G \curvearrowright \Omega$ have locally finite algebraic closure and non-trivial centre. Then, the semigroup Zariski topology τ_{Z} on $\overline{G}$ is not Hausdorff.

Proof sketch

Proof sketch of the theorem.

A semigroup disequality $w(s)$

$$t_k s t_{k-1} s \dots t_1 s t_0 \neq q_l s q_{k-1} s \dots q_1 s q_0$$

is **singular** if $k = l$ and **non-singular** otherwise.

Write $\mathcal{O}_{w(s)}$ for the τ_{Z}-open set of solutions to $w(s)$.

- if $w(s)$ is singular, for $\gamma \in Z(G)$, $1 \in \mathcal{O}_{w(s)}$ if and only if $\gamma \in \mathcal{O}_{w(s)}$;
- if $w(s)$ is non-singular, then $\mathcal{O}_{w(s)}$ is dense in $\overline{G}$ (wrt τ_{pw}) (use Neumann's Lemma).
- Because of this any $\gamma \in Z(G)$ cannot be separated from 1 by τ_{Z}-open sets (in neither G nor $\overline{G}$).

□

Proof sketch

Proof sketch of the theorem.

A semigroup disequality $w(s)$

$$t_k s t_{k-1} s \dots t_1 s t_0 \neq q_l s q_{k-1} s \dots q_1 s q_0$$

is **singular** if $k = l$ and **non-singular** otherwise.
Write $\mathcal{O}_{w(s)}$ for the τ_{Z}-open set of solutions to $w(s)$.

- if $w(s)$ is singular, for $\gamma \in Z(G)$, $1 \in \mathcal{O}_{w(s)}$ if and only if $\gamma \in \mathcal{O}_{w(s)}$;
- if $w(s)$ is non-singular, then $\mathcal{O}_{w(s)}$ is dense in $\overline{G}$ (wrt τ_{pw}) (use Neumann's Lemma).
- Because of this any $\gamma \in Z(G)$ cannot be separated from 1 by τ_{Z}-open sets (in neither G nor $\overline{G}$).



Proof sketch

Proof sketch of the theorem.

A semigroup disequality $w(s)$

$$t_k s t_{k-1} s \dots t_1 s t_0 \neq q_l s q_{k-1} s \dots q_1 s q_0$$

is **singular** if $k = l$ and **non-singular** otherwise.
Write $\mathcal{O}_{w(s)}$ for the τ_{Z}-open set of solutions to $w(s)$.

- if $w(s)$ is singular, for $\gamma \in Z(G)$, $1 \in \mathcal{O}_{w(s)}$ if and only if $\gamma \in \mathcal{O}_{w(s)}$;
- if $w(s)$ is non-singular, then $\mathcal{O}_{w(s)}$ is dense in $\overline{G}$ (wrt τ_{pw}) (use Neumann's Lemma).
- Because of this any $\gamma \in Z(G)$ cannot be separated from 1 by τ_{Z}-open sets (in neither G nor $\overline{G}$).



Proof sketch

Proof sketch of the theorem.

A semigroup disequality $w(s)$

$$t_k s t_{k-1} s \dots t_1 s t_0 \neq q_l s q_{k-1} s \dots q_1 s q_0$$

is **singular** if $k = l$ and **non-singular** otherwise.
Write $\mathcal{O}_{w(s)}$ for the τ_{Z}-open set of solutions to $w(s)$.

- if $w(s)$ is singular, for $\gamma \in Z(G)$, $1 \in \mathcal{O}_{w(s)}$ if and only if $\gamma \in \mathcal{O}_{w(s)}$;
- if $w(s)$ is non-singular, then $\mathcal{O}_{w(s)}$ is dense in $\overline{G}$ (wrt τ_{pw}) (use Neumann's Lemma).
- Because of this any $\gamma \in Z(G)$ cannot be separated from 1 by τ_{Z}-open sets (in neither G nor $\overline{G}$).



Proof sketch

Proof sketch of the theorem.

A semigroup disequality $w(s)$

$$t_k s t_{k-1} s \dots t_1 s t_0 \neq q_l s q_{k-1} s \dots q_1 s q_0$$

is **singular** if $k = l$ and **non-singular** otherwise.
Write $\mathcal{O}_{w(s)}$ for the τ_{Z}-open set of solutions to $w(s)$.

- if $w(s)$ is singular, for $\gamma \in Z(G)$, $1 \in \mathcal{O}_{w(s)}$ if and only if $\gamma \in \mathcal{O}_{w(s)}$;
- if $w(s)$ is non-singular, then $\mathcal{O}_{w(s)}$ is dense in $\overline{G}$ (wrt τ_{pw}) (use Neumann's Lemma).
- Because of this any $\gamma \in Z(G)$ cannot be separated from 1 by τ_{Z}-open sets (in neither G nor $\overline{G}$).



What about minimality of τ_{pw} with algebraicity?

By our theorem, the standard method to prove minimality for τ_{pw} cannot not work in many natural ω-categorical structures with algebraicity:

- countably infinite vector spaces over finite fields;
- $\aleph_0$ disjoint copies of K_2.

What about minimality of τ_{pw} with algebraicity?

By our theorem, the standard method to prove minimality for τ_{pw} cannot not work in many natural ω-categorical structures with algebraicity:

- countably infinite vector spaces over finite fields;
- $\aleph_0$ disjoint copies of K_2.

More generally: the standard strategy of $\tau_{\mathrm{pw}} = \tau_{\mathrm{Z}}$ easily fails the moment we have algebraicity.

What about minimality of τ_{pw} with algebraicity?

By our theorem, the standard method to prove minimality for τ_{pw} cannot not work in many natural ω-categorical structures with algebraicity:

- countably infinite vector spaces over finite fields;
- $\aleph_0$ disjoint copies of K_2.


More generally: the standard strategy of $\tau_{\mathrm{pw}} = \tau_{\mathrm{Z}}$ easily fails the moment we have algebraicity.
Can we still prove minimality by other means?

Modularity

Definition (Modularity)

$G \curvearrowright \Omega$ with locally finite acl.
acl is **modular** if for finite algebraically closed A, B, C s.t. $A \subseteq B$,

$$\mathrm{acl}(A \cup (C \cap B)) = \mathrm{acl}(A \cup C) \cap B \ .$$




Modularity

Definition (Modularity)

$G \curvearrowright \Omega$ with locally finite acl.
acl is **modular** if for finite algebraically closed A, B, C s.t. $A \subseteq B$,

$$\mathrm{acl}(A \cup (C \cap B)) = \mathrm{acl}(A \cup C) \cap B \ .$$




Modularity

Definition (Modularity)

$G \curvearrowright \Omega$ with locally finite acl.
acl is **modular** if for finite algebraically closed A, B, C s.t. $A \subseteq B$,

$$\mathrm{acl}(A \cup (C \cap B)) = \mathrm{acl}(A \cup C) \cap B \ .$$




Modularity

Definition (Modularity)

$G \curvearrowright \Omega$ with locally finite acl.
acl is **modular** if for finite algebraically closed A, B, C s.t. $A \subseteq B$,

$$\mathrm{acl}(A \cup (C \cap B)) = \mathrm{acl}(A \cup C) \cap B \ .$$



Modularity

Definition (Modularity)

$G \curvearrowright \Omega$ with locally finite acl.
acl is **modular** if for finite algebraically closed A, B, C s.t. $A \subseteq B$,

$$\mathrm{acl}(A \cup (C \cap B)) = \mathrm{acl}(A \cup C) \cap B \ .$$

Examples

- whenever $\mathbb{A}$ has no algebraicity, acl is modular;
- vector spaces or projective spaces over finite fields (possibly with forms, or densely ordered).
- sometimes one needs to be more subtle (affine space over $\mathbb{F}_q$): there is $\mathbb{A}'$ with $\mathrm{EEmb}(\mathbb{A}) = \mathrm{EEmb}(\mathbb{A}')$ (with τ_{pw}) and acl is modular in $\mathbb{A}'$ (but not in $\mathbb{A}$).

Modularity

Definition (Modularity)

$G \curvearrowright \Omega$ with locally finite acl.
acl is **modular** if for finite algebraically closed A, B, C s.t. $A \subseteq B$,

$$\mathrm{acl}(A \cup (C \cap B)) = \mathrm{acl}(A \cup C) \cap B \ .$$

Examples

- whenever $\mathbb{A}$ has no algebraicity, acl is modular;
- vector spaces or projective spaces over finite fields (possibly with forms, or densely ordered).
- sometimes one needs to be more subtle (affine space over $\mathbb{F}_q$): there is $\mathbb{A}'$ with $\mathrm{EEmb}(\mathbb{A}) = \mathrm{EEmb}(\mathbb{A}')$ (with τ_{pw}) and acl is modular in $\mathbb{A}'$ (but not in $\mathbb{A}$).

Modularity

Definition (Modularity)

$G \curvearrowright \Omega$ with locally finite acl.
acl is **modular** if for finite algebraically closed A, B, C s.t. $A \subseteq B$,

$$\mathrm{acl}(A \cup (C \cap B)) = \mathrm{acl}(A \cup C) \cap B \ .$$

Examples

- whenever $\mathbb{A}$ has no algebraicity, acl is modular;
- vector spaces or projective spaces over finite fields (possibly with forms, or densely ordered).
- sometimes one needs to be more subtle (affine space over $\mathbb{F}_q$): there is $\mathbb{A}'$ with $\mathrm{EEmb}(\mathbb{A}) = \mathrm{EEmb}(\mathbb{A}')$ (with τ_{pw}) and acl is modular in $\mathbb{A}'$ (but not in $\mathbb{A}$).

Modularity

Definition (Modularity)

$G \curvearrowright \Omega$ with locally finite acl.
acl is **modular** if for finite algebraically closed A, B, C s.t. $A \subseteq B$,

$$\mathrm{acl}(A \cup (C \cap B)) = \mathrm{acl}(A \cup C) \cap B \ .$$

Morally: acl is modular when it is
"not more complicated than in a vector space".

Modularity

Definition (Modularity)

$G \curvearrowright \Omega$ with locally finite acl.
acl is **modular** if for finite algebraically closed A, B, C s.t. $A \subseteq B$,

$$\mathrm{acl}(A \cup (C \cap B)) = \mathrm{acl}(A \cup C) \cap B \ .$$

Theorem (de la Nuez Gonzalez, Ghadernezhad, M., & Pinsker 2026)

Let $G \curvearrowright \Omega$ be oligomorphic and such that acl is modular.
Then, τ_{pw} is minimal on $\overline{G}$.

Modularity

Definition (Modularity)

$G \curvearrowright \Omega$ with locally finite acl.
acl is **modular** if for finite algebraically closed A, B, C s.t. $A \subseteq B$,

$$\mathrm{acl}(A \cup (C \cap B)) = \mathrm{acl}(A \cup C) \cap B \ .$$

Theorem (de la Nuez Gonzalez, Ghadernezhad, M., & Pinsker 2026)

Let $G \curvearrowright \Omega$ be oligomorphic and such that acl is modular.
Then, τ_{pw} is minimal on $\overline{G}$.

This subsumes Pinsker and Schindler 2023.

Rigidity and dispersivity

$G \curvearrowright \Omega$. Consider a semigroup topology $\tau \subseteq \tau_{\mathrm{pw}}$ on $\overline{G}$.

Definitions (Rigidity and dispersivity)

$t \in \overline{G}$ is **τ-rigid** on $a \in \Omega$ if there is some τ-neighbourhood $\mathcal{V}$ of t such that, for all $s \in \mathcal{V}$, $s(a) \in \mathrm{acl}(t(a))$.

$t \in \overline{G}$ is **τ-dispersive** on $a \in \Omega$ if for every finite $B \subseteq \Omega$ and any τ-neighbourhood $\mathcal{V}$ of t there is $s \in \mathcal{V}$ such that $a \notin s(B)$.

- $t \in \overline{G}$ is τ_{pw}-rigid on Ω and not τ_{pw}-dispersive on $\mathrm{Im}(t)$.

Rigidity and dispersivity

$G \curvearrowright \Omega$. Consider a semigroup topology $\tau \subseteq \tau_{\mathrm{pw}}$ on $\overline{G}$.

Definitions (Rigidity and dispersivity)

$t \in \overline{G}$ is **τ-rigid** on $a \in \Omega$ if there is some τ-neighbourhood $\mathcal{V}$ of t such that, for all $s \in \mathcal{V}$, $s(a) \in \mathrm{acl}(t(a))$.
$t \in \overline{G}$ is **τ-dispersive** on $a \in \Omega$ if for every finite $B \subseteq \Omega$ and any τ-neighbourhood $\mathcal{V}$ of t there is $s \in \mathcal{V}$ such that $a \notin s(B)$.

- $t \in \overline{G}$ is τ_{pw}-rigid on Ω and not τ_{pw}-dispersive on $\mathrm{Im}(t)$.

Rigidity and dispersivity

$G \curvearrowright \Omega$. Consider a semigroup topology $\tau \subseteq \tau_{\mathrm{pw}}$ on $\overline{G}$.

Definitions (Rigidity and dispersivity)

$t \in \overline{G}$ is **τ-rigid** on $a \in \Omega$ if there is some τ-neighbourhood $\mathcal{V}$ of t such that, for all $s \in \mathcal{V}$, $s(a) \in \mathrm{acl}(t(a))$.
$t \in \overline{G}$ is **τ-dispersive** on $a \in \Omega$ if for every finite $B \subseteq \Omega$ and any τ-neighbourhood $\mathcal{V}$ of t there is $s \in \mathcal{V}$ such that $a \notin s(B)$.

- $t \in \overline{G}$ is τ_{pw}-rigid on Ω and not τ_{pw}-dispersive on $\mathrm{Im}(t)$.

Rigidity and dispersivity

$G \curvearrowright \Omega$. Consider a semigroup topology $\tau \subseteq \tau_{\mathrm{pw}}$ on $\overline{G}$.

Definitions (Rigidity and dispersivity)

$t \in \overline{G}$ is **τ-rigid** on $a \in \Omega$ if there is some τ-neighbourhood $\mathcal{V}$ of t such that, for all $s \in \mathcal{V}$, $s(a) \in \mathrm{acl}(t(a))$.
$t \in \overline{G}$ is **τ-dispersive** on $a \in \Omega$ if for every finite $B \subseteq \Omega$ and any τ-neighbourhood $\mathcal{V}$ of t there is $s \in \mathcal{V}$ such that $a \notin s(B)$.

- $t \in \overline{G}$ is τ_{pw}-rigid on Ω and not τ_{pw}-dispersive on $\mathrm{Im}(t)$.

Lemma (de la Nuez Gonzalez, Ghadernezhad, M., and Pinsker 2026)

$G \curvearrowright \Omega$ with locally finite acl. Suppose that $\tau \subsetneq \tau_{\mathrm{pw}}$ is a semigroup topology and every $t \in \overline{G}$ is τ-rigid on Ω. Then, τ is not Hausdorff.

Proof idea of the main theorem

Proof idea.

Suppose $\tau \subsetneq \tau_{\mathrm{pw}}$ is a Hausdorff semigroup topology. Consider

$$\Delta := \{a \in \Omega | \text{ every } s \in \overline{G} \text{ is } \tau\text{-rigid on } a\} .$$

By previous Lemma, $\Delta \subsetneq \Omega$.
We can find $t \in \overline{G}$ satisfying a strong form of τ-dispersivity on $\Omega \setminus \Delta$.
By **modularity**, find a "generic" copy $\mathbb{A}^\star$ of $\mathbb{A}$ within itself.
In particular,

$$\mathcal{U} := \{s \in \overline{G} | \operatorname{Im}(s) \subsetneq \mathbb{A}^\star\}$$

is τ_{Z}-open (and so τ-open) and a neighbourhood of t.
We exploit τ-dispersivity and genericity of $\mathbb{A}^\star$ to find:
$u \in \mathcal{U}$ such that $\operatorname{Im}(u) \subseteq \mathbb{A}^\star$. (CONTRADICTION!)

□

Proof idea of the main theorem

Proof idea.

Suppose $\tau \subsetneq \tau_{\mathrm{pw}}$ is a Hausdorff semigroup topology. Consider

$$\Delta := \{a \in \Omega | \text{ every } s \in \overline{G} \text{ is } \tau\text{-rigid on } a\} .$$

By previous Lemma, $\Delta \subsetneq \Omega$.

We can find $t \in \overline{G}$ satisfying a strong form of τ-dispersivity on $\Omega \setminus \Delta$.

By **modularity**, find a "generic" copy $\mathbb{A}^\star$ of $\mathbb{A}$ within itself.

In particular,

$$\mathcal{U} := \{s \in \overline{G} | \operatorname{Im}(s) \subsetneq \mathbb{A}^\star\}$$

is τ_{Z}-open (and so τ-open) and a neighbourhood of t.

We exploit τ-dispersivity and genericity of $\mathbb{A}^\star$ to find:

$u \in \mathcal{U}$ such that $\operatorname{Im}(u) \subseteq \mathbb{A}^\star$. (CONTRADICTION!)

□

Proof idea of the main theorem

Proof idea.

Suppose $\tau \subsetneq \tau_{\mathrm{pw}}$ is a Hausdorff semigroup topology. Consider

$$\Delta := \{a \in \Omega | \text{ every } s \in \overline{G} \text{ is } \tau\text{-rigid on } a\} .$$

By previous Lemma, $\Delta \subsetneq \Omega$.

We can find $t \in \overline{G}$ satisfying a strong form of τ-dispersivity on $\Omega \setminus \Delta$.

By **modularity**, find a "generic" copy $\mathbb{A}^\star$ of $\mathbb{A}$ within itself.

In particular,

$$\mathcal{U} := \{s \in \overline{G} | \operatorname{Im}(s) \subsetneq \mathbb{A}^\star\}$$

is τ_{Z}-open (and so τ-open) and a neighbourhood of t.

We exploit τ-dispersivity and genericity of $\mathbb{A}^\star$ to find:

$u \in \mathcal{U}$ such that $\operatorname{Im}(u) \subseteq \mathbb{A}^\star$. (CONTRADICTION!)

□

Proof idea of the main theorem

Proof idea.

Suppose $\tau \subsetneq \tau_{\mathrm{pw}}$ is a Hausdorff semigroup topology. Consider

$$\Delta := \{a \in \Omega | \text{ every } s \in \overline{G} \text{ is } \tau\text{-rigid on } a\} .$$

By previous Lemma, $\Delta \subsetneq \Omega$.
We can find $t \in \overline{G}$ satisfying a strong form of τ-dispersivity on $\Omega \setminus \Delta$.
By **modularity**, find a "generic" copy $\mathbb{A}^\star$ of $\mathbb{A}$ within itself.[‡]
In particular,

$$\mathcal{U} := \{s \in \overline{G} | \operatorname{Im}(s) \subsetneq \mathbb{A}^\star\}$$

is τ_{Z}-open (and so τ-open) and a neighbourhood of t.
We exploit τ-dispersivity and genericity of $\mathbb{A}^\star$ to find:
$u \in \mathcal{U}$ such that $\operatorname{Im}(u) \subseteq \mathbb{A}^\star$. (CONTRADICTION!)

□

[‡]A **lovely pair** in model theory ▸ More on this.

Proof idea of the main theorem

Proof idea.

Suppose $\tau \subsetneq \tau_{\mathrm{pw}}$ is a Hausdorff semigroup topology. Consider

$$\Delta := \{a \in \Omega | \text{ every } s \in \overline{G} \text{ is } \tau\text{-rigid on } a\} .$$

By previous Lemma, $\Delta \subsetneq \Omega$.
We can find $t \in \overline{G}$ satisfying a strong form of τ-dispersivity on $\Omega \setminus \Delta$.
By **modularity**, find a "generic" copy $\mathbb{A}^\star$ of $\mathbb{A}$ within itself.[‡]
In particular,

$$\mathcal{U} := \{s \in \overline{G} | \ \mathrm{Im}(s) \subsetneq \mathbb{A}^\star\}$$

is τ_{Z}-open (and so τ-open) and a neighbourhood of t.
We exploit τ-dispersivity and genericity of $\mathbb{A}^\star$ to find:
$u \in \mathcal{U}$ such that $\mathrm{Im}(u) \subseteq \mathbb{A}^\star$. (CONTRADICTION!)

□

[‡]A **lovely pair** in model theory ▸ More on this.

Proof idea of the main theorem

Proof idea.

Suppose $\tau \subsetneq \tau_{\mathrm{pw}}$ is a Hausdorff semigroup topology. Consider

$$\Delta := \{a \in \Omega | \text{ every } s \in \overline{G} \text{ is } \tau\text{-rigid on } a\} .$$

By previous Lemma, $\Delta \subsetneq \Omega$.
We can find $t \in \overline{G}$ satisfying a strong form of τ-dispersivity on $\Omega \setminus \Delta$.
By **modularity**, find a "generic" copy $\mathbb{A}^\star$ of $\mathbb{A}$ within itself.[‡]
In particular,

$$\mathcal{U} := \{s \in \overline{G} | \ \mathrm{Im}(s) \subsetneq \mathbb{A}^\star\}$$

is τ_{Z}-open (and so τ-open) and a neighbourhood of t.
We exploit τ-dispersivity and genericity of $\mathbb{A}^\star$ to find:
$u \in \mathcal{U}$ such that $\mathrm{Im}(u) \subseteq \mathbb{A}^\star$. (CONTRADICTION!)

□

[‡]A **lovely pair** in model theory ▸ More on this.

Proof idea of the main theorem

Proof idea.

Suppose $\tau \subsetneq \tau_{\mathrm{pw}}$ is a Hausdorff semigroup topology. Consider

$$\Delta := \{a \in \Omega | \text{ every } s \in \overline{G} \text{ is } \tau\text{-rigid on } a\} .$$

By previous Lemma, $\Delta \subsetneq \Omega$.
We can find $t \in \overline{G}$ satisfying a strong form of τ-dispersivity on $\Omega \setminus \Delta$.
By **modularity**, find a "generic" copy $\mathbb{A}^\star$ of $\mathbb{A}$ within itself.[‡]
In particular,

$$\mathcal{U} := \{s \in \overline{G} | \ \mathrm{Im}(s) \subsetneq \mathbb{A}^\star\}$$

is τ_{Z}-open (and so τ-open) and a neighbourhood of t.
We exploit τ-dispersivity and genericity of $\mathbb{A}^\star$ to find:
$u \in \mathcal{U}$ such that $\mathrm{Im}(u) \subseteq \mathbb{A}^\star$. (CONTRADICTION!)

□

[‡]A **lovely pair** in model theory ▸ More on this.

Thank you!

For $\mathbb{A}$ ω-categorical

- We show τ_{Z} is not Hausdorff on $\mathrm{EEmb}(\mathbb{A})$ whenever $\mathrm{Aut}(\mathbb{A})$ has non-trivial centre;
- We find new ways to prove minimality of τ_{pw} as long as acl is sufficiently tame (i.e., modular).

Survey:

Paper:

Bibliography I

- Ahlbrandt, Gisela and Martin Ziegler (1986). "Quasi-finitely axiomatizable totally categorical theories". In: *Annals of Pure and Applied Logic* 30.1, pp. 63–82.
- Ben Yaacov, Itaï and Todor Tsankov (2016). "Weakly almost periodic functions, model-theoretic stability, and minimality of topological groups". In: *Transactions of the American Mathematical Society* 368.11, pp. 8267–8294.
- Ben-Yaacov, Itay, Anand Pillay, and Evgueni Vassiliev (2003). "Lovely pairs of models". In: *Annals of pure and applied logic* 122.1-3, pp. 235–261.
- Berenstein, Alexander and Evgueni Vassiliev (2010). "On lovely pairs of geometric structures". In: *Annals of pure and applied logic* 161.7, pp. 866–878.

Bibliography II

- Chang, Xiao and Paul Gartside (2017). "Minimum topological group topologies". In: *Journal of Pure and Applied Algebra* 221.8, pp. 2010–2024.
- Dikranjan, Dikran and Michael Megrelishvili (2013). "Minimality conditions in topological groups". In: *Recent progress in general topology III*. Springer, pp. 229–327.
- Elliott, L., Julius Jonušas, Zachary Mesyan, James Mitchell, Michał Morayne, and Yann Peresse (2023). "Automatic continuity, unique Polish topologies, and Zariski topologies on monoids and clones". In: *Transactions of the American Mathematical Society* 376.11, pp. 8023–8093.
- Elliott, L., Julius Jonušas, James Mitchell, Yann Peresse, and Michael Pinsker (2023). "Polish topologies on endomorphism monoids of relational structures". In: *Advances in Mathematics* 431, p. 109214.

Bibliography III

- Gaughan, Edward D (1967). "Topological group structures of infinite symmetric groups". In: *Proceedings of the National Academy of Sciences* 58.3, pp. 907–910.
- Ghadernezhad, Zaniar and Javier de la Nuez González (2024). "Group topologies on automorphism groups of homogeneous structures". In: *Pacific Journal of Mathematics* 327.1, pp. 83–105.
- Lascar, Daniel (1991). "Autour de la propriété du petit indice". In: *Proceedings of the London Mathematical Society* 62.1, pp. 25–53.
- Marimon, Paolo and Michael Pinsker (2025). *A guide to topological reconstruction on endomorphism monoids and polymorphism clones*. arXiv: 2512.01086 [math.LO]. URL: https://arxiv.org/abs/2512.01086.

Bibliography IV

- Nuez Gonzalez, J. de la, Zaniar Ghadernezhad, Paolo Marimon, and Michael Pinsker (2026). *Minimal and intrinsic topologies on monoids of elementary embeddings*. arXiv: 2603.28419 [math.LO]. URL: https://arxiv.org/abs/2603.28419.
- Pinsker, Michael and Clemens Schindler (2023). "On the Zariski topology on endomorphism monoids of omega-categorical structures". In: *The Journal of Symbolic Logic*, pp. 1–19.
- Poizat, Bruno (1983). "Paires de structures stables". In: *The Journal of Symbolic Logic* 48.2, pp. 239–249.

Bi-interpretations

Definition

An **interpretation** of $\mathbb{B}$ in $\mathbb{A}$ is a partial surjection $I : A^d \to B$ for some $d \in \mathbb{N}$ such that for each relation R of $\mathbb{B}$ defined by an atomic formula, $I^{-1}(R)$ is definable in $\mathbb{A}$ (without parameters).

Bi-interpretations

Definition

An **interpretation** of $\mathbb{B}$ in $\mathbb{A}$ is a partial surjection $I : A^d \to B$ for some $d \in \mathbb{N}$ such that for each relation R of $\mathbb{B}$ defined by an atomic formula, $I^{-1}(R)$ is definable in $\mathbb{A}$ (without parameters).

Example

$(\mathbb{C}; 0; 1; \times; +)$ is interpretable in $(\mathbb{R}; 0; 1; \times; +)$ (by the Argand plane).

Bi-interpretations

Definition

An **interpretation** of $\mathbb{B}$ in $\mathbb{A}$ is a partial surjection $I : A^d \to B$ for some $d \in \mathbb{N}$ such that for each relation R of $\mathbb{B}$ defined by an atomic formula, $I^{-1}(R)$ is definable in $\mathbb{A}$ (without parameters).

Interpretations compose naturally.
If I is an interpretation of $\mathbb{B}$ in $\mathbb{A}$ and J is an interpretation if $\mathbb{A}$ in $\mathbb{B}$, $\mathbb{A}$ and $\mathbb{B}$ are **bi-interpretable** if $I \circ J$ and $J \circ I$ are definable in $\mathbb{B}$ and $\mathbb{A}$ respectively.

Bi-interpretations

Definition

An **interpretation** of $\mathbb{B}$ in $\mathbb{A}$ is a partial surjection $I : A^d \to B$ for some $d \in \mathbb{N}$ such that for each relation R of $\mathbb{B}$ defined by an atomic formula, $I^{-1}(R)$ is definable in $\mathbb{A}$ (without parameters).

Interpretations compose naturally.
If I is an interpretation of $\mathbb{B}$ in $\mathbb{A}$ and J is an interpretation if $\mathbb{A}$ in $\mathbb{B}$, $\mathbb{A}$ and $\mathbb{B}$ are **bi-interpretable** if $I \circ J$ and $J \circ I$ are definable in $\mathbb{B}$ and $\mathbb{A}$ respectively.

Example

$(\mathbb{N}; =)$ is bi-interpretable with the **Johnson graph** $\mathfrak{J}(2)$:

- Domain: $[\mathbb{N}]^2$ (unordered pairs);
- Edges $E := \{(\{a, b\}, \{c, d\}) \in ([\mathbb{N}]^2)^2 |\ |\{a, b\} \cap \{c, d\}| = 1\}$.

Bi-interpretations

Definition

An **interpretation** of $\mathbb{B}$ in $\mathbb{A}$ is a partial surjection $I : A^d \to B$ for some $d \in \mathbb{N}$ such that for each relation R of $\mathbb{B}$ defined by an atomic formula, $I^{-1}(R)$ is definable in $\mathbb{A}$ (without parameters).

Interpretations compose naturally.
If I is an interpretation of $\mathbb{B}$ in $\mathbb{A}$ and J is an interpretation if $\mathbb{A}$ in $\mathbb{B}$, $\mathbb{A}$ and $\mathbb{B}$ are **bi-interpretable** if $I \circ J$ and $J \circ I$ are definable in $\mathbb{B}$ and $\mathbb{A}$ respectively.

Theorem (Coquand in Ahlbrandt and Ziegler 1986, cf. Lascar 1991)

Let $\mathbb{A}$ and $\mathbb{B}$ be ω-categorical structures. Then, TFAE:

- *$\mathbb{A}$ and $\mathbb{B}$ are bi-interpretable;*
- *$\mathrm{Aut}(\mathbb{A})$ and $\mathrm{Aut}(\mathbb{B})$ are isomorphic as topological groups;*
- *$\mathrm{EEmb}(\mathbb{A})$ and $\mathrm{EEmb}(\mathbb{B})$ are isomorphic as topological monoids.*

Bi-interpretations

Definition

An **interpretation** of $\mathbb{B}$ in $\mathbb{A}$ is a partial surjection $I : A^d \to B$ for some $d \in \mathbb{N}$ such that for each relation R of $\mathbb{B}$ defined by an atomic formula, $I^{-1}(R)$ is definable in $\mathbb{A}$ (without parameters).

Interpretations compose naturally.
If I is an interpretation of $\mathbb{B}$ in $\mathbb{A}$ and J is an interpretation if $\mathbb{A}$ in $\mathbb{B}$, $\mathbb{A}$ and $\mathbb{B}$ are **bi-interpretable** if $I \circ J$ and $J \circ I$ are definable in $\mathbb{B}$ and $\mathbb{A}$ respectively.

Theorem (Coquand in Ahlbrandt and Ziegler 1986, cf. Lascar 1991)

Let $\mathbb{A}$ and $\mathbb{B}$ be ω-categorical structures. Then, TFAE:

- *$\mathbb{A}$ and $\mathbb{B}$ are bi-interpretable;*
- *$\mathrm{Aut}(\mathbb{A})$ and $\mathrm{Aut}(\mathbb{B})$ are isomorphic as topological groups;*
- *$\mathrm{EEmb}(\mathbb{A})$ and $\mathrm{EEmb}(\mathbb{B})$ are isomorphic as topological monoids.*

Lovely pairs

Definition (Lovely pair[§])

$\mathbb{A}$ ω-categorical. $L \subseteq \mathbb{A}$ algebraically closed is a **lovely pair** if

- (coheir) for finite algebraically closed A, B with $\operatorname{acl}(B) \cap \operatorname{acl}((L \cap B) \cup A) = \operatorname{acl}(L \cap B)$, there is $g \in G_B$ such that $g(A) \subseteq L$;
- (extension) for finite algebraically closed A, B there is $g \in G_B$ with $\operatorname{acl}(g(A) \cup B) \cap L = B$.

[§]Berenstein and Vassiliev 2010 (cf. Ben-Yaacov, Pillay, and Vassiliev 2003; Poizat 1983)

Lovely pairs

Definition (Lovely pair[§])

$\mathbb{A}$ ω-categorical. $L \subseteq \mathbb{A}$ algebraically closed is a **lovely pair** if

- (coheir) for finite algebraically closed A, B with $\operatorname{acl}(B) \cap \operatorname{acl}((L \cap B) \cup A) = \operatorname{acl}(L \cap B)$, there is $g \in G_B$ such that $g(A) \subseteq L$;
- (extension) for finite algebraically closed A, B there is $g \in G_B$ with $\operatorname{acl}(g(A) \cup B) \cap L = B$.

Examples

- in $(\mathbb{N}; =)$ this is an infinite-coinfinite set;
- in $(\mathbb{Q}; <)$ this is a dense codense set;
- in vector space $\mathbb{F}_q^{\aleph_0}$ this is an infinite-dimensional subspace of infinite codimension.

[§]Berenstein and Vassiliev 2010 (cf. Ben-Yaacov, Pillay, and Vassiliev 2003; Poizat 1983)

What is left out?

- Droste's almost homogeneous semilinear orders
  (but for these we can prove $\tau_{\mathrm{pw}} = \tau_{\mathrm{Z}}$);
- Hrushovski constructions
  (but these seem to be amenable to a variant of our proof);
- atomless Boolean algebras
  (τ_{pw} should be minimal by other methods).