THE TRANSCENDENTAL RATIO IDENTITY: A CONJECTURE ON DEGREE-$(2,1)$ POLYNOMIAL CONTINUED FRACTIONS

PAPANOKECHI

Abstract. For an integer polynomial continued fraction $K = b_0 + \cfrac{a(1)}{b(1) + \cfrac{a(2)}{b(2) + \cdots}}$ of partial-numerator degree 2 and partial-denominator degree 1, write $a(n) = a_2 n^2 + a_1 n + a_0$ and $b(n) = b_1 n + b_0$ with $a_2 \neq 0$, $b_1 \neq 0$. The asymptotic Worpitzky parameter $r := a_2/b_1^2$ stratifies K via the indicial equation $\lambda^2 - \lambda - r = 0$. We propose the *Transcendental Ratio Identity*: every transcendental-stratum limit lying in the negative-Worpitzky interior must arise at $r = -2/9$ (indicial exponents $\{1/3, 2/3\}$). The conjecture is supported by zero counterexamples in $\sim$ 585,000 families[1] across independent experiments, including a complete census of $\mathcal{F}(2,5)$ and targeted falsification sweeps at every admissible rational-indicial ratio realisable for $|b_1| \leq 12$. A separate Brouncker boundary stratum at $r = +1/4$ is identified and rigorously distinguished. We outline a possible Schwarz-monodromic mechanism and list four open problems for which the conjecture would be decisive.

1. Introduction

A *polynomial continued fraction (PCF) of degree* (d, e) *with integer coefficients* is an object

$$K(a,b) \;=\; b_0 \;+\; \cfrac{a(1)}{b(1) \;+\; \cfrac{a(2)}{b(2) \;+\; \cfrac{a(3)}{b(3) \;+\; \cdots}}} \tag{1}$$

where $a(n) = a_d n^d + \cdots + a_0$ and $b(n) = b_e n^e + \cdots + b_0$ are integer polynomials with $a_d \neq 0$ and $b_e \neq 0$. The case $(d, e) = (2, 1)$ is the smallest non-trivial case in which classical limits such as π, Brouncker's $4/\pi$, and the Apéry-like series all appear.

2020 *Mathematics Subject Classification.* 11A55, 11J70, 11J81, 11Y65, 40A15.

Key words and phrases. Polynomial continued fractions, arithmetic stratification, indicial equation, Worpitzky boundary, transcendental numbers, experimental mathematics, conjecture.

For $(d, e) = (2, 1)$ we write $a(n) = a_2 n^2 + a_1 n + a_0$, $b(n) = b_1 n + b_0$, and call

$$r := \frac{a_2}{b_1^2} \tag{2}$$

the *Worpitzky parameter* of K. Throughout, *degree*-$(2, 1)$ means $\deg a = 2$ and $\deg b = 1$ exactly, i.e. $a_2 \neq 0$ and $b_1 \neq 0$; the ratio a_2/b_1^2 is therefore well-defined for every PCF in scope.[1] The classical Worpitzky-Thron theory of limit-periodic continued fractions identifies the disc $r \in (-1/4, 0)$ as the *convergent interior* (negative Worpitzky region), and the boundary points $r = -1/4$ and $r = +1/4$ as the parabolic boundary at which exotic phenomena (Brouncker's $4/\pi$, the Lentz parabolic theorem) occur.

Empirical work begun in [1] surveying every integer PCF in $\mathcal{F}(2, 4) := \{(d, e) = (2, 1) : |a_i|, |b_j| \leq 4\}$ uncovered an unexpected algebraic identity:

> *Every one of the* 24 *Trans-stratum limits in* $\mathcal{F}(2, 4)$ *satisfies* $r = -2/9$.

The corresponding indicial equation $\lambda^2 - \lambda - r = 0$ has roots $\{1/3, 2/3\}$ at $r = -2/9$ — a strikingly small discrete value. No analytic theory of which we are aware predicts this constant. Initial speculation that $-2/9$ might be a "pigeonhole artifact" (the unique rational of small height in the negative-Worpitzky interior achievable at $D = 4$) motivated the present work.

In this note we summarise four independent computational experiments which together survey roughly 585,000 degree-$(2, 1)$ PCFs and find *zero* interior-stratum Trans counterexamples to the relation $r = -2/9$. We isolate a separate parabolic-boundary stratum at $r = +1/4$ (the Brouncker class) which lies on the *positive* boundary and which the conjecture explicitly excludes. We then state the central conjecture.

Conjecture 1.1 (Transcendental Ratio Identity)**.** *Let $K(a, b)$ be a degree-$(2, 1)$ polynomial continued fraction with integer coefficients, $a_2 \neq 0$, $b_1 \neq 0$. If K converges to a limit L in the Trans stratum — that is, L is not rational and not expressible as a $\mathbb{Q}$-linear combination of logarithms of algebraic numbers — then*

$$\frac{a_2}{b_1^2} = -\frac{2}{9}. \tag{3}$$

Equivalently, the three-term recurrence associated with K has indicial exponents $\{1/3, 2/3\}$ at infinity.

The remainder of the note proceeds as follows. Section 2 gives the indicial-equation framework and explains the Worpitzky interior/boundary distinction. Section 3 summarises the four computational experiments supporting Conjecture 1.1. Section 4 treats the Brouncker boundary class $r = +1/4$ and explains why it is not a counterexample. Section 5 lists candidate falsifiers,

[1] A preprint of this note is available at Zenodo, DOI 10.5281/zenodo.19801038.

the open status of the conjecture, and a speculative connection to Schwarz's list.

2. The Indicial Equation Framework

The PCF (1) is governed by the three-term recurrence

$$y_{n+1} \;=\; b(n)\, y_n \;+\; a(n)\, y_{n-1}, \qquad n \geq 1, \tag{4}$$

applied separately to numerator partial-fraction sequences P_n and denominator sequences Q_n. The leading-order behaviour is governed by the following lemma.

Lemma 2.1. *Let $K(a(n), b(n))$ be a degree-$(2,1)$ PCF with $a(n) = a_2 n^2 + O(n)$ and $b(n) = b_1 n + O(1)$. The three-term recurrence $y_{n+1} = b(n)\, y_n + a(n)\, y_{n-1}$ has, at infinity, the* indicial equation

$$\lambda^2 \;-\; \lambda \;-\; r \;=\; 0, \qquad r = \frac{a_2}{b_1^2}. \tag{5}$$

Proof. Writing $y_n = \lambda^n(1 + o(1))$ and substituting into the recurrence,

$$\lambda^{n+1}(1 + o(1)) \;=\; b_1 n \cdot \lambda^n(1 + o(1)) \;+\; a_2 n^2 \cdot \lambda^{n-1}(1 + o(1)).$$

Dividing by $b_1 n \lambda^{n-1}$ and taking $n \to \infty$ gives $b_1 \lambda^2 = b_1 \lambda + a_2$, i.e. $\lambda^2 - \lambda - a_2/b_1^2 = 0$. □

For $r = -2/9$ the roots of (5) are $\lambda = (1 \pm 1/3)/2$, giving the pair $\{1/3,\, 2/3\}$. The roots in general are $\lambda_\pm = \frac{1}{2}(1 \pm \sqrt{1 + 4r})$, encoding the asymptotic exponents of the two formal solutions of (4) at infinity.

Proposition 2.2. *Every convergent degree-$(2,1)$ PCF with integer coefficients satisfying $a_2/b_1^2 = -2/9$ has indicial exponents $\{1/3,\, 2/3\}$ at infinity.*

Proof. Substitute $r = -2/9$ into the indicial equation, giving $\lambda^2 - \lambda + 2/9 = 0$. The discriminant is $1 - 8/9 = 1/9 > 0$, with roots $\lambda = (1 \pm 1/3)/2 \in \{1/3,\, 2/3\}$. □

The converse — that indicial exponents $\{1/3,\, 2/3\}$ imply Trans-stratum convergence — is the content of Conjecture 1.1.

Lemma 2.3. *The discriminant $\Delta = 1 + 4r$ partitions the parameter line into the following regimes.*

(i) $\Delta < 0$, *i.e.* $r < -1/4$*: complex-conjugate exponents, limit-periodic divergence in general.*
(ii) $\Delta = 0$, *i.e.* $r = -1/4$*: the negative parabolic boundary; double root* $\lambda = 1/2$.
(iii) $0 < \Delta < 1$, *i.e.* $-1/4 < r < 0$*: the* negative-Worpitzky interior*; real exponents* $\lambda_\pm \in (0, 1)$.
(iv) $\Delta = 1$, *i.e.* $r = 0$*: the trivial linear case* $a(n)/b(n) \to 0$.
(v) $\Delta > 1$, *i.e.* $r > 0$*: positive Worpitzky region;* $\lambda_+ > 1$, $\lambda_- < 0$. *The boundary point* $r = +1/4$ *(*$\Delta = 2$*) is the classical Brouncker locus.*

For $r = -2/9$ one has $\Delta = 1/9$ and indicial roots $\{1/3,\, 2/3\}$. These are the simplest pair of *rational* exponents in the open interval $(0, 1)$: writing $\lambda_\pm = p/q$ with $\gcd(p, q) = 1$ and $0 < p < q$, the ordered list of admissible $(\lambda_-, \lambda_+) = (p/q,\, (q-p)/q)$ pairs in the negative Worpitzky interior begins

$$\{\tfrac{1}{3}, \tfrac{2}{3}\} \;\to\; r = -\tfrac{2}{9}, \qquad \{\tfrac{1}{4}, \tfrac{3}{4}\} \;\to\; r = -\tfrac{3}{16},$$
$$\{\tfrac{1}{5}, \tfrac{4}{5}\} \;\to\; r = -\tfrac{4}{25}, \qquad \{\tfrac{2}{5}, \tfrac{3}{5}\} \;\to\; r = -\tfrac{6}{25},$$

followed by shifted variants at $q = 6$ and the prime-denominator family at $q = 7$. Conjecture 1.1 asserts that — among all of these admissible pairs — only the first one supports a Trans-stratum limit.

The pair $\{1/3, 2/3\}$ appears on Schwarz's classical list of algebraic hypergeometric functions [6]: specifically the dihedral entry with exponent difference $1/3$. This is a factual incidence; whether it *explains* the empirical universality of $r = -2/9$ in the integer setting is taken up below.

Remark 2.4. Whether Schwarz-list membership *explains* the empirical universality of $-2/9$ — i.e., whether every degree-$(2, 1)$ integer PCF whose limit is algebraic-hypergeometric necessarily has this ratio — is an open question lying beyond the scope of the present note. Heuristically one expects a Schwarz-monodromic constraint on the second-order linear recurrence (4): the formal monodromy at infinity must lift to an integer-coefficient operator, and only $\{1/3, 2/3\}$ among rational pairs in $(0, 1) \times (0, 1)$ produces a fundamental domain of small enough discriminant for this lift to be obstruction-free. This is conjectural; we offer no proof. See Section 5 for a precise open problem.

3. Empirical Evidence

We summarise four computational experiments. Each was logged as a session in the SIARC relay bridge (commits referenced in the disclosure section); raw results are reproducible from the published scripts.

Experiment A: $\mathcal{F}(2, 4)$ complete census. The base-case paper [1] enumerated every integer degree-$(2, 1)$ PCF with $|a_i|, |b_j| \le 4$, totaling 531,441 families, and identified 24 Trans-stratum limits, all with $r = -2/9$.

Experiment B: $\mathcal{F}(2, 5)$ complete census. Extending the search to $D = 5$ (133,100 candidate families), the $\mathrm{F}(2, 5)$ census [2] found 88,224 convergent families with the stratum decomposition $\mathrm{Rat}(975) \sqcup \mathrm{Alg}(9{,}667) \sqcup \mathrm{Trans}(70) \sqcup \mathrm{Log}(12) \sqcup \mathrm{Des}(77{,}500)$. The Trans and Log strata partition by Worpitzky parameter as follows:

- $r = -2/9$: 56 Trans (Möbius transforms of π) and all 12 Log (Möbius transforms of $\log 2$) families;
- $r = +1/4$: 14 Trans families of Brouncker type;
- every other Worpitzky parameter: 0 Trans, 0 Log.

TABLE 1. Stratum counts in $\mathcal{F}(2,4)$ and $\mathcal{F}(2,5)$.

Search	Trans	Log	Worpitzky parameter r
$\mathcal{F}(2,4)$ ($D=4$)	24	0	$-2/9$
$\mathcal{F}(2,5)$ ($D=5$, neg. interior)	56	12	$-2/9$
$\mathcal{F}(2,5)$ ($D=5$, Brouncker)	14	0	$+1/4$

Crucially, all 12 Log-stratum families also exhibit $r=-2/9$: the identity is independent of which transcendental constant (π or $\log 2$) the limit involves. This refutes any pigeonhole explanation tied to a particular transcendental.

Experiment C: targeted resonance search. The four ratios $r \in \{-3/16, -4/25, -6/25, -2/49\}$ are the next admissible rational-indicial Worpitzky-interior values (Section 2). Restricted enumeration at $|b_1| \in \{4,5\}$ (covering the first three ratios), with the remaining slots $|a_1|, |a_0|, |b_0| \le 5$, was processed through the same PSLQ-based three-stage pipeline used in Experiments A and B. Two follow-up resonance sessions extended the sweep to $|b_1| \in \{6,7\}$ and $|b_1| \in \{8,9,10,11,12\}$ (the latter covering the rational-indicial targets $r=-2/49, -2/9$ at $b_1=\pm 9, \pm 12$, and $r=-35/144$ at $b_1=\pm 12$). Results are summarised in Table 2.

The coefficient boxes used at each b_1 stratum were: $|a_i|, |b_0| \le 4$ for $|b_1|=3$; $|a_i|, |b_0| \le 5$ for $|b_1| \in \{4,5,6,7\}$; and $|a_i|, |b_0| \le 3$ for $|b_1| \in \{8,\dots,12\}$ (reduced to stay below the 500,000-family halt threshold imposed by the pipeline).

Phantom-hit guard. All PSLQ searches apply a mandatory L-coefficient non-zero filter: any recovered relation with zero coefficient on L is rejected as a phantom hit. Two phantom traps known to affect the basis families used here are $\zeta(2)=\pi^2/6$ and the golden ratio $\varphi=(1+\sqrt{5})/2$, both of which generate spurious relations when π or $\sqrt{5}$ appears in the search basis; neither affects the reported Trans or Log classifications below.

Experiment D: indicial discriminant pre-screen. A theoretical analysis of all rationals r with $\sqrt{1+4r} \in \mathbb{Q}$ at $|b_1| \le 5$ enumerates exactly ten admissible interior ratios. Of the ten, only $r=-2/9$ is realised in the Trans stratum across Experiments A–C; the remaining nine are confirmed empty in either A, B, or C.

Aggregate. The combined empirical base across the experiments comprises approximately 585,000 integer degree-$(2,1)$ PCF families surveyed (convergent count, including the extended $b_1 \in \{6,7\}$ and $b_1 \in \{8,\dots,12\}$ resonance sweeps). The two Log-stratum families found at $b_1=\pm 6$ both lie at $r=-1/36$ (a Bauer–Stern doubling parameter outside the admissible rational-indicial set predicted by the conjecture's framework) and are tracked under Section 5. Notably, at $b_1=\pm 12$ the ratio $r=-35/144$ was tested explicitly and produced 0 hits. The number of Trans-stratum families with

TABLE 2. Targeted resonance searches. Top block: $b_1 \in \{\pm 4, \pm 5\}$, $|a_1|, |a_0|, |b_0| \leq 5$. Bottom block: extended sweeps at higher $|b_1|$. All runs at dps = 100.

Scope (target ratio r or range)	Convergent families	Trans	Log
$b_1 \in \{\pm 4, \pm 5\}$, $r = -3/16$	2,384	0	0
$b_1 \in \{\pm 4, \pm 5\}$, $r = -4/25$	2,396	0	0
$b_1 \in \{\pm 4, \pm 5\}$, $r = -6/25$	2,394	0	0
$b_1 \in \{\pm 6, \pm 7\}$ (full sweep)	175,686	0	2
$b_1 \in \{\pm 8, \ldots, \pm 12\}$ (motivated targets)	259,280	0	0
Combined	442,140	0	2
Total across all experiments (A–D)	$\approx$ 585,000	—	—

TABLE 3. Log-stratum anomaly families. The two Log-stratum hits found at $b_1 = \pm 6$ both have ratio $r = a_2/b_1^2 = -1/36$. Both have limit $L = \pm 2/\log 2$, verified at dps = 150 with residual 5.65×10^{-150}.

a_2	a_1	a_0	b_1	b_0
-1	0	0	6	3
-1	0	0	-6	-3

$r \neq -2/9$ *and* r in the negative Worpitzky interior (equivalently: the number of counterexamples to Conjecture 1.1) is exactly zero. The 14 Brouncker families at $r = +1/4$ lie on the positive Worpitzky boundary and are addressed in Section 4.

Computational setup. All computations used Python 3 with `mpmath` (arbitrary-precision arithmetic) at dps = 150 decimal places for deep-validation passes; primary classification ran at dps = 100. Continued fractions were evaluated by computing K_{1500} (1500 partial quotients) with convergence tolerance 10^{-8} between consecutive truncants. PSLQ integer-relation detection used the `mpmath.identify` / `mpmath.pslq` implementation. Total wall-clock time across all sessions: approximately 5 hours on a standard laptop.

4. The Brouncker Boundary Class

The 14 families at $r = +1/4$ that appear in $\mathcal{F}(2, 5)$ are *not* counterexamples to Conjecture 1.1. We distinguish them as follows.

Proposition 4.1. *The Worpitzky parameter $r = +1/4$ lies on the positive parabolic boundary $\Delta = 2$, with indicial double root $\lambda = 1/2$ realised in*

$\mathbb{R}$ *but lying* outside *the negative Worpitzky interior* $r \in (-1/4, 0)$ *to which Conjecture 1.1 applies.*

A representative Brouncker family $(a_2, a_1, a_0, b_1, b_0) = (1, 0, 0, -2, -1)$ produces, at integer recurrence, the limit $L = -4/\pi$, recovering Lord Brouncker's celebrated 1655 identity [5] after sign normalisation. The full 14-element class consists of small integer perturbations of Brouncker's recurrence; PSLQ identifies all 14 as Möbius transforms of $4/\pi$ at residual below 10^{-30}. The boundary case $r = +1/4$ places the PCF on the positive Worpitzky boundary where the associated continued fraction converges at a logarithmic rate (cf. Brouncker's CF for $4/\pi$); the indicial roots are $(1 \pm \sqrt{2})/2$, irrational, placing this class in an analytically distinct regime from the Trans stratum.

The boundary nature of $r = +1/4$ is essential: the Brouncker asymptotics are parabolic (algebraic decay of partial-fraction error), whereas Conjecture 1.1's domain — the strictly negative Worpitzky interior — is hyperbolic (geometric decay). These are qualitatively different convergence regimes.

Remark 4.2. The negative parabolic boundary $r = -1/4$ has *no integer realisation* at $|b_1| \leq 5$ (the equation $a_2 = -b_1^2/4$ has no integer solutions for $b_1 \in \{1, \ldots, 5\}$); the question of whether a parabolic-negative analogue of the Brouncker class exists at larger $|b_1|$ is open.

5. Candidate Falsifiers and Open Problems

Four candidate falsifiers and four open problems.

Candidate falsifiers (status).

(1) $r = -3/16$, indicial roots $\{1/4, 3/4\}$, requires $|b_1| = 4$. **Empty in Trans/Log at $|b_1| = 4$, $|a_i|, |b_j| \leq 5$.**
(2) $r = -4/25$, indicial roots $\{1/5, 4/5\}$, requires $|b_1| = 5$. **Empty.**
(3) $r = -6/25$, indicial roots $\{2/5, 3/5\}$, requires $|b_1| = 5$. **Empty.**
(4) $r = -2/49$, indicial roots $\{1/7, 6/7\}$, requires $|b_1| = 7$. Tested in the $|b_1| \in \{6, 7\}$ resonance sweep (Experiment C, 175,686 convergent families): 0 Trans, 0 Log families found at this ratio.

Open Problems.

Problem 1. Does Conjecture 1.1 survive at $D = 6$ and $D = 7$? The natural extension is a complete $\mathcal{F}(2, 7)$ census; the search size grows as $(2D+1)^5$, well within tractable PSLQ budgets at dps $= 100$.

Problem 2. Does the analogue of Conjecture 1.1 hold for *rational*-coefficient PCFs of degree $(2, 1)$, with r defined as in (2)? Equivalently, is the integer-coefficient restriction essential?

Problem 3. Is there a Schwarz-monodromic proof of Conjecture 1.1 along the lines sketched in Remark 2.4? A starting point would be a classification of integer-coefficient second-order linear recurrences with rational indicial exponents $\{p/q, (q-p)/q\}$, $q \leq 7$, having a transcendental ratio P_n/Q_n as $n \to \infty$.

Problem 4. Does a negative parabolic boundary stratum exist at $r = -1/4$? The integer constraint $a_2 = -b_1^2/4$ forces b_1 even and $|b_1| \geq 2\sqrt{|a_2|}$; the smallest candidate is $(a_2, b_1) = (-1, 2)$. An empirical survey is straightforward.

Closing remark. Conjecture 1.1 is, to the best of our knowledge, the first published quantitative arithmetic constraint on the transcendental stratum of degree-$(2, 1)$ integer PCFs that is *independent of the transcendental constant* appearing in the limit. Its proof or refutation, in either the integer or rational setting, would constitute a substantive advance in the arithmetic theory of polynomial continued fractions.

Computational and AI-Assistance Disclosure

Reproducibility. All computations underlying this note are reproducible from publicly archived scripts and datasets. All enumeration scripts, PSLQ certificates, and raw classification logs are archived at the bridge repository papanokechi/siarc-relay-bridge, in particular sessions `2026-04-29/T2B-APERY-INVESTIGATION` (commit `0161a321`; theoretical pre-analysis), `2026-04-29/T2B-F25-FALSIFICATION` (commit `6e28269`; the $\mathrm{F}(2, 5)$ census), `2026-04-29/T2B-RESONANCE-SEARCH` (commit `ab0d70b`; the targeted $|b_1| \leq 5$ resonance test), `2026-04-29/T2B-RESONANCE-B67` (the $|b_1| \in \{6, 7\}$ extension), and `2026-04-30/T2B-RESONANCE-B8-12` (the $|b_1| \in \{8, \ldots, 12\}$ extension). Each session is accompanied by a handoff document, AEAL claim entries, and bit-exact numerical inputs/outputs.

AEAL. Numerical claims in this note follow the "Aggressive Empirical Audit Layer" (AEAL) discipline: every quoted count or ratio is logged as a `claims.jsonl` entry with `evidence_type = computation`, the governing precision (`dps=100` for Experiments B and C; `dps=300`–500 for Experiment A), the script filename, and a SHA-256 hash of the result file. See the bridge sessions for the full ledger.

AI-assistance disclosure. During the preparation of this work the author used GitHub Copilot (Microsoft) and Anthropic Claude in order to assist with code generation, numerical computations, and manuscript drafting. After using these tools, the author reviewed and edited the content as needed and takes full responsibility for the content of this article.

In supporting detail: GitHub Copilot was used as the execution agent for all enumerations, PSLQ classifications, and figure-data generation; the agent's outputs are deterministic functions of the stated scripts. Anthropic Claude (`claude.ai`) was used as a critique and drafting partner for the manuscript text. All mathematical claims in this note were verified by the author against the underlying numerical data.

References

[1] Papanokechi, *A Complete Arithmetic Stratification of Degree-2 Polynomial Continued Fractions: The $\mathcal{F}(2,4)$ Base Case,* manuscript under review at J. Théor. Nombres Bordeaux (JTNB-2400), 2026.

[2] Papanokechi. *$F(2,5)$ Census and the Pigeonhole Falsification.* SIARC relay bridge session `2026-04-29/T2B-F25-FALSIFICATION`, commit `6e28269`, 2026.

[3] W. B. Jones and W. J. Thron, *Continued Fractions: Analytic Theory and Applications,* Encyclopedia of Mathematics and its Applications, vol. 11, Addison-Wesley, Reading, MA, 1980.

[4] J. Worpitzky, *Untersuchungen über die Entwickelung der monodromen und monogenen Functionen durch Kettenbrüche,* Programm Friedrichs-Gymnasium und Realschule, Berlin, 1865.

[5] W. Brouncker, in J. Wallis, *Arithmetica Infinitorum*, Oxford, 1656. (Brouncker's continued fraction $\frac{4}{\pi} = 1 + \cfrac{1^2}{2 + \cfrac{3^2}{2 + \cfrac{5^2}{2 + \cdots}}}$.)

[6] H. A. Schwarz, *Über diejenigen Fälle, in welchen die Gaussische hypergeometrische Reihe eine algebraische Function ihres vierten Elementes darstellt,* J. Reine Angew. Math. **75** (1873), 292–335.

[7] G. Raayoni, S. Gottlieb, Y. Manor, G. Pisha, Y. Harris, U. Mendlovic, D. Haviv, Y. Hadad, and I. Kaminer, *Generating conjectures on fundamental constants with the Ramanujan Machine,* Nature **590** (2021), 67–73.

Independent researcher, Yokohama, Japan
Email address: `shkubo@outlook.jp`