COMPLEX MULTIPLICATION AS A TRANSCENDENCE PREDICATE FOR DEGREE-2 POLYNOMIAL CONTINUED FRACTIONS

PAPANOKECHI

Date: May 1, 2026.

2020 *Mathematics Subject Classification.* Primary 11J70, 11J81; Secondary 34M55, 11G15.

Key words and phrases. polynomial continued fractions, complex multiplication, Painlevé transcendents, Spec(K), balanced discriminant, Stokes phenomenon, experimental number theory.

ORCID: 0009-0000-6192-8273. **Version 1.3 (2026-05-01):** folds in the WKB exponent identity (Theorem 5.1, Sessions E and E′) and the channel-scope subsection for the V_quad prototype (Sessions D, E, E′); Conjecture A part (iv) is restated per the post-E′ decision matrix (Variant A: both $K = 12$ marginals, QL15 and QL26, confirmed as Padé–Borel artefacts). **Version 1.2 (2026-05-01):** added Session C (STOKES-FAMILY-EXTEND-3X) results for QL02, QL06, QL26; Stokes predicate now confirmed across all six known $\Delta < 0$ degree-2 PCF families. **Version 1.1 (2026-05-01):** added Session B results for family QL15 ($\Delta = -20$, non-Heegner). Version 1.0 published as DOI 10.5281/zenodo.19931636.

AI use disclosure. This research was conducted using the AEAL/SIARC multi-agent methodology. GitHub Copilot (powered by Anthropic Claude Opus 4.7) and Anthropic Claude were used to assist with code generation for numerical computations, cross-checking of intermediate results, and drafting of expository prose. Sessions D, E, and E′ of the AEAL chain were executed via Anthropic Claude and GitHub Copilot agents. All conjecture formulations, mathematical interpretations, evidence-class assignments, and final editorial decisions were made by the human author, who takes full responsibility for the content of this manuscript. The AI tools are not authors. Session-level computation logs and prompts are archived in the SIARC research workspace for reproducibility per the AEAL methodology of [5].

Abstract. We propose a transcendence predicate for degree-two polynomial continued fractions (PCFs) based on the sign of the balanced discriminant $\Delta = \beta^2 - 4\alpha\gamma$ of the denominator polynomial $b_n = \alpha n^2 + \beta n + \gamma$. Working inside the $\mathrm{Spec}(K)$ classification framework of Papanokechi (2026), we present a v5 upgrade of the schema that fixes two cross-paper inconsistencies and adds the Stokes exponent and connection-coefficient proxy as components 6 and 7. Our central empirical finding is a sharp dichotomy across 30 degree-two families: 24 $\mathrm{F}(2,4)$ Trans-stratum families have $\Delta \in \{+1, +8\}$ and admit elementary closed forms, whereas 6 candidates with $\Delta \in \{-3, -4, -7, -11, -20, -28\}$ all return PSLQ non-detection against an 18-element constants basis at working precision 220 digits and integer bound 10^{10}. The V_quad family ($\Delta = -11$) is the explicit Painlevé $\mathrm{III}(D_6)$ prototype with parameters $\alpha = 1/6$, $\beta = \gamma = 0$, $\delta = -1/2$. For QL01 ($\Delta = -3$) the Eisenstein-root deformation yields a Stokes exponent $S \in \{0.770, 0.819\} < 1$, a fractional-power singularity that is the qualitative signature of a Painlevé/Stokes obstruction. Conjecture A v5 formalises the dichotomy. Version 1.3 folds in the WKB exponent identity (Theorem 5.1) and the revised Conjecture A part (iv) per Sessions D, E, and E′. The session chain that produced these results is itself an instance of the AEAL four-class evidence gate methodology.

Series label: P12 / PCF-1. This paper is P12 in the AI Epistemic Governance Stack (Zenodo preprints P7–P11, DOIs 10.5281/zenodo.19562751, 10.5281/zenodo.19565086, 10.5281/zenodo.19591564, 10.5281/zenodo.19591685, 10.5281/zenodo.19717115) and simultaneously PCF-1 in a mathematical companion track documenting the PCF research program.

1. Introduction

A central question in the analytic theory of polynomial continued fractions (PCFs) is: *what determines whether the limit $L = K(a_n, b_n)$ is elementary or transcendental?* The Ramanujan-Machine search programme has produced a rapidly growing list of empirically identified PCFs with closed forms [13], but no spectral invariant has been proposed that *predicts* which families escape elementary identification. The present paper addresses that gap for the degree-two regime $a_n \in \mathbb{Z}[n]_{\leq 1}$, $b_n \in \mathbb{Z}[n]_{\leq 2}$, and proposes the sign of the balanced discriminant $\Delta = \beta^2 - 4\alpha\gamma$ as the controlling invariant.

We work inside the $\mathrm{Spec}(K)$ classification framework introduced in [1], which assigns to each PCF the invariant

$$\mathrm{Spec}(K) = (d, \Lambda, \Delta, \rho, \tau).$$

The first contribution of this paper is a v5 upgrade of the schema: we resolve two cross-paper inconsistencies (a clash between the $d = 1$ and $d = 2$ growth-degree conventions of [3] and [2], and a vocabulary mismatch between Schwarz-triangle and Takeuchi terminology in the arithmetic sector τ), give an explicit numbered definition of Δ that was previously only recoverable from the Catalan example, and append a sixth and seventh component (Stokes exponent S and connection-coefficient proxy σ) which we use below to detect non-analytic transcendental obstructions even when no specific Painlevé ansatz fits.

The second contribution is empirical and is the core mathematical content. We exhibit a sharp dichotomy across 30 degree-two families:

- All 24 F$(2, 4)$ Trans-stratum families have $\Delta \in \{+1, +8\} > 0$; their normalized Poincaré roots Λ lie in $\mathbb{Q}$ or $\mathbb{Q}(\sqrt{2})$, and their limits admit explicit closed forms via Lindemann–Weierstrass.
- All 6 tested $\Delta < 0$ families ($\Delta \in \{-3, -4, -7, -11, -20, -28\}$) return PSLQ non-detection against an 18-element basis at 220 digits with integer bound 10^{10}, and have Λ in $\mathbb{Q}(\sqrt{\Delta})$ by construction.

The first row anchors the elementary side; the second row anchors the transcendental side. Zero counterexamples have been found across the sample. Notably, one of the new $\Delta < 0$ candidates (QL15, $\Delta = -20$) has class number $h = 2$ and is therefore non-Heegner, yet still exhibits PSLQ non-detection — which suggests the controlling invariant is the sign of Δ rather than the class number.

The third contribution is a structural-evidence layer for the analytic side of the conjecture. For the prototype V_quad family ($\Delta = -11$) the limit is established as a Painlevé III(D_6) transcendent with explicit parameters [2]. For the simplest CM companion QL01 ($\Delta = -3$, recurrence $a_n = n$, $b_n = n^2 - n + 1$) we report a 3×3 residual matrix (three CM-respecting deformations $\times$ three Painlevé ansätze) with best validation residual 0.00840 (D2 + Painlevé III), an 18.7$\times$ improvement over the universal $b_n \to b_n + tn^2$ baseline. No specific Painlevé ansatz fits below the 10^{-20} threshold; nonetheless the Eisenstein-root deformation D3 yields a Stokes exponent $S \in \{0.770, 0.819\} < 1$. This sub-linear decay of $|L(t) - L(0)|$ near $t = 0$ is precisely the qualitative signature of a fractional-power Painlevé/Stokes singularity, and it is detectable even when the explicit ansatz fits all fail. The structural evidence therefore supports a weakened version of the analytic conjecture in which the specific Painlevé type is allowed to be family-dependent.

The paper is organised as follows. Section 2 states the v5 upgrade of Spec(K), including the explicit definition of the balanced discriminant. Section 3 presents the full 30-family evidence table and states parts (i)–(ii) of Conjecture A v5. Section 4 reports the QL01 deformation residual matrix and the Stokes-exponent diagnostic, and states parts (iii)–(iv). Section 6 is the V_quad explicit prototype. Section 7 maps the session chain producing this paper to the AEAL four evidence classes. Section 8 records open questions.

This paper is positioned as the mathematical instantiation of the AEAL evidence chain documented in P7–P11. The session chain PSLQ $\to \Delta$ computation $\to$ Painlevé fitting $\to$ Stokes diagnostics is itself an instance of the AEAL four-class evidence-gate methodology applied to a previously open question of analytic number theory: every numerical claim in this paper is independently reproduced across at least two distinct sessions, every conjecture is supported by explicit empirical or algebraic evidence, and every gap (e.g. the absence of a universal Painlevé type for $\Delta < 0$ families) is flagged rather than glossed.

2. The Spec(K) Framework — v5 Upgrade

We work with PCFs of the form

$$K(a_n, b_n) = b_0 + \cfrac{a_1}{b_1 + \cfrac{a_2}{b_2 + \cdots}},$$

together with the Wallis convergent recurrence $p_n = b_n p_{n-1} + a_n p_{n-2}$ and seeds $(p_{-1}, p_0, q_{-1}, q_0) = (1, b_0, 0, 1)$.

The Spec(K) invariant of [1] is

$$\mathrm{Spec}(K) = (d, \Lambda, \Delta, \rho, \tau),$$

with d the dominant growth degree, Λ the normalized Perron–Poincaré root, Δ the balanced discriminant, ρ the Wallis tail-decay rate, and τ the arithmetic sector. The framework was proposed in [1] together with the Poincaré characteristic lemma, which in the balanced degree-one class ($a_n = \delta n^2 + O(n)$, $b_n = \alpha n + O(1)$) forces Λ to satisfy the quadratic

$$\Lambda^2 - \alpha\Lambda - \delta = 0. \tag{1}$$

Two cross-paper inconsistencies, resolved in v5. A coherence audit across [1], [3], and [2] surfaced two gaps:

(I-1) The growth-degree convention d is degree-one in [3] (where a_n is quadratic and b_n linear) but degree-two in [2] (where a_n is linear and b_n quadratic). v5 explicitly distinguishes these via

the *class label* "$d = k$ balanced" meaning the leading-degree pair $(\deg a, \deg b)$ is $(k+1, k)$ in the δ-direction and $(k-1, k)$ in the α-direction.

(I-2) The arithmetic-sector vocabulary τ in [1] mixes Schwarz-triangle and Takeuchi terminology in a way that does not consistently distinguish monodromy type from arithmetic field. v5 splits τ into τ_{mono} (monodromy type) and τ_{arith} (arithmetic field).

The balanced discriminant. The discriminant Δ was used throughout [1] without an explicit numbered definition; it was recovered in [3] from the Catalan example ($a_n = -4n^2$, $b_n = 4n+1$, leading to $\Lambda = 2$ as a repeated root, hence $\Delta = 0$). v5 promotes this to a definition.

Definition 2.1 (Balanced discriminant)**.** For a balanced PCF in the d=1 class with leading data (α, δ) as in (1),

$$\Delta \;=\; \alpha^2 + 4\delta.$$

For a balanced PCF in the d=2 class with $b_n = \alpha n^2 + \beta n + \gamma$ and $a_n = \delta n + \varepsilon$,

$$\Delta \;=\; \beta^2 - 4\alpha\gamma \;=\; \operatorname{disc}(b).$$

In both cases $\Delta < 0$ is called *the CM (complex multiplication) case*; the splitting field of the characteristic polynomial is then the imaginary-quadratic field $\mathbb{Q}(\sqrt{\Delta})$.

v5 additions: Stokes exponent and connection proxy. Given a one-parameter deformation $K(t)$ of K with limit $L(t)$ satisfying $L(0) = L$, define

$$S(t) \;=\; \frac{\log|L(t) - L(0)|}{\log|t|}, \tag{2}$$

$$\sigma(t) \;=\; \frac{L(t) - L(-t)}{2t\,L'(0)}. \tag{3}$$

The exponent S records the local power-law behaviour of $|L(t) - L(0)|$; analytic dependence forces $S \to 1$ as $t \to 0$. A fractional limit $\lim_{t\to 0} S(t) \in (0,1)$ signals a non-analytic, fractional-power singularity at $t = 0$, which is the qualitative signature of a Painlevé/Stokes connection. The proxy σ measures the antisymmetric deviation from the linear Taylor approximation; analytic dependence forces $\sigma \to 1$. We add S and σ as $\operatorname{Spec}(K)$ components 6 and 7.

3. The Sharp Dichotomy

We tested 30 degree-two PCF families: 24 from the $\mathrm{F}(2,4)$ Trans-stratum table of [3] and 6 from the curated $\Delta < 0$ database underlying [2, 14]. Every limit was independently recomputed from scratch via

the forward Wallis recurrence with seeds $(p_{-1}, p_0, q_{-1}, q_0) = (1, b_0, 0, 1)$ at working precision 320 digits, depth 2500, and verified to agree with the source database to at least 28 digits. The full evidence is summarised in Table 1.

Family	Δ	Λ field	Limit type	PSLQ status	Stokes signal
F(2, 4) Trans (Main) $\times 22$	$+1$	$\mathbb{Q}$	elementary	relation found	—
F(2, 4) Trans (Outlier) $\times 2$	$+8$	$\mathbb{Q}(\sqrt{2})$	elementary	relation found	—
QL01	-3	$\mathbb{Q}(\sqrt{-3})$	transcendental	no relation	$S \in [0.77, 0.82]$ ✓
QL02	-4	$\mathbb{Q}(i)$	transcendental	no relation	$S \in [0.56, 0.93]$ ✓
QL06	-7	$\mathbb{Q}(\sqrt{-7})$	transcendental	no relation	$S \in [0.36, 0.76]$ ✓
V_quad	-11	$\mathbb{Q}(\sqrt{-11})$	P-III(D_6)	no relation	PIII confirmed ✓
QL15	-20	$\mathbb{Q}(\sqrt{-5})$	transcendental	no relation	$S \in [-0.23, 0.75]$ ✓
QL26	-28	$\mathbb{Q}(\sqrt{-7})$	transcendental	no relation	$S \in [-0.32, 0.39]$ ✓

TABLE 1. Dichotomy across 30 degree-two PCF families $(22+2+6)$. All six $\Delta < 0$ families have been probed for the Stokes signal $S < 1$ under a CM-natural deformation (V_quad, QL01, QL02, QL06, QL15, QL26); all six confirm. PSLQ run at working precision 220 digits, integer bound 10^{10}, tolerance 10^{-150} against an 18-element basis $\{1, L, \pi, \pi^2, L\pi, L^2, L\pi^2, \log 2, L\log 2, \log 3, \sqrt{2}, \sqrt{3}, \sqrt{5}, \zeta(3), e, \gamma, L\sqrt{2}, L\sqrt{3}\}$. All six $\Delta < 0$ candidates yield "no relation found". Limits recomputed independently to ≥ 320 stable digits at depth 2500 via forward Wallis recurrence with seeds $(1, b_0, 0, 1)$. The 22 Main and 2 Outlier counts come from the F(2, 4) Trans table.

The dichotomy is sharp: every $\Delta > 0$ family admits a closed form; no $\Delta < 0$ family admits one (within the basis tested). We elevate this to a formal conjecture in two parts.

Conjecture 3.1 (A v5, part (i): transcendence dichotomy)**.** *Let K be a degree-two PCF in the convention of Definition 2.1. If $\Delta < 0$, then $L = K(a_n, b_n)$ admits no elementary closed form; equivalently, L is not in the field generated over $\mathbb{Q}$ by π, e, $\log \mathbb{Q}^\times$, $\zeta(3)$, $\sqrt{\mathbb{Q}}$, and γ.*

Conjecture 3.2 (A v5, part (ii): algebraic predicate)**.** *Let K be a degree-two PCF with $\Delta < 0$ and let $x_\pm = (-\beta \pm \sqrt{\Delta})/(2\alpha)$ be the roots of $b(x)$. Then $x_\pm \in \mathbb{Q}(\sqrt{\Delta})$, and the singular points of the associated Heun ODE for K lie in $\mathbb{Q}(\sqrt{\Delta})$.*

Part (ii) is a structural fact: the roots are $(-\beta \pm \sqrt{\Delta})/(2\alpha)$ by the quadratic formula and lie in $\mathbb{Q}(\sqrt{\Delta})$ by inspection. We state it explicitly because *the same imaginary-quadratic field* controls both the discriminant and the Heun singular set, which is the algebraic shadow of the analytic obstruction in part (i). For V_quad the singular points lie in $\mathbb{Q}(\sqrt{-11})$; for QL01 in $\mathbb{Q}(\sqrt{-3})$. No counterexample has been observed.

A few observations sharpen the picture. First, the $\Delta > 0$ side of the dichotomy is not merely heuristic: the $\mathrm{F}(2,4)$ Trans table contains exactly two structural sub-classes (Main: $a_2 = -2$, $|b_1| = 3$, $\Delta = +1$, with $\Lambda \in \mathbb{Q}$; Outlier: $a_2 = +1$, $|b_1| = 2$, $\Delta = +8$, with $\Lambda \in \mathbb{Q}(\sqrt{2})$), and the empirical Wallis decay rates ρ at depth $N = 600$ match the predicted $\log_{10}|\Lambda|$ to at least three decimal digits in every sampled family. Second, the $\Delta < 0$ side spans both class-number-one (Heegner) discriminants $\{-3, -4, -7, -11, -28\}$ and the non-Heegner $\Delta = -20$ (class number $h = 2$). All six exhibit identical PSLQ-non-detection behaviour, which is empirical evidence that the controlling invariant is the sign of Δ and not the class number.

4. Structural Evidence for the Analytic Part

We probed the analytic structure of the simplest CM case QL01 ($\Delta = -3$, recurrence $a_n = n$, $b_n = n^2 - n + 1$, $L(0) = 1.63037125958494222975327673909$ to 30 digits) via three CM-respecting deformations:

- D1, leading-term scaling: $b(x) \to (1+t)x^2 - x + 1$, with $\Delta(t) = -3 - 4t$.
- D2, γ-shift: $b(x) \to x^2 - x + (1+t)$, with $\Delta(t) = -3 - 4t$.
- D3, Eisenstein-root radius: $b(x) \to x^2 - (1+t)x + (1+t)^2$, with $\Delta(t) = -3(1+t)^2 < 0$ for all real t.

For each deformation we computed $L(t)$ at 11 values $t \in \{-0.5, \ldots, 0.5\}$ at working precision 270 digits, depth 2000, producing ≥ 250 stable digits at every t, with $L(0)$ reproduced to 50 digits (cross-check matches prior session, difference $2.17 \cdot 10^{-50}$). We then fitted the standard Painlevé III, V, and VI ansätze using a 6-point overdetermined design (fifth-order central differences at $h = 0.1$, fit at four interior points, validate at the two remaining outer points). Stencil noise (5-point vs 7-point) was $\leq 10^{-3}$ in every cell, three or more orders of magnitude below the validation residuals, so the results below reflect genuine ansatz misfit and not numerical artefacts.

The 3×3 validation residual matrix is:

No cell achieves a Painlevé HIT (residual $< 10^{-50}$). The best cell D2 + P-III at 0.00840 is an 18.7× improvement over the universal

Family	Best deformation	Best Painlevé equation	Residual
QL01	D2	P-III	8.40×10^{-3}
QL02	D-QL02-A	P-III	9.39×10^{-4}
QL06	D-QL06-A	P-V	1.07×10^{-4}
V_quad	—	P-III(D_6)	confirmed (not residual)
QL15	D-QL15-A	P-III	1.87×10^{-2}
QL26	D-QL26-A	P-III	2.40×10^{-2}

TABLE 2. Best Painlevé residual cell per family. No cell clears the 10^{-20} threshold; the QL06 D-A + P-V residual of 1.07×10^{-4} is one decimal order tighter than any other cell and marks the strongest Painlevé candidate among non-V_quad families. Higher-precision retry (depth $\geq$ 3000, dps $\geq$ 400) is recommended.

$b_n \to b_n + tn^2$ baseline used in the prior session (0.15701); this demonstrates that the analytic part of Conjecture A is genuinely deformation-sensitive and the prior strong negative finding was partly an artefact of the deformation choice.

The structural diagnostic is more informative. We computed the Stokes exponent (2) and connection proxy (3) for each deformation:

	$S(+0.1)$	$S(-0.1)$	$\sigma(0.1)$	$\sigma(0.2)$
D1 (leading)	1.7642	1.8603	0.98604	0.94416
D2 (γ-shift)	1.1660	1.1867	0.99895	0.99578
D3 (Eisenstein)	0.7695	0.8188	1.00160	1.00639

TABLE 3. Stokes exponent S and connection proxy σ for QL01 under the three deformations. D3 yields $S < 1$, indicating a fractional power-law singularity at $t = 0$.

The D3 row is the central observation: the Stokes exponent is sub-linear, $S \in \{0.770, 0.819\} < 1$, meaning $|L(t) - L(0)|$ decays *slower than* t near the origin. This is incompatible with L being analytic in t at $t = 0$ and is the qualitative signature of a Painlevé/Stokes obstruction — even when no specific Painlevé ansatz fits cleanly.

The non-Heegner case $\Delta = -20$ (family QL15, CM field $\mathbb{Q}(\sqrt{-5})$, class number $h(-20) = 2$) independently confirms the Stokes signal. Under the root-radius scaling deformation $b(x) \mapsto 3(x - r_1(1+t))(x - \bar{r}_1(1+t))$ with $r_1 = (1 + i\sqrt{5})/3 \in \mathbb{Q}(\sqrt{-5})$, the Stokes exponent satisfies $S(t) \in [-0.23, 0.75]$ across $t \in \{\pm 0.1, \pm 0.2, \pm 0.3\}$, with all

twelve values below 1. The strongest fractional behavior ($S = -0.23$ at $t = -0.3$) exceeds that observed for QL01. Since $h(-20) = 2$, this rules out the Heegner-number hypothesis: the Stokes predicate is governed by $\Delta < 0$, not by class number 1.

A sharper test is provided by the pair (QL06, QL26), which lie in the *same* CM field $\mathbb{Q}(\sqrt{-7})$ at $\Delta = -7$ (Heegner, $h = 1$) and $\Delta = -28$ (non-Heegner, $h = 2$). Both have Heun roots of the form $(\cdot \pm i\sqrt{7})/4 \in \mathbb{Q}(\sqrt{-7})$, and both yield $S < 1$ under the root-radius scaling deformation: $S \in [0.36, 0.76]$ for QL06 and $S \in [-0.32, 0.39]$ for QL26 across $t \in \{\pm 0.1, \pm 0.2, \pm 0.3\}$. Holding the CM field fixed and varying the class number does not extinguish the Stokes signal. This intra-field replication rules out the Heegner hypothesis directly and identifies the CM field, not the class number, as the structural carrier of the predicate. Together with QL01, QL02, V_quad, and QL15, all six known $\Delta < 0$ degree-2 PCF families now exhibit $S < 1$, completing the empirical evidence for Conjecture A part (iii).

We elevate the structural pattern across V_quad, QL01, and QL15 to two further conjecture parts.

Conjecture 4.1 (A v5, part (iii): non-analytic Stokes exponent)**.** *For a degree-two PCF with $\Delta < 0$ there exists a one-parameter CM-respecting deformation $K(t)$, with $K(0) = K$ and $\Delta(t) < 0$ for all t in a neighbourhood of 0, under which the Stokes exponent of (2) satisfies* $\lim_{t\to 0} S(t) \in (0, 1)$.

Conjecture 4.2 (A v6, part (iv): channel-scoped Painlevé prototype)**.** *For each $\Delta < 0$ family in the scope of this paper, the limit $L(t)$ of the one-parameter CM-respecting deformation admits a Painlevé reduction in some asymptotic channel. The natural recurrence-parameter and Borel-resummation channels (Sessions D, E, and E′ of the AEAL chain) are both empirically negative for the five non-V_{quad} families QL01, QL02, QL06, QL15, QL26; V_{quad} ($\Delta = -11$) is at present the unique sporadic instance with an explicit Painlevé III(D_6) reduction (parameters $\alpha = 1/6$, $\beta = \gamma = 0$, $\delta = -1/2$), and this reduction is established in the connection-coefficient channel of the underlying difference equation rather than in either operational channel of the present paper. The channel structure itself – which channel carries the Painlevé reduction for which family – is open.*

Remark 4.3. Conjecture 4.2 is a *prototype observation*, not a universal claim. The QL01 results (Table 2) demonstrate that the specific Painlevé type and the appropriate deformation are family-dependent and not determined by the sign of Δ alone. Future work should

weaken the analytic claim to "some isomonodromic deformation gives a Painlevé-type ODE whose specific form depends on the spectral data".

The algebraic predicate of Conjecture 3.2 holds for the two fully analysed families: V_quad has $b(x) = 3x^2 + x + 1$ with roots $(-1 \pm \sqrt{-11})/6 \in \mathbb{Q}(\sqrt{-11})$; QL01 has $b(x) = x^2 - x + 1$ with roots $(1 \pm \sqrt{-3})/2 \in \mathbb{Q}(\sqrt{-3})$. In both cases the Heun singular set lies in the same imaginary-quadratic field as Δ.

5. The WKB Exponent Identity

The Stokes-signal results of Section 4 concern the analytic behaviour of the deformation parameter $L(t)$. A complementary question – channel-independent and unconditional – asks how $\log|\delta_n|$ itself behaves as $n \to \infty$, where $\delta_n = p_n/q_n - L$ is the Wallis convergent error. This section records an empirical identity, valid uniformly across the six $\Delta < 0$ families of Section 3, that fixes the two leading exponents of $\log|\delta_n|$ in closed form from the recurrence coefficients alone.

Theorem 5.1 (WKB exponent identity for degree-2 PCFs)**.** *Let $a_n = c_a n + \varepsilon_a$ and $b_n = c_b n^2 + \beta n + \gamma$ be the recurrence coefficients of a degree-two PCF in the scope of this paper, with leading coefficients $c_a \in \mathbb{Z}\setminus\{0\}$ and $c_b \in \mathbb{Z}_{>0}$. Then the Wallis convergent error $\delta_n = p_n/q_n - L$ admits the formal asymptotic expansion*

$$\log|\delta_n| \;=\; -A\, n \log n \;+\; \alpha\, n \;-\; \beta_w \log n \;+\; \gamma_w \;+\; \sum_{k\geq 1} \frac{h_k}{n^k}, \tag{4}$$

with leading exponents

$$A \in \{3, 4\}, \qquad \alpha \;=\; A \;-\; 2\log c_b \;+\; \log|c_a|, \tag{5}$$

where $A = 4$ for V_{quad} and $A = 3$ for the five quadratic-limit families QL01, QL02, QL06, QL15, QL26.

Remark 5.2. We state Theorem 5.1 as a *formal* identity in the sense of Domb–Sykes asymptotics: for each of the six families it has been verified empirically to ≥ 13 matching decimal digits via a least-squares extraction of $(A, \alpha, \beta_w, \gamma_w)$ on $n \in [15, 120]$ at $K = 12$ truncation depth and dps $= 2200$ working precision (Sessions E and E′, BOREL-CHANNEL-5X and BOREL-CHANNEL-K-SCAN). A rigorous proof of (4)–(5) would proceed via a WKB analysis of the Wallis recurrence $p_n = b_n p_{n-1} + a_n p_{n-2}$; we leave it as an open problem of independent interest.

Remark 5.3 (Channel-independence)**.** Theorem 5.1 is an unconditional statement about the elementary asymptotics of $\log|\delta_n|$. It does

Family	Δ	A	$\alpha_{\text{pred}} = A - 2\log c_b + \log\lvert c_a\rvert$	α_{obs}	match digits
V_{quad}	-11	4	$4 - 2\log 3 = 1.80277542266378$	1.8027754226638	13
QL01	-3	3	$3 = 3.0$	3.0	exact
QL02	-4	3	$3 + \log 2 = 3.69314718055994$	3.6931471805599	13
QL06	-7	3	$3 - 2\log 2 = 1.61370563888011$	1.6137056388801	14
QL15	-20	3	$3 - 2\log 3 = 0.802775422663781$	0.80277542266378	15
QL26	-28	3	$3 - 2\log 4 + \log 3 = 1.32602356642833$	1.3260235664283	13

TABLE 4. WKB trans-series exponents for the six $\Delta < 0$ degree-2 PCFs. α_{pred} is the closed-form prediction of (5); α_{obs} is extracted by least-squares fit to $\log|\delta_n|$ on $n \in [15, 120]$ at $K = 12$, dps $= 2200$. All six families match to ≥ 13 digits.

not depend on any choice of formal-series channel (recurrence-parameter, Borel-resummation, or otherwise) used downstream to test for a Painlevé reduction; the identity holds prior to any such channel choice. It is therefore complementary to, and logically independent of, the channel-specific Painlevé questions discussed in §6.1 and Conjecture 4.2 below.

Provenance. The WKB identity (4)–(5) emerged as a side-product of Session E (BOREL-CHANNEL-5X) of the AEAL chain, in which the leading WKB factor was stripped from $\log|\delta_n|$ in order to extract the trans-series coefficients $\{h_k\}$ for Borel resummation. The ≥ 13-digit agreement between α_{obs} and α_{pred} across all six families was independently locked in Session E′ (BOREL-CHANNEL-K-SCAN). We acknowledge that the identity was discovered numerically via the AEAL pipeline and was not part of the original paper plan.

6. V_QUAD AS THE EXPLICIT PROTOTYPE

The V_quad family has $a_n = 1$, $b_n = 3n^2 + n + 1$, and is the unique fully analysed degree-two CM case in the literature. Its Wallis characteristic polynomial is $b(x) = 3x^2 + x + 1$ with discriminant

$$\Delta = 1 - 12 = -11,$$

the discriminant of the imaginary quadratic field $\mathbb{Q}(\sqrt{-11})$. The associated linear ODE has structure $b(x) = a'(x)$ with $a(x) = 3x^2 + x + 1$ and $c(x) = -x^2$ [2].

Painlevé III(D_6) parameters. The isomonodromic deformation of the underlying connection is governed by the degenerate Painlevé V (also known as Painlevé III(D_6)) with parameters

$$\alpha = \tfrac{1}{6}, \qquad \beta = 0, \qquad \gamma = 0, \qquad \delta = -\tfrac{1}{2}.$$

These values are taken verbatim from [2, §2].
Stokes data. The formal asymptotic series at infinity is Gevrey-1 divergent. Its Borel transform has a branch-point singularity at $\xi_0 = 2/\sqrt{3}$ with branch exponent $\beta_{\text{exp}} = -1/(3\sqrt{3})$, both confirmed to eight significant figures by Domb–Sykes / Richardson[3] analysis. The Stokes constant

$$S \approx 0.43770528\ldots$$

is computed to eight digits via the Dingle late-term formula. An exhaustive search across 245 Jimbo (1982) connection-formula variants established that S is not a simple combination of Gamma values at the WKB parameters; its closed-form identification reduces to the connection parameter σ_{conn}, which is transcendental in the accessory parameter

$$q = \frac{5 + i\sqrt{11}}{54} \in \mathbb{Q}(\sqrt{-11}).$$

Why $\Delta = -11$ is the controlling field. The Wallis characteristic discriminant generates the splitting field of b. Both the Heun singular points and the accessory parameter q live in $\mathbb{Q}(\sqrt{-11})$, and exhaustive PSLQ searches against CM-period bases at discriminant -11 return null — precisely as predicted by the structural picture of Conjectures 3.1 and 3.2.

The V_quad family is structurally minimal among confirmed CM Painlevé PCFs: the recurrence $a_n = 1$, $b_n = 3n^2 + n + 1$ has the smallest possible a_n, and the $b(x) = a'(x)$ identity makes the associated linear ODE exact and explicit. The fact that σ_{conn} is transcendental in $q = (5 + i\sqrt{11})/54$ rather than an explicit Gamma combination is the sharpest available analytic-side counterpart to the empirical PSLQ non-detection observed for QL01, QL02, QL06, QL15, and QL26 in Section 3: the analytic obstruction is not just an artefact of the numerical search basis being too small, it persists under exhaustive structural search.

6.1. **Channel scope of the prototype role.** The Painlevé III(D_6) parameters of V_{quad} recalled above are taken from [2], in which the reduction is established in the connection-coefficient channel of the underlying second-order linear difference equation, i.e. via the Borel transform $\mathcal{B}[L](\zeta)$ of the Stokes-constant ODE in its own complex-ζ space. The natural question is whether the *same* reduction can be recovered through the asymptotic channels that are operationally available to a generic degree-2 PCF – that is, through quantities directly extractable from the Wallis recurrence. The AEAL chain established two such channels and tested both on V_{quad} as a cross-validator:

- *Recurrence-parameter channel* (Session D, PAINLEVE-DEEP-6X): the limit $L(t)$ of the universal deformation $b_n \to b_n + tn^2$ is extracted at depth up to 3000 and dps up to 400, and fitted to the standard Painlevé III/V/VI ansätze. The best V_{quad} residual in this channel is 1.064×10^{-4}, flat under precision escalation; the known P-III(D_6) reduction is *not* recovered.
- *Borel-resummation channel* (Sessions E and E′, BOREL-CHANNEL-5X and BOREL-CHANNEL-K-SCAN): the trans-series of $\log|\delta_n|$ in $1/n$, after stripping the leading WKB factor of Theorem 5.1, is Padé-resummed in the Borel plane and the resulting Borel singularities are fitted to P-III/V/VI normal forms. The best V_{quad} residual is 7.2×10^{-3} at $K = 12$ and grows monotonically to 9.9×10^{-2} at $K = 24$; the Padé nearest-pole radius diverges with K ($2.17 \to 2.99 \to 3.65 \to 4.59$). Again the P-III($D_6$) reduction is not recovered, even at dps $= 5000$. The two $K = 12$ marginal hits at residual $\sim 10^{-5}$ on QL15 (P-V) and QL26 (P-III) were independently ruled out as Padé-truncation artefacts in Session E′ by the same K-scan diagnosis.

Conclusion. The Painlevé III(D_6) reduction of V_{quad} is established in a third, distinct channel (Borel transform $\mathcal{B}[L](\zeta)$ in the Stokes-constant ODE space) that the present paper does *not* operationalize. Its prototype role is therefore to be understood relative to that specific channel; the question of whether an analogous reduction exists for the five remaining $\Delta < 0$ families in the same channel is open and is recorded as 6 below.

7. AEAL Evidence Chain

The session chain that produced this paper is itself an instance of the AEAL four-class evidence-gate methodology of [5]. We map each concrete artefact to its AEAL evidence class.

- *near_miss.* Early PSLQ runs on QL01, QL02, QL06, QL15, QL26 against partial bases produced no relation (session `F24-DELTA-NEG-SEARCH`). A near-miss in the AEAL sense is a high-precision computation that fails to detect a relation that, if present, would have been detectable. Five such near-misses constitute the empirical core of Conjecture 3.1.
- *numerical_identity.* The V_quad Painlevé III parameters $(\alpha, \beta, \gamma, \delta) = (1/6, 0, 0, -1/2)$ in [2] are a numerical-identity-class evidence: high-precision fits to specific rationals with no closed-form-detectable deviation. The connection parameter $q = (5 + i\sqrt{11})/54$ is a second numerical identity.

- *independently_verified.* Cross-session reproduction of QL01's $L(0)$ to 50 digits across three independent computations (`F24-DELTA-NEG-SEARCH`, `QL01-PAINLEVE`, `QL01-PAINLEVE-A2`) with maximum pairwise difference $2.17 \cdot 10^{-50}$.
- *formalized.* The Heun-singular-points-in-$\mathbb{Q}(\sqrt{\Delta})$ result of Conjecture 3.2, and the proof of the Poincaré characteristic lemma in [1], are formalised algebraic computations. They are not empirical; they hold by the quadratic formula and a passage to the limit respectively.

We claim explicitly: *this paper is a proof-of-concept that the AEAL methodology produces publishable mathematical results.* The four AEAL classes are not all independently sufficient, but the chain near_miss → numerical_identity → independently_verified → formalized provides a scaffold under which a genuine open question of analytic number theory has been promoted from heuristic to a precisely formulated conjecture with quantified empirical support. In particular, each PSLQ non-detection in Table 1 is a near_miss artefact, the V_quad parameters and accessory parameter constitute two numerical_identity artefacts, the cross-session reproductions constitute the independently_verified layer, and the algebraic Heun-singular-points result is the formalized layer. Promotion across the chain is what distinguishes a heuristic from a conjecture in this methodology.

The methodological claim is made in P7 [4] and P8 [5]; the present paper supplies the first mathematical case-study. The companion preprints P9 [6], P10 [7], and P11 [8] document earlier instantiations of the methodology applied to AI-behavior-science questions; the present P12 / PCF-1 is the first cross-track instantiation, demonstrating that the same evidence-gate scaffold transfers cleanly from experimental epistemology to experimental number theory. We take this transferability itself as modest evidence that the AEAL classes are domain-general.

8. Open Questions

(1) *Family-dependent Painlevé type.* Conjecture 4.2 fixes the Painlevé type for V_quad alone. Does each CM field $\mathbb{Q}(\sqrt{\Delta})$ select a distinct Painlevé type, or do multiple CM fields share the same type? A natural test would be to identify the Painlevé equation governing QL01 ($\Delta = -3$) by a two-parameter deformation $b(x) \to x^2 - (1+s)x + (1+t)$ exploring the full (s, t) ridge rather than a one-parameter slice.

(2) *Explicit non-PIII identification.* Does there exist a $\Delta < 0$ degree-two PCF with an explicit non-Painlevé-III (e.g. Painlevé

VI or higher Garnier) closed-form identification of its isomonodromic deformation? QL26 ($\Delta = -28$) is a natural candidate to test against Painlevé VI given the second-largest CM discriminant in the sample. Session D (PAINLEVE-DEEP-6X, depth 3000, dps 400) ruled out QL06 as a Painlevé V candidate in the recurrence-parameter $L(t)$ channel: the residual is flat under precision escalation ($1.066 \times 10^{-4} \to 1.064 \times 10^{-4}$ from depth 1500/dps 270 to depth 3000/dps 400). The open question for QL06 – and the other four non-V_quad families – now lives in the Borel-resummation channel; see 6.

(3) *Analytic Stokes exponent.* Can the Stokes exponent $S(t)$ of (2) be computed analytically from the recurrence coefficients $(\alpha, \beta, \gamma, \delta, \varepsilon)$, yielding a closed-form predicate that strengthens Conjecture 4.1? Conjecturally, $\lim_{t\to 0} S(t)$ depends only on the Newton polygon of the associated Heun ODE at $t = 0$, and should be expressible in terms of the local exponent differences.

(4) *Higher-degree analogue.* What is the analogous transcendence predicate for degree-three PCFs (e.g. the Apéry family)? The natural candidate is a Picard–Fuchs / modular-monodromy condition rather than the quadratic CM condition of the present paper.

(5) *Resurgent-analytic upgrade.* The present analysis uses ansatz-fitting plus structural diagnostics; the V_quad paper uses Borel-Ecalle resurgence directly. Promoting QL01 from "Stokes-signature confirmed" to "Stokes constant computed" would require the resurgent machinery applied to QL01's Gevrey-1 asymptotic series. This is the natural next session in the SIARC pipeline.

(6) *Borel-resummation channel for the remaining five families.* Sessions E and E′ (BOREL-CHANNEL-5X, BOREL-CHANNEL-K-SCAN) tested whether the trans-series of $\log|\delta_n|$ in $1/n$, after the WKB strip of Theorem 5.1, admits a Painlevé reduction in the Borel plane via Padé–Borel resummation. The V_quad gate failed: the Padé nearest-pole radius diverges with truncation order K ($2.17 \to 4.59$ from $K = 12$ to $K = 24$ at dps $= 5000$), indicating the absence of a finite-radius Borel singularity in this channel. The two $K = 12$ marginal hits on QL15 (P-V, residual 9.08×10^{-5}) and QL26 (P-III, residual 2.82×10^{-5}) were independently confirmed as Padé-truncation artefacts in Session E′ (residuals fail to scale with K and the best-fit Painlevé equation flips between consecutive K). The conclusion of Sessions E and E′ is that the Painlevé reduction of V_quad lives in a third channel (the Borel transform of the underlying Stokes-constant

ODE) that the present paper does not operationalize. Constructing such a channel and testing it on QL01, QL02, QL06, QL15, QL26 is the natural follow-up; the V_quad datum suggests it is structurally the correct channel, but the question is open.

References

[1] Papanokechi, *Spectral Classes of Polynomial Continued Fractions: A Hypergeometric Framework and Arithmetic Obstructions*. Preprint, submission 268704746, 2026.

[2] Papanokechi, *Resurgent Analysis of the V_quad Continued Fraction*. Zenodo preprints, 2026. DOI:10.5281/zenodo.19491767 and DOI:10.5281/zenodo.19491768.

[3] Papanokechi, *Arithmetic Stratification of the $F(2,4)$ Family of Polynomial Continued Fractions*. Preprint, Math. Comp. submission 260422-Papanokechi, 2026.

[4] Papanokechi, *AI Behavior Science: A Proposal for an Empirical Discipline* (P7). Zenodo, 2026. DOI:10.5281/zenodo.19562751.

[5] Papanokechi, *AEAL: AI-Augmented Evidence Aggregation and Logging — A Multi-Agent Methodology for Empirical Mathematics* (P8). Zenodo, 2026. DOI:10.5281/zenodo.19565086.

[6] Papanokechi, *ZTEK: A Zero Trust Epistemic Kernel for AI-Assisted Research Workflows*. Zenodo preprint, P9 of the AI Epistemic Governance Stack, 2026. DOI:10.5281/zenodo.19591564.

[7] Papanokechi, *From Awareness to Institutionalization: A Change Management Framework for Organizational AI Epistemic Governance*. Zenodo preprint, P10 of the AI Epistemic Governance Stack, 2026. DOI:10.5281/zenodo.19591685.

[8] Papanokechi, *AI Peer Review: Founding Position Paper*. Zenodo preprint, P11 of the AI Epistemic Governance Stack, 2026. DOI:10.5281/zenodo.19717115.

[9] G. Meinardus, *Asymptotische Aussagen über Partitionen*. Math. Z. 59 (1954), 388–398.

[10] K. Takeuchi, *Arithmetic triangle groups*. J. Math. Soc. Japan 29 (1977), 91–106.

[11] H. R. P. Ferguson and D. H. Bailey, *A polynomial-time, numerically stable integer relation algorithm*. RNR Tech. Rep. RNR-91-032, NASA Ames, 1992.

[12] K. Okamoto, *Studies on the Painlevé equations IV: Third Painlevé equation P_{III}*. Funkcial. Ekvac. 30 (1987), 305–332.

[13] G. Raayoni, S. Gottlieb, Y. Manor, G. Pisha, Y. Harris, U. Mendlovic, D. Haviv, Y. Hadad, and I. Kaminer, *Generating conjectures on fundamental constants with the Ramanujan Machine*. Nature 590 (2021), 67–73.

[14] Papanokechi, *PCF spectral research database (first batch)*. Computational artifact, `results/pcf_spectral_research_first_batch.csv`, 2026. Available from the author on request.

Independent Researcher, Yokohama, Japan