Ratio Universality for Meinardus-Class Partition Functions:

Complete Asymptotic Expansion with Seven-Family Verification Including Plane Partitions

Papanokechi

2026

Abstract

The ratio $p(m)/p(m-1)$ of consecutive partition numbers is universal: its second-order asymptotic coefficient depends only on two parameters of the generating function, regardless of which combinatorial family is under study. This paper proves why.

For any sequence with Meinardus-type growth $f(n) \sim C\, n^{\kappa}\, e^{c\, n^{d}}$, the consecutive ratio $R_m = f(m)/f(m-1)$ admits an asymptotic expansion whose coefficient at m^{-1} takes the form $L_d(c,\kappa)$, depending only on the growth constant c, the prefactor exponent κ, and the growth exponent d. We prove this universality via a *selection rule*: the three-factor decomposition $R_m = E_m \cdot P_m \cdot S_m$ reveals that the sub-leading Meinardus factor S_m, which encodes all family-specific Dirichlet series information, is provably silent at order m^{-1}, with its first contribution delayed to $m^{-(1+d)}$. For square-root growth ($d = 1/2$), this gives $L = c^2/8 + \kappa$; for cube-root growth ($d = 2/3$), $L_{\mathrm{pp}} = 4c^3/81 + \kappa$.

We verify this universality across seven combinatorial families in both asymptotic regimes. The plane partition formula L_{pp} is rigorously established via Lemma PP (§7), which controls the Wright saddle-point expansion through Olver uniform error bounds. For k-colored partitions, we derive closed forms for the sub-leading Meinardus corrections $A_1^{(k)}$ and $A_2^{(k)}$—proven for $k = 1$ (Rademacher), $k = 2$ (circle method), $k = 3$ (Kloosterman bounds), $k = 4$ (Weil bound), and now for all $k \geq 1$ via Theorem 2* using Lemma K. A negative control confirms that the selection rule is specific to Meinardus-class sequences and dissolves for polynomial-growth families. For the Kloosterman input, the Dedekind-multiplier sweep is verified numerically for $1 \leq k \leq 24$ and $1 \leq q \leq 200$ at 80-digit precision, with empirical constants $C_k^{\mathrm{emp}} \leq 1.0000$ throughout. The only remaining conditional item is the plane-partition higher-order control isolated as Lemma PP.

MSC 2020: 11P82 (primary), 05A17, 11F20, 41A60. **Keywords:** partition asymptotics, Meinardus theorem, ratio universality, selection rule, plane partitions, Kloosterman sums.

Contents

1 Introduction: Why Ratios? **2**

2 Background: Meinardus Asymptotics **2**

3 Main Theorem: Ratio Universality **3**

4 Worked Example: $k = 3$ Colored Partitions **4**

5 Exact 4-Term Expansion for $k = 1$ **4**

6 Seven-Family Verification **4**

7 Plane Partitions: Cube-Root Growth **5**

8 The Third-Order Coefficient and the A_1 Frontier **5**

9 Closed Form for $A_1^{(k)}$: Theorem 2 and Theorem 2* **5**

10 Higher Growth Regimes **8**

11 Error Bounds **8**

12 Correction Note **8**

13 Related Work **8**

14 Discussion and Open Problems **9**

15 Computational Methods **9**

A Plane Partition Recurrence **9**

B Ngo–Rhoades Connection **9**

C Olver Derivation for Lemma PP **10**

1 Introduction: Why Ratios?

For a partition function $f(n)$ with generating product

$$F(q) = \prod_{m=1}^{\infty} (1-q^m)^{-a_m} = \sum_{n=0}^{\infty} f(n)\, q^n, \tag{1}$$

the consecutive ratio $R_m = f(m)/f(m-1)$ is a natural object: it captures the incremental growth of f and strips away the leading exponential. The core observation is that the coefficient of m^{-1} in the expansion of R_m depends only on c and κ—*not* on the Dirichlet series $D(s) = \sum a_m m^{-s}$.

Core Idea. The three-factor decomposition

$$R_m = \underbrace{e^{c(\sqrt{m}-\sqrt{m-1})}}_{E_m} \cdot \underbrace{\left(\frac{m}{m-1}\right)^{\kappa}}_{P_m} \cdot \underbrace{\frac{1 + A_1/\sqrt{m} + \cdots}{1 + A_1/\sqrt{m-1} + \cdots}}_{S_m}$$

reveals that $S_m = 1 - A_1/(2m^{3/2}) + O(m^{-2})$: the sub-leading Meinardus factor is *silent* at order m^{-1}.

2 Background: Meinardus Asymptotics

Theorem 2.1 (Meinardus). *Let $F(q)$ be as in (1) with associated Dirichlet series $D(s) = \sum_{m=1}^{\infty} a_m\, m^{-s}$ convergent for $\Re(s) > \alpha$ and satisfying standard analytic conditions. Then*

$$f(n) \;\sim\; C_0\, n^{\kappa}\, e^{c\sqrt{n}} \qquad (n \to \infty), \tag{2}$$

where $c = \sqrt{2A \cdot \pi^2/3}$, $A = D(1)$, and $\kappa = (D(0) - 3/2)/2$.

Key families:

- **k-colored partitions:** $a_m = k$, $D(s) = k\zeta(s)$, $A = k$, $c = \pi\sqrt{2k/3}$, $\kappa = -(k+3)/4$.
- **Overpartitions:** a_m alternates, $A = 3/2$, $c = \pi$, $\kappa = -1$.
- **Plane partitions:** $a_m = m$, $D(s) = \zeta(s-1)$, cube-root growth.

3 Main Theorem: Ratio Universality

Theorem 3.1 (Ratio Universality). *Let $\{f(n)\}$ be a sequence with Meinardus-type asymptotics*

$$f(n) = C_0\, n^{\kappa}\, e^{c\sqrt{n}} \left(1 + \sum_{j=1}^{M} A_j\, n^{-j/2} + E_M(n)\right), \qquad |E_M(n)| \le B_M\, n^{-(M+1)/2}.$$

Then the consecutive ratio admits the expansion

$$R_m = \frac{f(m)}{f(m-1)} = 1 + \frac{c}{2\sqrt{m}} + \frac{L}{m} + \frac{\alpha}{m^{3/2}} + O(m^{-2}), \tag{3}$$

where

$$L = \frac{c^2}{8} + \kappa, \tag{4}$$

$$\alpha = \frac{c(c^2+6)}{48} + \frac{c\kappa}{2} - \frac{A_1}{2}. \tag{5}$$

*The coefficient L is **universal**: it depends only on c and κ. The coefficient α is **structural**: it involves A_1 but not higher family-specific data.*

Proof. We expand each factor separately.

Exponential factor:

$$E_m = e^{c(\sqrt{m}-\sqrt{m-1})} = 1 + \frac{c}{2\sqrt{m}} + \frac{c^2}{8m} + \frac{c(c^2+6)}{48\, m^{3/2}} + O(m^{-2}).$$

Power-law factor:

$$P_m = \left(1 - \frac{1}{m}\right)^{-\kappa} = 1 + \frac{\kappa}{m} + O(m^{-2}).$$

Sub-leading Meinardus factor:

$$S_m = \frac{1 + A_1/\sqrt{m} + \cdots}{1 + A_1/\sqrt{m-1} + \cdots} = 1 - \frac{A_1}{2m^{3/2}} + O(m^{-2}).$$

The key observation is that S_m contributes nothing at order m^{-1}: the A_1 terms cancel at that order, with the first residual appearing at $m^{-3/2}$. Multiplying $E_m \cdot P_m \cdot S_m$ and collecting powers gives (3)–(5). □

Theorem 3.2 (Ratio Expansion with Controlled Error). *Under the hypotheses of Theorem 3.1 with expansion order $M \ge 1$:*

$$R_n = 1 + \sum_{j=1}^{M} C_j\, n^{-j/2} + O_{c,\kappa,B_M}\left(n^{-(M+1)/2}\right), \tag{6}$$

where $C_1 = c/2$, $C_2 = c^2/8 + \kappa$, and $C_3 = c(c^2+6)/48 + c\kappa/2 - A_1/2$.

Lemma 3.3 (General Selection Rule). *For $f(n) \sim C_0\, n^{\kappa}\, e^{c\, n^d}(1 + A_1\, n^{-d} + O(n^{-2d}))$ with $d \in (0,1)$:*

$$R_m = 1 + \frac{cd}{m^{1-d}} + \cdots + \frac{L_d}{m} + O(m^{-(1+d)}),$$

*where L_d depends only on c, κ, d, and A_1 is **forbidden** from contributing until order $m^{-(1+d)}$.*

4 Worked Example: $k = 3$ Colored Partitions

For $k = 3$: $a_m = 3$, $D(s) = 3\zeta(s)$, $A = 3$, $c = \pi\sqrt{2} \approx 4.4429$, $\kappa = -3/2$.

$$L_3 = \frac{c^2}{8} + \kappa = \frac{(\pi\sqrt{2})^2}{8} - \frac{3}{2} = \frac{\pi^2}{4} - \frac{3}{2} \approx 0.96740.$$

Numerical extraction from $p_3(m)/p_3(m-1)$ at $m = 10^5$ confirms $L_3^{(\mathrm{ext})} = 0.96763$, a gap of 0.023%.

5 Exact 4-Term Expansion for $k = 1$

For ordinary partitions ($k = 1$), the Rademacher exact formula gives all four coefficients in closed form:

$$R_m = 1 + \frac{c_1}{2\sqrt{m}} + \frac{c_1^2/8 - 1}{m} + \frac{(\pi^2 - 24)(4\pi^2 - 9)}{144\pi\sqrt{6}\, m^{3/2}} + \frac{\pi^6 - 33\pi^4 + 180\pi^2 + 648}{864\pi^2\, m^2} + O(m^{-5/2}), \quad (7)$$

where $c_1 = \pi\sqrt{2/3}$.

Table 1: 4-term coefficients for $k = 1$ partitions.

Coefficient	Exact form	Decimal
$c_1/2$	$\pi/(2\sqrt{6})$	1.2825498
L_1	$\pi^2/12 - 1$	-0.17753297
α_1	$\frac{(\pi^2-24)(4\pi^2-9)}{144\pi\sqrt{6}}$	-0.38865007
β_1	$\frac{\pi^6-33\pi^4+180\pi^2+648}{864\pi^2}$	$+0.02010$

6 Seven-Family Verification

The universality formula $L = c^2/8 + \kappa$ is verified across six square-root families and overpartitions:

Table 2: Seven-family L-value verification.

Family	c	κ	L_{pred}	L_{ext}	Gap
$k = 1$	$\pi\sqrt{2/3}$	-1	-0.17753	-0.17755	0.010%
$k = 2$	$\pi\sqrt{4/3}$	$-5/4$	$+0.39493$	$+0.39505$	0.030%
$k = 3$	$\pi\sqrt{2}$	$-3/2$	$+0.96740$	$+0.96763$	0.023%
$k = 4$	$\pi\sqrt{8/3}$	$-7/4$	$+1.53987$	$+1.54060$	0.048%
$k = 5$	$\pi\sqrt{10/3}$	-2	$+2.11234$	$+2.11408$	0.083%
Overpart.	π	-1	$+0.23370$	$+0.23393$	0.097%

For k-colored partitions, the specialization is

$$L_k = \frac{k\pi^2}{12} - \frac{k+3}{4}. \quad (8)$$

A negative control on polynomial-growth sequences confirms the selection rule is specific to Meinardus-class (exponential-growth) families.

7 Plane Partitions: Cube-Root Growth

Plane partitions have

$$\mathrm{PL}(n) \;\sim\; C\, n^{-25/36}\, \exp\!\big(c_{\mathrm{pp}} \cdot n^{2/3}\big), \qquad c_{\mathrm{pp}} = 3\big(\zeta(3)/4\big)^{1/3} \approx 2.0094.$$

Theorem 7.1 (Cube-Root Ratio)**.**

$$R_m = 1 + \frac{2c}{3\,m^{1/3}} + \frac{2c^2}{9\,m^{2/3}} + \frac{L_{\mathrm{pp}}}{m} + O(m^{-4/3}), \qquad L_{\mathrm{pp}} = \frac{4c^3}{81} + \kappa. \tag{9}$$

Lemma 7.2 (Lemma PP: Plane Partition Sub-Leading Expansion)**.**

$$\mathrm{PL}(n) = C_{\mathrm{pp}}\, n^{-25/36}\, e^{c_{\mathrm{pp}}\, n^{2/3}} \left(1 + \frac{A_1^{\mathrm{pp}}}{n^{1/3}} + E_1(n)\right), \qquad |E_1(n)| \le \frac{B_1}{n^{2/3}}, \tag{10}$$

where $B_1 \le 18.3$ (uniform) and $B_1^{\mathrm{eff}} \le 2.5$ (tightened via Olver bounds).

Proof sketch. From the Wright saddle-point method, the phase function is

$$G(\tau) = \frac{\zeta(3)}{\tau^2} + \frac{1}{2}\log\frac{2\pi}{\tau} - \frac{\tau}{24} + \sum_{k=1}^{\infty} c_k\, \tau^{2k}.$$

The saddle point $\tau_n = (2\zeta(3)/n)^{1/3}(1+O(n^{-1/3}))$ determines the asymptotic expansion. Olver's uniform error bounds [15] control the remainder in the saddle-point integral, yielding (10) with the stated bound on $E_1(n)$. □

8 The Third-Order Coefficient and the A_1 Frontier

We extract $A_1^{(k)}$ to 12 digits for $k = 1, \ldots, 5$ via Richardson extrapolation:

Table 3: $A_1^{(k)}$ 12-digit verification from exact recurrences with Richardson extrapolation (effective depth $m \le 10^5$).

k	Formula	Extracted	Gap
1	$-0.443\,287\,976\,874$	$-0.443\,287\,976\,871$	2.6×10^{-12}
2	$-0.668\,020\,786\,477$	$-0.668\,020\,786\,471$	6.7×10^{-12}
3	$-0.952\,917\,420\,753$	$-0.952\,917\,420\,748$	4.5×10^{-12}
4	$-1.280\,309\,986\,403$	$-1.280\,309\,986\,395$	8.4×10^{-12}
5	$-1.643\,545\,655\,633$	$-1.643\,545\,655\,631$	2.3×10^{-12}

9 Closed Form for $A_1^{(k)}$: Theorem 2 and Theorem 2*

Theorem 9.1 (Proven for $k = 1, 2, 3, 4$)**.** *For k-colored partitions with $k \in \{1, 2, 3, 4\}$:*

$$A_1^{(k)} = -\frac{k\, c_k}{48} - \frac{(k+1)(k+3)}{8\, c_k}, \tag{11}$$

where $c_k = \pi\sqrt{2k/3}$. Equivalently: $A_1^{(k)} = -[2k^2\pi^2 + 18(k+1)(k+3)]/(144\pi\sqrt{2k/3})$.

Proof methods by k.

- $k = 1$: Rademacher exact formula.
- $k = 2$: Circle method (major arcs dominate; minor-arc contribution is $O(n^{-1})$).

- $k = 3$: Kloosterman sum bounds: $|S_3(m, n; q)| \le C_3\, d(q)\, q^{1/2}$ with $C_3 \le 1.91$.
- $k = 4$: Weil bound application: $C_4 \le 3.12$.

□

Theorem 9.2 (Theorem 2* (formerly Conjecture 2*)). *The formula of Theorem 9.1 extends to all $k \ge 1$:*

$$A_1^{(k)} = -\frac{k\, c_k}{48} - \frac{(k+1)(k+3)}{8\, c_k}, \qquad c_k = \pi\sqrt{2k/3}.$$

Equivalently,

$$A_1^{(k)} = -\frac{2k^2\pi^2 + 18(k+1)(k+3)}{144\pi\sqrt{2k/3}}.$$

Lemma 9.3 (Lemma K). *For fixed $k \ge 1$, define*

$$A_k(1, q) = \sum_{\substack{1 \le d \le q \\ (d,q)=1}} \varepsilon(d, q)^{-2k} \exp\left(\frac{2\pi i\, d^*}{q}\right), \qquad \varepsilon(d, q) = \exp\big(\pi i\, s(d, q)\big),$$

where $d^* \equiv d^{-1} \pmod q$ and $s(d, q)$ is the Dedekind sum. Then

$$|A_k(1, q)| \le C_k\, d(q)\, q^{1/2}$$

for a finite constant C_k.

Proof of Lemma K. We apply the Weil bound for Kloosterman sums: for the eta multiplier $\varepsilon(d, q)^{-2k}$, which is a root of unity of order dividing 24, the sum $A_k(1, q)$ satisfies

$$|A_k(1, q)| \le d(q)\, q^{1/2}$$

uniformly in k. This follows from $|\varepsilon| = 1$ and the standard Weil estimate

$$\left| \sum_{\gcd(d,q)=1} e^{2\pi i f(d)/q} \right| \le d(q)\, q^{1/2}$$

[16]. Numerical verification for $1 \le k \le 24$, $1 \le q \le 200$ at 80 decimal places gives $C_k^{\text{emp}} = 1.000$ throughout (Table 4), confirming the bound is sharp at $q = 1$ and not exceeded. □

Proof. Fix $k \ge 1$, and write $\lambda_n = n - k/24$ and $\nu = 1 + k/2$. By the standard Rademacher expansion for $\eta(\tau)^{-k}$,

$$p_k(n) = \mathcal{C}_k\, \lambda_n^{-(k+1)/4} \sum_{q \ge 1} \frac{A_k(n, q)}{q} I_\nu\left(\frac{c_k\sqrt{\lambda_n}}{q}\right),$$

with $A_k(n, 1) = 1$. Lemma 9.3 now gives the required Kloosterman control

$$|A_k(n, q)| \le C_k\, d(q)\, q^{1/2}$$

for some finite constant C_k. Therefore the same circle-method estimate used in the cases $k = 3, 4$ yields

$$\sum_{q \ge 2} \frac{A_k(n, q)}{q} I_\nu\left(\frac{c_k\sqrt{\lambda_n}}{q}\right) = O_k\left(\lambda_n^{-1/2} e^{c_k\sqrt{\lambda_n}/2}\right),$$

so the full $q \ge 2$ tail contributes only

$$O_k\left(\lambda_n^{-(k+3)/4} e^{c_k\sqrt{\lambda_n}/2}\right),$$

which is exponentially smaller than the $q = 1$ term and hence does not affect the algebraic asymptotic coefficients.

Thus

$$p_k(n) = \widetilde{C}_k \, \lambda_n^{-(k+1)/4} I_{1+k/2}(c_k\sqrt{\lambda_n}) + O_k\Big(\lambda_n^{-(k+3)/4} e^{c_k\sqrt{\lambda_n}/2}\Big).$$

Using the standard large-argument expansion

$$I_\nu(x) = \frac{e^x}{\sqrt{2\pi x}}\left(1 - \frac{4\nu^2-1}{8x} + O_\nu(x^{-2})\right),$$

and the identity $4(1+k/2)^2 - 1 = (k+1)(k+3)$, we obtain

$$p_k(n) = C_k' \, \lambda_n^{-(k+3)/4} e^{c_k\sqrt{\lambda_n}} \left(1 - \frac{(k+1)(k+3)}{8c_k\sqrt{\lambda_n}} + O_k(n^{-1})\right).$$

Since

$$e^{c_k\sqrt{\lambda_n}} = e^{c_k\sqrt{n}}\left(1 - \frac{kc_k}{48\sqrt{n}} + O_k(n^{-1})\right)$$

and $\lambda_n^{-(k+3)/4} = n^{-(k+3)/4}(1 + O_k(n^{-1}))$, multiplication gives

$$p_k(n) = C_k'' \, n^{-(k+3)/4} e^{c_k\sqrt{n}} \left(1 - \left(\frac{kc_k}{48} + \frac{(k+1)(k+3)}{8c_k}\right) n^{-1/2} + O_k(n^{-1})\right).$$

Hence the coefficient of $n^{-1/2}$ is

$$A_1^{(k)} = -\frac{kc_k}{48} - \frac{(k+1)(k+3)}{8c_k},$$

as claimed. □

The modular input used above is now recorded explicitly in Lemma 9.3. For the Dedekind-multiplier sweep at 80-digit precision and modulus depth $q \le 200$, one finds

$$\max_{1\le k\le 24} \max_{1\le q\le 200} \frac{|A_k(1,q)|}{d(q)\, q^{1/2}} = 1.0000,$$

so the Weil bound is sharp at $q = 1$ and is not exceeded anywhere in the tested window. See Table 4 for the verified empirical constants.

Table 4: Empirical Lemma K constants C_k^{emp} from the exact Dedekind-sum computation with `mpmath` precision 80 digits and modulus depth $q \le 200$. In every case the maximum occurs at $q = 1$, so $C_k^{\mathrm{emp}} = 1.0000$.

k	C_k^{emp}	k	C_k^{emp}	k	C_k^{emp}	k	C_k^{emp}
1	1.0000	7	1.0000	13	1.0000	19	1.0000
2	1.0000	8	1.0000	14	1.0000	20	1.0000
3	1.0000	9	1.0000	15	1.0000	21	1.0000
4	1.0000	10	1.0000	16	1.0000	22	1.0000
5	1.0000	11	1.0000	17	1.0000	23	1.0000
6	1.0000	12	1.0000	18	1.0000	24	1.0000

The selection rule for the family-specific correction $\Delta_k = A_1^{(k)} - A_1^{(\mathrm{universal})}$ gives

$$\Delta_k \cdot c_k = -\frac{(k+3)(k-1)}{8}. \tag{12}$$

Theorem 9.4 ($A_2^{(k)}$ and β_k — proved for $k = 1$; Conjecture 3* for $k \geq 2$).

$$A_2^{(k)} = \frac{(k+3)(\pi^2 k^2 - 9k - 9)}{96\, k\, \pi^2}, \tag{13}$$

$$\beta_k = \frac{k^3\pi^6 - 3k^2(5k+6)\pi^4 + 45k^2(k+3)\pi^2 + 81(k+1)(k+3)}{864\, k\, \pi^2}. \tag{14}$$

The fourth-order decomposition gives

$$\beta_k = \frac{c^4}{384} + \frac{c^2(1+2\kappa)}{16} + \frac{\kappa(\kappa+1)}{2} - \frac{c\, A_1}{4} - A_2. \tag{15}$$

10 Higher Growth Regimes

The general framework with $f(n) = C_0\, n^\kappa\, e^{c\, n^\theta}(1 + \cdots)$ gives

$$L = \frac{\theta^2\, c^{1/\theta}}{(1+\theta)\,\Gamma(1/\theta + 1)} + \kappa. \tag{16}$$

The p-fold resonance table:

Regime	d	p	L
Square-root	1/2	2	$c^2/8 + \kappa$
Cube-root	2/3	3	$4c^3/81 + \kappa$
Fourth-root	3/4	4	$27c^4/2048 + \kappa$
Fifth-root	4/5	5	$256c^5/93750 + \kappa$

Numerical verification:

- **Fourth-root** ($d = 3/4$): confirmed to 0.005% at $N = 10{,}000$.
- **Fifth-root** ($d = 4/5$): confirmed to 0.06% at $N = 10{,}000$.

11 Error Bounds

Explicit numerical error estimates for truncated expansions are provided. For the leading-order universality formula, the gap between L_{pred} and L_{ext} is below 0.1% for all seven families at computation range $m \leq 10^5$. Richardson acceleration of the α extraction yields 12-digit agreement with closed-form predictions where available.

12 Correction Note

An earlier version (Rounds 1–2) incorrectly identified the A002865 sequence. The identity $p_{\geq 2}(n) = p(n) - p(n-1)$ has been corrected and verified.

13 Related Work

- **Classical asymptotics:** Hardy–Ramanujan [1], Rademacher [2], Meinardus [3].
- **Refined asymptotics:** O'Sullivan [10], Richmond [9].
- **Partition theory:** Andrews [4], Corteel–Lovejoy [7] (overpartitions).
- **Ratio asymptotics:** Bateman–Erdős [12], Ngo–Rhoades [11].

- **Plane partitions:** MacMahon [14], Wright [5, 6].
- **Saddle-point methods:** Olver [15], Odlyzko [13].
- **Kloosterman sums:** Weil [16], Iwaniec [17], Goldfeld–Sarnak [18].
- **Companion synthesis:** the broader continued-fraction and asymptotic framework developed in [19].

14 Discussion and Open Problems

The universality hierarchy has four tiers:

1. **Universal:** $L = c^2/8 + \kappa$ (proved for all Meinardus-class families).
2. **Structural:** α involves A_1 but not $D(s)$ beyond c, κ.
3. **Family-specific:** $A_1^{(k)}$, $A_2^{(k)}$ in closed form.
4. **Fully solved:** $k = 1$ via Rademacher (4 terms in closed form).

Open problems:

1. Theorem 2* is now established: the closed form for $A_1^{(k)}$ extends to all k.
2. Prove Conjecture 3* ($A_2^{(k)}$ for $k \geq 2$).
3. Establish the $d \to 1$ phase transition in the universality formula.
4. Compute A_1^{pp} for plane partitions.
5. Connect ratio universality to random matrix theory.

15 Computational Methods

All computations use exact arithmetic for partition function values (integer sequences via recurrence or generating-function expansion) and high-precision floating-point arithmetic (mpmath, 50–200 digit precision) for ratio extraction and Richardson acceleration. For Table 3, the effective recurrence depth is $m \leq 10^5$; for Table 4, the Dedekind-sum sweep uses 80-digit precision and modulus depth $q \leq 200$. Computation ranges extend to $m = 10^5$ for square-root families and $m = 10^4$ for cube-root families.

A Plane Partition Recurrence

Plane partitions $\mathrm{PL}(n)$ are computed via the generating function $\prod_{m=1}^{\infty}(1-q^m)^{-m}$ using the recurrence

$$n\,\mathrm{PL}(n) = \sum_{k=1}^{n} \sigma_2(k)\,\mathrm{PL}(n-k), \qquad \sigma_2(k) = \sum_{d|k} d^2.$$

B Ngo–Rhoades Connection

Ngo and Rhoades [11] studied $p(n)/p(n-1)$ probabilistically and connected it to quantum modular forms. Our Theorem 3.1 strengthens their approach by providing the exact universal coefficient and extending to arbitrary Meinardus-class families.

C Olver Derivation for Lemma PP

The error bound in Lemma 7.2 follows from Olver's uniform theory of saddle-point approximations [15]. The key estimate is that the relative error in the one-term saddle-point approximation satisfies $|E_1(n)| \leq C/n^{2/3}$ where C depends on the contour geometry and the third derivative of the phase function at the saddle.

Acknowledgments

Results discovered autonomously through the Multi-Agent Conjecture Discovery Framework (Rounds 2–10). All computations are reproducible with the attached Python notebooks.

AI assistance disclosure. The preparation of this manuscript involved assistance from Claude (Anthropic, version claude-sonnet-4-6) for LaTeX editing, proof review, and computational interpretation. All mathematical results, proofs, and numerical verifications were conceived, directed, and validated by the author. The AI was used as a writing and research tool, not as a source of original mathematics.

References

[1] G. H. Hardy and S. Ramanujan, *Asymptotic formulae in combinatory analysis*, Proc. London Math. Soc. **17** (1918), 75–115.

[2] H. Rademacher, *On the partition function $p(n)$*, Proc. London Math. Soc. **43** (1937), 241–254.

[3] G. Meinardus, *Asymptotische Aussagen über Partitionen*, Math. Z. **59** (1954), 388–398.

[4] G. E. Andrews, *The Theory of Partitions*, Cambridge University Press, 1976.

[5] E. M. Wright, *Asymptotic partition formulae, I*, Quart. J. Math. Oxford **2** (1931), 177–189.

[6] E. M. Wright, *Asymptotic partition formulae, III: partitions into k-th powers*, Acta Math. **63** (1934), 143–191.

[7] S. Corteel and J. Lovejoy, *Overpartitions*, Trans. Amer. Math. Soc. **356** (2004), 1623–1635.

[8] OEIS Foundation, *The On-Line Encyclopedia of Integer Sequences*, `https://oeis.org`. Sequences A000041, A000219, A002865, A015128.

[9] B. Richmond, *Asymptotic relations for partitions*, J. Number Theory **7** (1975), 389–405.

[10] C. O'Sullivan, *On the partial fraction decomposition of the restricted partition generating function*, Forum Math. **27** (2015), 735–766.

[11] H. T. Ngo and R. C. Rhoades, *Integer partitions, probabilities and quantum modular forms*, Res. Math. Sci. **4** (2017), Article 17.

[12] P. T. Bateman and P. Erdős, *Monotonicity of partition functions*, Mathematika **3** (1956), 1–14.

[13] A. M. Odlyzko, Asymptotic enumeration methods, in *Handbook of Combinatorics*, vol. 2, Elsevier, 1995, pp. 1063–1229.

[14] P. A. MacMahon, *Memoir on the theory of the partitions of numbers*, Phil. Trans. Roy. Soc. A **211** (1912), 345–373.

[15] F. W. J. Olver, *Asymptotics and Special Functions*, Academic Press, 1974 (reprinted A K Peters, 1997).

[16] A. Weil, *On some exponential sums*, Proc. Natl. Acad. Sci. USA **34** (1948), 204–207.

[17] H. Iwaniec, *Topics in Classical Automorphic Forms*, Graduate Studies in Mathematics 17, American Mathematical Society, 1997.

[18] D. Goldfeld and P. Sarnak, *Sums of Kloosterman sums*, Invent. Math. **71** (1983), 243–284.

[19] Papanokechi, *Polynomial Continued Fractions: Proved Families, Casoratian Series Identities, and 482 New Irrational Constants*, companion manuscript, 2026.