

A
Immune-high vs Immune-low
reference = Immune-low | up 3,718; down 228
−log10(BH q)
Molecular-group-adjusted log2 fold change
B
Immune-intermediate vs Immune-low
reference = Immune-low | up 431; down 69
−log10(BH q)
Molecular-group-adjusted log2 fold change
C
Immune-high vs Immune-intermediate
reference = Immune-intermediate | up 808; down 117
−log10(BH q)
Molecular-group-adjusted log2 fold change
D
404/431 (93.7%) of the Immune-intermediate set
is contained in the Immune-high set
Immune-high vs Immune-low
404
3,314 unique
total 3,718
27 unique
Immune-intermediate vs Immune-low
404
total 431
Up-regulated genes (BH q < 0.05, adj. log2FC > 1)
E
All 18,564 genes; Pearson r = 0.78
slope above identity is consistent with an ordered
shared immune-presence axis
identity (y = x)
orthogonal fit (slope = 1.90)
adj. log2FC Immune-high vs Immune-low
adj. log2FC Immune-intermediate vs Immune-low
F
Median 1.51× larger in Immune-high
paired Wilcoxon signed-rank P = 1.6 × 10−65
adj. log2FC on the 404 shared genes
Immune-intermediate
vs Immune-low
Immune-high
vs Immune-low