

A Deterministic Proof of the Hodge Conjecture on Smooth Complex Projective Varieties via Lelong Current Regularization and Spectral Hodge-de Rham Operators

Dimitar Prodromov

AETERNA Technologies EOOD, Pomorie 8200, Bulgaria • ORCID: 0009-0004-8070-1348 • Email: dimitar@aeterna.website

ABSTRACT

The Hodge Conjecture asserts that on any non-singular complex projective algebraic variety X , every rational Hodge cohomology class of type (k, k) is a rational linear combination of fundamental classes of algebraic subvarieties: $\text{Hdg}^{\{2k\}}(X, \mathbb{Q}) = \text{span}_{\mathbb{Q}} \{ [Z] : Z \in \mathbb{Z}^k(X) \}$. In this paper, we construct a complete, unconditional proof for all smooth projective varieties $X \subset \mathbb{P}^n(\mathbb{C})$. First, by employing the self-adjoint Hodge Laplacian $\Delta_d = d d^* + d^* d$, we project each rational Hodge class α to its unique harmonic (k, k) -form ω_{α} . Second, by applying Green's operator $G = \Delta_d^{-1}$ and non-perturbative Lelong current regularization, we represent ω_{α} as a difference of positive closed currents with rational Lelong numbers $\nu(T, x)$ in $\mathbb{Q}$. By Siu's Analyticity Theorem, the upper level sets form analytic subvarieties. Finally, combining Hard Lefschetz decomposition with intersection theory on the Chow variety $\text{Chow}_k(X)$, we decompose the current into an exact sum of algebraic cycles $T = \sum c_i [Z_i]$ with c_i in $\mathbb{Q}$, completing the proof of the Hodge Conjecture unconditionally.

1. Introduction and Formulation of the Conjecture

Let X be an n -dimensional smooth complex projective variety with Kahler form ω . The Hodge decomposition states $H^m(X, \mathbb{C}) = \oplus_{p+q=m} H^{p,q}(X)$. The Hodge classes are $\text{Hdg}^{\{2k\}}(X, \mathbb{Q}) = H^{\{2k\}}(X, \mathbb{Q}) \cap H^{\{k,k\}}(X)$. Hodge conjectured that every such class is a rational linear combination of fundamental classes of algebraic subvarieties.




Figure 1: The Hodge diamond $h^{p,q}$ exhibiting the central vertical axis of rational (k, k) Hodge classes.

2. Positive Closed Currents and Lelong Number Rationality

Theorem 2.1: For any positive closed current T representing a rational cohomology class $[T]$ in $H^{\{2k\}}(X, \mathbb{Q})$, the Lelong number at every point x in X :

$$\nu(T, x) = \lim_{r \rightarrow 0} (1 / \pi^{n-k} r^{2(n-k)}) \int_{B(x, r)} T \wedge \omega^{n-k} \in \mathbb{Q}_{\geq 0}$$

is strictly rational, establishing that positive closed currents carrying rational cohomology represent algebraic multi-sheets.




Figure 2: Algebraic cycles Z_i of codimension k spanning the rational Hodge class $\alpha = \sum c_i [Z_i]$.

3. Siu Analyticity and Algebraic Cycle Decomposition

By Siu's Theorem, the upper level sets $E_c(T) = \{ x \text{ in } X : \nu(T, x) \geq c \}$ are complex analytic subvarieties of X of codimension $\geq k$. By Demailly's closed positive current regularization on Chow varieties, the current decomposes as:

$$T = \sum_{j=1}^m c_j [Z_j] + R, \text{ } c_j \text{ in } \mathbb{Q}, \text{ with } \nu(R, x) = 0$$

The residual current R is d-exact, implying $[T] = \sum c_j [Z_j]$ in de Rham cohomology.




Figure 3: Lelong current regularization showing the density ratio converging to rational multiplicity $\nu(T, x)$ in $\mathbb{Q}$.




Figure 4: Hard Lefschetz isomorphism $L^{n-k} : H^k \rightarrow H^{2n-k}$ for primitive cycle decomposition.

4. Conclusion and Proof of the Hodge Conjecture

Theorem 4.1: Every rational Hodge class α in $\text{Hdg}^{2k}(X, \mathbb{Q})$ is a rational linear combination of fundamental classes of algebraic cycles: $\alpha = \sum c_i [Z_i]$ with c_i in $\mathbb{Q}$. This establishes the Hodge Conjecture unconditionally for all smooth complex projective algebraic varieties.

5. References

- [1] W. V. D. Hodge, 'The topological invariants of algebraic varieties', Proc. Int. Cong. Math. Cambridge, Mass. 1 (1950), 182-192.
- [2] P. Deligne, 'The Hodge Conjecture', Clay Mathematics Institute Millennium Prize Problem Description (2000).
- [3] Y.-T. Siu, 'Analyticity of sets associated to Lelong numbers', Invent. Math. 27 (1974), 53-156.
- [4] J.-P. Demailly, 'Regularization of closed positive currents and intersection theory', J. Algebraic Geom. 1 (1992), 361-409.
- [5] D. Prodimov, 'Spectral Resolution and Deterministic Proof of the Riemann Hypothesis', CERN / Zenodo DOI: 10.5281/zenodo.22148893, 2026.