

1 The Portmanteau Theorem

Theorem 1.1 (The Portmanteau Theorem, Theorem 2.1, [1])

Suppose:

- (S, ρ) is a metric space, $\mathcal{B}(S)$ its the Borel σ -algebra, $(S, \mathcal{B}(S))$ is the corresponding measurable space.
- $P, P_1, P_2, \dots \in \mathcal{M}_1(S, \mathcal{B}(S))$ are probability measures on $(S, \mathcal{B}(S))$.

Then, the following are equivalent:

- (i) P_n converges weakly to P , i.e. for each bounded continuous \mathbb{R} -valued function $f : S \rightarrow \mathbb{R}$, we have

$$\lim_{n \rightarrow \infty} \int_S f(s) dP_n(s) = \int_S f(s) dP(s).$$

- (ii) For each closed set $F \subset S$, we have

$$\limsup_{n \rightarrow \infty} P_n(F) \leq P(F).$$

- (iii) For each open set $G \subset S$, we have

$$\limsup_{n \rightarrow \infty} P_n(G) \geq P(G).$$

- (iv) For each P -continuity set $A \in \mathcal{B}(S)$, i.e. $P(\partial A) = 0$, we have

$$\lim_{n \rightarrow \infty} P_n(A) = P(A).$$

PROOF

(i) \implies (ii)

For each $\varepsilon > 0$, by Lemma A.2(ii), choose a bounded continuous functions $f_\varepsilon : S \rightarrow [0, 1]$ such that

$$I_F \leq f_\varepsilon \leq I_{F^\varepsilon}.$$

This implies, for each $\varepsilon > 0$, we have

$$P_n(F) = \int_S I_F(x) dP_n(x) \leq \int_S f_\varepsilon(x) dP_n(x).$$

By (i), we thus have

$$\limsup_{n \rightarrow \infty} P_n(F) \leq \lim_{n \rightarrow \infty} \int_S f_\varepsilon(x) dP_n(x) = \int_S f_\varepsilon(x) dP(x) \leq \int_S I_{F^\varepsilon}(x) dP(x) = P(F^\varepsilon).$$

By Lemma A.2(i), we have $F^\varepsilon \downarrow F$ as $\varepsilon \downarrow 0$. Hence, $P(F^\varepsilon) \downarrow P(F)$ as $\varepsilon \downarrow 0$. We may now conclude:

$$\limsup_{n \rightarrow \infty} P_n(F) \leq \lim_{\varepsilon \rightarrow 0^+} P(F^\varepsilon) = P(F).$$

□

A Technical Lemmas

Lemma A.1 Suppose (S, ρ) is a metric space, and $A \subset S$ is an arbitrary non-empty subset. Define

$$\rho(\cdot, A) : S \longrightarrow \mathbb{R} : x \longmapsto \inf_{y \in A} \{ \rho(x, y) \}$$

Then,

- (i) $\rho(\cdot, A)$ is a continuous \mathbb{R} -valued function on S .
- (ii) For each $x \in S$, $\rho(x, A) = 0$ if and only if $x \in \overline{A}$.

PROOF

- (i) Suppose $x_n \longrightarrow x$. We need to prove $\rho(x_n, A) \longrightarrow \rho(x, A)$, which follows immediately from the following two Claims:

Claim 1: $\rho(x, A) \leq \liminf_{n \rightarrow \infty} \rho(x_n, A)$.

Claim 2: $\limsup_{n \rightarrow \infty} \rho(x_n, A) \leq \rho(x, A)$.

Proof of Claim 1: For each $y \in S$, we have:

$$\rho(x, y) \leq \rho(x, x_n) + \rho(x_n, y).$$

Hence,

$$\rho(x, A) = \inf_{y \in A} \rho(x, y) \leq \rho(x, x_n) + \inf_{y \in A} \rho(x_n, y) = \rho(x, x_n) + \rho(x_n, A).$$

Since $\rho(x, x_n) \longrightarrow 0$, the preceding inequality implies

$$\rho(x, A) \leq \liminf_{n \rightarrow \infty} \rho(x_n, A).$$

This proves Claim 1.

Proof of Claim 2: For each $y \in S$, we have:

$$\rho(x_n, y) \leq \rho(x_n, x) + \rho(x, y).$$

Hence,

$$\rho(x_n, A) = \inf_{y \in A} \rho(x_n, y) \leq \rho(x_n, x) + \inf_{y \in A} \rho(x, y) = \rho(x_n, x) + \rho(x, A).$$

Since $\rho(x, x_n) \longrightarrow 0$, the preceding inequality implies

$$\limsup_{n \rightarrow \infty} \rho(x_n, A) \leq \rho(x, A).$$

This proves Claim 2.

- (ii)

$$\begin{aligned} \rho(x, A) = 0 &\iff \inf_{y \in A} \rho(x, y) = 0 \\ &\iff \text{For each } \varepsilon > 0, \text{ there exists } y \in A \text{ such that } \rho(x, y) < \varepsilon \\ &\iff y \in \overline{A} \end{aligned}$$

□

Lemma A.2 Suppose (S, ρ) is a metric space, and $A \subset S$ is a non-empty subset. For each $\varepsilon > 0$, define

$$A^\varepsilon := \{ s \in S \mid \rho(s, A) < \varepsilon \}.$$

Then the following are true:

- (i) $A^\varepsilon \downarrow \overline{A}$, as $\varepsilon \downarrow 0$.
- (ii) There exists a bounded continuous \mathbb{R} -valued function $f : S \rightarrow \mathbb{R}$ such that

$$I_{\overline{A}}(x) \leq f(x) \leq I_{A^\varepsilon}(x), \quad \text{for each } x \in S.$$

PROOF

- (i)
- (ii) Define $f : S \rightarrow \mathbb{R}$ as follows:

$$f(x) := \max \left\{ 0, 1 - \frac{\rho(x, A)}{\varepsilon} \right\}.$$

Then, by Lemma A.1(i), f is continuous \mathbb{R} -valued function on S . Clear, $0 \leq f(x) \leq 1$, for each $x \in S$. By Lemma A.1(ii), we have

$$x \in \overline{A} \iff \rho(x, F) = 0 \iff f(x) = 1.$$

This proves $I_{\overline{A}}(x) \leq 1 = f(x)$, for each $x \in \overline{A}$, and hence for each $x \in S$ (since $I_{\overline{A}}(x) = 0$ for $x \in S \setminus \overline{A}$, and the inequality holds trivially). On the other hand,

$$x \in S \setminus A^\varepsilon \iff \varepsilon \leq \rho(x, A) \iff 1 - \frac{\rho(x, A)}{\varepsilon} \leq 0 \implies f(x) = 0.$$

This proves $f(x) = 0 \leq I_{A^\varepsilon}(x)$, for each $x \in S \setminus A^\varepsilon$, and hence for each $x \in S$ (since $I_{A^\varepsilon}(x) = 1$ for each $x \in A^\varepsilon$ and the inequality holds trivially). This completes the proof of (ii). □

References

- [1] BILLINGSLEY, P. *Convergence of Probability Measures*, second ed. John Wiley & Sons, 1999.