

# Unit 3 :: Complex Analysis

☞ PHY0400404: Mathematical Physics ☞

Dr. Parag Bhattacharya  
Department of Physics  
Rangapara College (Autonomous)



# Syllabus

## Unit 3: Complex analysis

Functions of complex variables, analyticity and Cauchy-Riemann conditions, examples of analytic functions; singular functions: poles and branch points, order of singularity; integration of functions with complex variable, Cauchy's integral theorem and Cauchy's integral formula, simply and multiply connected regions; Laurent and Taylor's series expansions, residue theorem with application.

## Complex trigonometric and hyperbolic identities

$$1. \cosh x = \frac{e^x + e^{-x}}{2}$$

$$2. \sinh x = \frac{e^x - e^{-x}}{2}$$

$$3. \cos x = \frac{e^{ix} + e^{-ix}}{2}$$

$$4. \sin x = \frac{e^{ix} - e^{-ix}}{2i}$$

Using the above identities, it can be shown that

$$1. \cosh ix = \cos x$$

$$2. \cos ix = \cosh x$$

$$3. \sinh ix = i \sin x$$


$$4. \sin ix = i \sinh x$$

## 1 Functions of complex variables

Suppose, to each value that a complex variable  $z$  can assume, there corresponds one or more values of a complex variable  $w$ . We then say that  $w$  is a function of  $z$  and write  $w = f(z)$ .

- The complex variable  $z$  is called an *independent variable*
- The complex variable  $w$  is called a *dependent variable*.

### 1.1 Complex derivatives



If a complex function  $f(z)$  is single-valued in some region  $\mathcal{R}$  of the  $z$  plane, the derivative of  $f(z)$  is defined as

$$f'(z) = \lim_{\Delta z \rightarrow 0} \frac{\Delta w}{\Delta z} = \lim_{\Delta z \rightarrow 0} \frac{f(z + \Delta z) - f(z)}{\Delta z}$$

provided that the limit exists independent of the manner in which  $\Delta z$  approaches 0. In such a case, we say that  $f(z)$  is *differentiable* at  $z$  and  $f'(z)$  is known as the *derivative* of the function  $f(z)$ .

## 1.2 Complex analytic functions

If the derivative  $f'(z)$  exists at all points  $z$  of a region  $\mathcal{R}$ , then  $f(z)$  is said to be analytic in  $\mathcal{R}$  and is referred to as an *analytic function* in  $\mathcal{R}$ .

A function  $f(z)$  is said to be analytic at a point  $z_0$  if there exists a neighborhood  $|z - z_0| < \delta$  at all points of which  $f'(z)$  exists.

**Note:** Analytic functions are sometimes also known as *regular* or *holomorphic* or *monogenic* functions.

## 1.3 Cauchy-Riemann conditions

Given a complex function which can be written as follows:

$$w = f(z) = u(x, y) + iv(x, y)$$

and is analytic over a region  $\mathcal{R}$  on the  $z$ -plane, then the 2D functions  $u(x, y)$  and  $v(x, y)$  satisfy the following Cauchy-Riemann conditions in  $\mathcal{R}$ :

$$\boxed{\frac{\partial u}{\partial x} = \frac{\partial v}{\partial y} \quad \text{and} \quad \frac{\partial u}{\partial y} = -\frac{\partial v}{\partial x}}$$

If the partial derivatives of  $u$  and  $v$  are continuous in  $\mathcal{R}$ , then the Cauchy-Riemann equations are sufficient conditions that  $f(z)$  be analytic in  $\mathcal{R}$ .

**Note:** The functions  $u(x, y)$  and  $v(x, y)$  are known as *conjugate functions*.

### Theorem 1

The necessary conditions for a function  $f(z) = u(x, y) + iv(x, y)$  to be analytic at all points in a region  $\mathcal{R}$  are

$$\text{a) } \frac{\partial u}{\partial x} = \frac{\partial v}{\partial y}$$

$$\text{b) } \frac{\partial u}{\partial y} = -\frac{\partial v}{\partial x}$$

provided that  $\frac{\partial u}{\partial x}$ ,  $\frac{\partial u}{\partial y}$ ,  $\frac{\partial v}{\partial x}$  and  $\frac{\partial v}{\partial y}$  exist.

### Theorem 2

The sufficient conditions for a function  $f(z) = u(x, y) + iv(x, y)$  to be analytic at all points in a region  $\mathcal{R}$  are

- a)  $\frac{\partial u}{\partial x} = \frac{\partial v}{\partial y}$  and  $\frac{\partial u}{\partial y} = -\frac{\partial v}{\partial x}$
- b)  $\frac{\partial u}{\partial x}$ ,  $\frac{\partial u}{\partial y}$ ,  $\frac{\partial v}{\partial x}$  and  $\frac{\partial v}{\partial y}$  are continuous functions of  $x$  and  $y$  in the region  $\mathcal{R}$ .

### Theorem 3

Given a function  $f(z) = u(x, y) + iv(x, y)$ , which is analytic over a region  $\mathcal{R}$ , its derivative is obtained as

$$f'(z) = \frac{df}{dz} = \frac{\partial u}{\partial x} + i\frac{\partial v}{\partial x} = \frac{\partial v}{\partial y} - i\frac{\partial u}{\partial y}$$

## 1.4 Examples of analyticity of complex functions

1. Determine whether  $f(z) = \frac{1}{z}$  is analytic or not.

**Solution:** Let us consider

$$\begin{aligned} f(z) &= \frac{1}{z} \\ &= \frac{1}{x + iy} \\ &= \left( \frac{1}{x + iy} \right) \times \left( \frac{x - iy}{x - iy} \right) \\ &= \frac{x - iy}{(x + iy)(x - iy)} \\ &= \frac{x - iy}{x^2 + y^2} \\ \Rightarrow f(z) &= \left( \frac{x}{x^2 + y^2} \right) + i \left( \frac{-y}{x^2 + y^2} \right) \\ &= u(x, y) + iv(x, y) \end{aligned}$$

This gives us

$$u(x, y) = \frac{x}{x^2 + y^2} \quad \text{and} \quad v(x, y) = \frac{-y}{x^2 + y^2}$$

We obtain the partial derivatives as

$$\begin{aligned} \frac{\partial u}{\partial x} &= \frac{(x^2 + y^2) \times 1 - x \times 2x}{(x^2 + y^2)^2} \\ &= \frac{x^2 + y^2 - 2x^2}{(x^2 + y^2)^2} \\ &= \frac{y^2 - x^2}{(x^2 + y^2)^2} \end{aligned} \qquad \begin{aligned} \frac{\partial u}{\partial y} &= \frac{(x^2 + y^2) \times 0 - x \times 2y}{(x^2 + y^2)^2} \\ &= \frac{-2xy}{(x^2 + y^2)^2} \end{aligned}$$

$$\begin{aligned}
\frac{\partial v}{\partial x} &= \frac{(x^2 + y^2) \times 0 - (-y) \times 2x}{(x^2 + y^2)^2} & \frac{\partial v}{\partial y} &= \frac{(x^2 + y^2) \times (-1) - (-y) \times 2y}{(x^2 + y^2)^2} \\
&= \frac{2xy}{(x^2 + y^2)^2} & &= \frac{2y^2 - x^2 - y^2}{(x^2 + y^2)^2} \\
& & &= \frac{y^2 - x^2}{(x^2 + y^2)^2}
\end{aligned}$$

From the above partial derivatives, we find that

$$\frac{\partial u}{\partial x} = \frac{\partial v}{\partial y} \quad \text{and} \quad \frac{\partial u}{\partial y} = -\frac{\partial v}{\partial x}$$

Hence the Cauchy-Riemann conditions are satisfied. Also the partial derivatives  $\frac{\partial u}{\partial x}$ ,  $\frac{\partial u}{\partial y}$ ,  $\frac{\partial v}{\partial x}$  and  $\frac{\partial v}{\partial y}$  are continuous everywhere except at  $(0, 0)$ .

Therefore the complex function  $f(z) = \frac{1}{z}$  is analytic everywhere except at  $z = 0$ .

2. Show that the function  $f(z) = e^x(\cos y + i \sin y)$  is analytic, and then find its derivative.

**Solution:** Let us consider

$$\begin{aligned}
f(z) &= e^x(\cos y + i \sin y) \\
\Rightarrow f(z) &= (e^x \cos y) + i(e^x \sin y) \\
&= u(x, y) + iv(x, y)
\end{aligned}$$

This gives us

$$u(x, y) = e^x \cos y \quad \text{and} \quad v(x, y) = e^x \sin y$$

We obtain the partial derivatives as

$$\begin{aligned}
\frac{\partial u}{\partial x} &= \frac{\partial}{\partial x}(e^x \cos y) & \frac{\partial u}{\partial y} &= \frac{\partial}{\partial y}(e^x \cos y) \\
&= e^x \cos y & &= -e^x \sin y \\
\\ 
\frac{\partial v}{\partial x} &= \frac{\partial}{\partial x}(e^x \sin y) & \frac{\partial v}{\partial y} &= \frac{\partial}{\partial y}(e^x \sin y) \\
&= e^x \sin y & &= e^x \cos y
\end{aligned}$$

From the above partial derivatives, we find that

$$\frac{\partial u}{\partial x} = \frac{\partial v}{\partial y} \quad \text{and} \quad \frac{\partial u}{\partial y} = -\frac{\partial v}{\partial x}$$

Hence the Cauchy-Riemann conditions are satisfied. Also the partial derivatives  $\frac{\partial u}{\partial x}$ ,  $\frac{\partial u}{\partial y}$ ,  $\frac{\partial v}{\partial x}$  and  $\frac{\partial v}{\partial y}$  are continuous everywhere.

Therefore the complex function  $f(z) = e^x(\cos y + i \sin y)$  is analytic everywhere.

Finally, the derivative of the given function is obtained as

$$\begin{aligned} f'(z) &= \frac{\partial u}{\partial x} + i \frac{\partial v}{\partial x} \\ &= e^x \cos y + i e^x \sin y \\ \implies f'(z) &= e^x(\cos y + i \sin y) \end{aligned}$$

3. Test the analyticity of the function  $w = \sin z$  and thereby show that  $\frac{d}{dz}(\sin z) = \cos z$ .

**Solution:** We have

$$\begin{aligned} w &= \sin z \\ &= \sin(x + iy) \\ &= \sin x \cos iy + \cos x \sin iy \\ &= \sin x(\cosh y) + \cos x(i \sinh y) \\ \implies w &= f(z) = \sin x \cosh y + i \cos x \sinh y \end{aligned}$$

This gives us

$$u(x, y) = \sin x \cosh y \quad \text{and} \quad v(x, y) = \cos x \sinh y$$

The partial derivatives are

$$\begin{aligned} \frac{\partial u}{\partial x} &= \frac{\partial}{\partial x}(\sin x \cosh y) & \frac{\partial u}{\partial y} &= \frac{\partial}{\partial y}(\sin x \cosh y) \\ &= \cos x \cosh y & &= \sin x \sinh y \\ \frac{\partial v}{\partial x} &= \frac{\partial}{\partial x}(\cos x \sinh y) & \frac{\partial v}{\partial y} &= \frac{\partial}{\partial y}(\cos x \sinh y) \\ &= -\sin x \sinh y & &= \cos x \cosh y \end{aligned}$$

From the above partial derivatives, we find that

$$\frac{\partial u}{\partial x} = \frac{\partial v}{\partial y} \quad \text{and} \quad \frac{\partial u}{\partial y} = -\frac{\partial v}{\partial x}$$

Hence the Cauchy-Riemann conditions are satisfied. Also the partial derivatives  $\frac{\partial u}{\partial x}$ ,  $\frac{\partial u}{\partial y}$ ,  $\frac{\partial v}{\partial x}$  and  $\frac{\partial v}{\partial y}$  are continuous everywhere.

Therefore the complex function  $f(z) = \sin z$  is analytic everywhere.

Now, we evaluate the derivative as follows

$$\begin{aligned} \frac{d}{dz}(\sin z) &= \frac{\partial u}{\partial x} + i \frac{\partial v}{\partial x} \\ &= \cos x \cosh y + i(-\sin x \sinh y) \\ &= \cos x \cosh y - \sin x(i \sinh y) \\ &= \cos x(\cos iy) - \sin x(\sin iy) \\ &= \cos(x + iy) \\ \implies \frac{d}{dz}(\sin z) &= \cos z \end{aligned}$$



4. Find the values of  $C_1$  and  $C_2$  such that the function

$$f(z) = x^2 + C_1 y^2 - 2xy + i(C_2 x^2 - y^2 + 2xy)$$

is analytic. Also show that  $f'(z) = 2(1+i)z$ .

**Solution:** The given function is

$$f(z) = x^2 + C_1 y^2 - 2xy + i(C_2 x^2 - y^2 + 2xy)$$

from which we obtain

$$u(x, y) = x^2 + C_1 y^2 - 2xy \quad \text{and} \quad v(x, y) = C_2 x^2 - y^2 + 2xy$$

For the given function to be analytic, it should satisfy the Cauchy-Riemann conditions, i.e.

$$\begin{aligned} \frac{\partial u}{\partial x} &= \frac{\partial v}{\partial y} \\ \implies \frac{\partial}{\partial x}(x^2 + C_1 y^2 - 2xy) &= \frac{\partial}{\partial y}(C_2 x^2 - y^2 + 2xy) \\ \implies 2x - 2y &= -2y + 2x \end{aligned}$$

and

$$\begin{aligned} \frac{\partial u}{\partial y} &= -\frac{\partial v}{\partial x} \\ \implies \frac{\partial}{\partial y}(x^2 + C_1 y^2 - 2xy) &= -\frac{\partial}{\partial x}(C_2 x^2 - y^2 + 2xy) \\ \implies 2C_1 y - 2x &= -(2C_2 x + 2y) \\ \implies 2C_1 y - 2x &= -2y - 2C_2 x \end{aligned}$$

Comparing the coefficients of  $y$  and  $x$  above, we obtain

$$\begin{aligned} 2C_1 &= -2 \quad \text{and} \quad -2C_2 = -2 \\ \implies C_1 &= -1 \quad \text{and} \quad C_2 = 1 \end{aligned}$$

Hence the given function is

$$f(z) = x^2 - y^2 - 2xy + i(x^2 - y^2 + 2xy)$$

Now, we have

$$\begin{aligned} \frac{\partial u}{\partial y} &= 2C_1 y - 2x \\ \implies \frac{\partial u}{\partial y} &= -2y - 2x \quad \text{and} \quad \frac{\partial v}{\partial x} = -2y + 2x \end{aligned}$$

Therefore, the derivative is

$$\begin{aligned} f'(z) &= \frac{\partial v}{\partial y} - i \frac{\partial u}{\partial x} \\ \implies f'(z) &= -2y + 2x - i(-2y - 2x) \\ &= -2y + 2x + 2i(x + y) \\ &= 2[(x + iy) + (-y + ix)] \\ &= 2[(1+i)x + i(1+i)y] \\ &= 2(1+i)(x + iy) \\ \implies f'(z) &= 2(1+i)z \end{aligned}$$

5. Obtain the Cauchy-Riemann conditions in polar form.

**Solution:** The polar representation of a complex number is

$$z = x + iy = re^{i\theta} \quad (1)$$

Differentiating equation (1) with respect to  $r$  and  $\theta$  give us

$$\frac{\partial z}{\partial r} = e^{i\theta} \quad (2)$$

$$\frac{\partial z}{\partial \theta} = ire^{i\theta} \quad (3)$$

Let us now consider an analytic function given as

$$f(z) = u + iv \quad (4)$$

Differentiating equation (4) with respect to  $r$  gives us

$$\begin{aligned} \frac{\partial}{\partial r}[f(z)] &= \frac{\partial}{\partial r}(u + iv) \\ \Rightarrow \frac{df}{dz} \frac{\partial z}{\partial r} &= \frac{\partial u}{\partial r} + i \frac{\partial v}{\partial r} \\ \Rightarrow f'(z)e^{i\theta} &= \frac{\partial u}{\partial r} + i \frac{\partial v}{\partial r} \quad [\text{using eqn. (2) on LHS}] \end{aligned}$$

Again, differentiating equation (4) with respect to  $\theta$  gives us

$$\begin{aligned} \frac{\partial}{\partial \theta}[f(z)] &= \frac{\partial}{\partial \theta}(u + iv) \\ \Rightarrow \frac{df}{dz} \frac{\partial z}{\partial \theta} &= \frac{\partial u}{\partial \theta} + i \frac{\partial v}{\partial \theta} \\ \Rightarrow irf'(z)e^{i\theta} &= \frac{\partial u}{\partial \theta} + i \frac{\partial v}{\partial \theta} \quad [\text{using eqn. (3) on LHS}] \\ \Rightarrow ir \left[ \frac{\partial u}{\partial r} + i \frac{\partial v}{\partial r} \right] &= \frac{\partial u}{\partial \theta} + i \frac{\partial v}{\partial \theta} \\ \Rightarrow ir \frac{\partial u}{\partial r} + i^2 r \frac{\partial v}{\partial r} &= \frac{\partial u}{\partial \theta} + i \frac{\partial v}{\partial \theta} \\ \Rightarrow -r \frac{\partial v}{\partial r} + ir \frac{\partial u}{\partial r} &= \frac{\partial u}{\partial \theta} + i \frac{\partial v}{\partial \theta} \end{aligned} \quad (5)$$

Equating the imaginary and real parts on either sides of equation (5), we obtain

$$\begin{aligned} r \frac{\partial u}{\partial r} &= \frac{\partial v}{\partial \theta} & -r \frac{\partial v}{\partial r} &= \frac{\partial u}{\partial \theta} \\ \Rightarrow \frac{\partial u}{\partial r} &= \frac{1}{r} \frac{\partial v}{\partial \theta} & \Rightarrow \frac{\partial u}{\partial \theta} &= -r \frac{\partial v}{\partial r} \end{aligned}$$

Therefore, the Cauchy-Riemann conditions in polar coordinates are

$$\boxed{\begin{aligned} \frac{\partial u}{\partial r} &= \frac{1}{r} \frac{\partial v}{\partial \theta} \\ \frac{\partial u}{\partial \theta} &= -r \frac{\partial v}{\partial r} \end{aligned}}$$

## 2 Singularities of complex functions

A point at which a complex function  $f(z)$  fails to be analytic is called a *singular point* or *singularity* of  $f(z)$ . Various types of singularities exist, which are discussed below:

### 2.1 Isolated singularities

A point  $z = z_0$  is called an *isolated singularity* or *isolated singular point* of a function  $f(z)$  if we can find  $\delta > 0$  such that the circle  $|z - z_0| = \delta$  encloses no singular point other than  $z_0$ .

- If no such  $\delta$  can be found, we call  $z_0$  a *non-isolated singularity*.
- If  $z_0$  is not a singular point and we can find  $\delta > 0$  such that  $|z - z_0| = \delta$  encloses no singular point, then we call  $z_0$  an *ordinary point* of  $f(z)$ .

#### Examples

1.  $f(z) = \frac{1}{z(z-2)}$  has isolated singularities at  $z_0 = 0$  and  $z_0 = 2$ .
2.  $f(z) = e^{1/z}$  has an isolated singularity at  $z = 0$ .

### 2.2 Poles

If  $z_0$  is an isolated singularity and we can find a positive integer  $n$  such that

$$\lim_{z \rightarrow z_0} (z - z_0)^n f(z) \neq 0$$


then  $z = z_0$  is called a *pole* of order  $n$ .

- Special case: If  $n = 1$ ,  $z_0$  is called a *simple pole*.
- If  $g(z) = (z - z_0)^n f(z)$ , where  $f(z_0) \neq 0$  and  $n$  is a positive integer, then  $z = z_0$  is called a *zero* of order  $n$  of  $g(z)$ . In such a case,  $z_0$  is a pole of order  $n$  of the function  $\frac{1}{g(z)}$ .
  - If  $n = 1$ ,  $z_0$  is called a *simple zero*.

#### Examples

1.  $f(z) = \frac{1}{(z-2)^3}$  has a pole of order 3 at  $z_0 = 2$ .
2.  $f(z) = \frac{3z-2}{(z-1)^2(z+5i)(z-4)}$  has a pole of order 2 at  $z_0 = 1$  and simple poles at  $z_0 = -5i$  and  $z_0 = 4$ .

## 2.3 Branch points



Suppose that we are given the function  $w = z^{1/2}$ . Further, we allow  $z$  to make a complete circuit (counterclockwise) around the origin starting from point  $A$ , as shown in the figure above.

We have  $z = re^{i\theta}$ , and so  $w = (re^{i\theta})^{1/2} = \sqrt{r}e^{i\theta/2}$ . Then at point  $A$ ,  $\theta = \theta_1$  and  $w = \sqrt{r}e^{i\theta_1/2}$ . After a complete circuit back to  $A$ ,  $\theta = \theta_1 + 2\pi$  and so

$$\begin{aligned} w &= \sqrt{r}e^{i(\theta_1+2\pi)/2} \\ &= \sqrt{r}e^{i\theta_1/2}e^{i\pi} \\ \implies w &= -\sqrt{r}e^{i\theta_1/2} \end{aligned}$$

Thus, we have not achieved the same value of  $w$  with which we started. However, by making a second complete circuit back to  $A$ , i.e.,  $\theta = \theta_1 + 4\pi$ , we have

$$\begin{aligned} w &= \sqrt{r}e^{i(\theta_1+4\pi)/2} \\ &= \sqrt{r}e^{i\theta_1/2}e^{i2\pi} \\ \implies w &= \sqrt{r}e^{i\theta_1/2} \end{aligned}$$

and we then do obtain the same value of  $w$  with which we started.

We can describe the above by stating that if  $0 \leq \theta < 2\pi$ , we are on one branch of the multi-valued function  $z^{1/2}$ , while if  $2\pi \leq \theta < 4\pi$ , we are on the other branch of the function.

It is clear that each branch of the function is single-valued. In order to keep the function single-valued, we set up an artificial barrier such as  $OB$  where  $B$  is at infinity which we agree not to cross.

This barrier is called a *branch line* or *branch cut*, and point  $O$  is called a *branch point*.

### Examples

1.  $f(z) = (z - 3)^{1/2}$  has a branch point at  $z_0 = 3$ .
2.  $f(z) = \ln(z^2 + z - 2)$  has branch points where  $z^2 + z - 2 = 0$ , i.e. at  $z_0 = 1$  and  $z_0 = -2$ .

## 2.4 Removable singularities

An isolated singular point  $z_0$  is called a *removable singularity* of  $f(z)$  if  $\lim_{z \rightarrow z_0} f(z)$  exists.

By defining  $f(z_0) = \lim_{z \rightarrow z_0} f(z)$ , it can then be shown that  $f(z)$  is not only continuous at  $z_0$  but is also analytic at  $z_0$ .

### Examples

1.  $f(z) = \frac{\sin z}{z}$  has a removable singularity at  $z_0 = 0$  because  $\lim_{z \rightarrow 0} \frac{\sin z}{z} = 1$ , which allows us to define  $f(0) = 1$ .
2.  $f(z) = \frac{e^{z-1} - 1}{z - 1}$  has a removable singularity at  $z_0 = 1$  because  $\lim_{z \rightarrow 1} \frac{e^{z-1} - 1}{z - 1} = 1$ , which allows us to define  $f(1) = 1$ .

## 2.5 Essential singularities

An isolated singularity that is not a pole or removable singularity is called an *essential singularity*.

### Examples

1.  $f(z) = e^{1/(z-2)}$  has an essential singularity at  $z_0 = 2$ .
2.  $f(z) = \sin\left(\frac{1}{z-i}\right)$  has an essential singularity at  $z_0 = i$ .


### 3 Complex integration

#### 3.1 Simply and multiply connected regions

A region  $\mathcal{R}$  is called *simply-connected* if any simple closed curve, which lies in  $\mathcal{R}$ , can be shrunk to a point without leaving  $\mathcal{R}$ .


A region  $\mathcal{R}$ , which is not simply-connected, is called *multiply-connected*.

##### 3.1.1 Simply connected region




Suppose  $\mathcal{R}$  is the region defined by  $|z| < 2$  shown shaded in the above figure. If  $\Gamma$  is any simple closed curve lying in  $\mathcal{R}$ , i.e. whose points are in  $\mathcal{R}$ , we see that it can be shrunk to a point that lies in  $\mathcal{R}$ , and thus does not leave  $\mathcal{R}$ , so that  $\mathcal{R}$  is simply-connected.

##### 3.1.2 Multiply connected region



If  $\mathcal{R}$  is the region defined by  $1 < |z| < 2$ , shown shaded in the above figure, then there is a

simple closed curve  $\Gamma$  lying in  $\mathcal{R}$  that cannot possibly be shrunk to a point without leaving  $\mathcal{R}$ , so that  $\mathcal{R}$  is multiply-connected.



Intuitively, a simply-connected region is one that does not have any “holes” in it, while a multiply-connected region is one that does. The multiply-connected region of the above figure has three holes and, hence, is said to be triply-connected.


### 3.2 Cauchy’s integral theorem

Let  $f(z)$  be analytic in a region  $\mathcal{R}$  and on its boundary  $C$ . Then

$$\oint_C f(z) \, dz = 0$$

This fundamental theorem, known as *Cauchy’s integral theorem*, is valid for both simply- and multiply-connected regions.

### 3.3 Cauchy’s integral formulas



Let  $f(z)$  be analytic inside and on a simple closed curve  $C$  and let  $a$  be any point inside  $C$ . Then

$$f(a) = \frac{1}{2\pi i} \oint_C \frac{f(z)}{z - a} \, dz$$

where  $C$  is traversed in the positive (counterclockwise) sense.

Also, the  $n^{\text{th}}$  derivative of  $f(z)$  at  $z = a$  is given by

$$f^{(n)}(a) = \frac{n!}{2\pi i} \oint_C \frac{f(z)}{(z-a)^{n+1}} dz, \quad n = 1, 2, 3, \dots$$



## **4 Complex series**

### **4.1 Taylor's series expansion of a complex function**

### **4.2 Laurent's series expansion of a complex function**

#### **4.2.1 Residue theorem**