Draft version October 9, 2026
Typeset using LaTeX **default** style in AASTeX7.0.1

Two Earth-sized Transiting Planet Candidates around Nearby K Dwarfs from a Multi-sector Search of TESS Full-frame Images

Paras Chopra[1]

[1] *Independent researcher*

ABSTRACT

We report two transit-like signals from a multi-sector box least squares search of TESS full-frame-image light curves of 526 nearby (< 60 pc) dwarf stars never observed at 2-minute cadence. TIC 370612437.01 has $P = 3.564921$ d, a depth of 210 ppm and an implied radius of $1.04 \pm 0.15\,R_\oplus$. TIC 252721481.01 (HIP 114417) has $P = 4.933533$ d, a depth of 279 ppm and an implied radius of $1.10 \pm 0.14\,R_\oplus$. Both signals appear in the SPOC and QLP reductions, show no secondary eclipse or odd–even difference, and give local depth-to-noise ratios of about 7–8. Neither source is localized, so blended eclipsing binaries are not excluded. We provide ephemerides and preregistered transit predictions for the 2027–2028 TESS sectors.

Keywords: Exoplanet detection methods (489) — Transit photometry (1709) — Exoplanets (498)

1. SEARCH

We searched TESS (G. R. Ricker et al. 2015) data for stars from the Gaia Catalogue of Nearby Stars (Gaia Collaboration et al. 2021) within 60 pc, matched to the TESS Input Catalog v8.2 (K. G. Stassun et al. 2019). We kept stars with $R_\star \leq 1.5\,R_\odot$, $T \leq 13$ and contamination ratio < 0.2, observed in at least three sectors spanning at least two years, never observed at 2-minute cadence, and not already a TOI, CTOI or confirmed-planet host. This left 526 stars.

We used one light curve per sector: TESS-SPOC full-frame-image PDCSAP photometry (D. A. Caldwell et al. 2020) where available, otherwise QLP (C. X. Huang et al. 2020). Each light curve was detrended with a 0.75 d biweight filter (M. Hippke et al. 2019), chosen by injection–recovery tests. We then ran an iterative box least squares search (G. Kovács et al. 2002) over 0.5–25 d with up to three signals per star. Signals with SDE ≥ 7, SNR ≥ 6 and at least three transits (743 in total) were triaged automatically. The flags tested odd–even depth differences, secondary eclipses, inconsistent depths between sectors, duration, the dominance of single events, edge effects, and periods shared across stars (cf. J. D. Twicken et al. 2018), using noise measured from 40 off-transit phases. Flag thresholds were calibrated on injected planets.

Thirty-one signals had no hard flags. The two reported here have the highest SDE (11.3 and 10.3). As an empirical null, we inverted all 526 light curves and repeated the identical search and triage (cf. S. E. Thompson et al. 2018). No flag-free inverted signal exceeded SDE 8.4. This is a single null realization, so it does not give a calibrated false-alarm probability.

2. CANDIDATES

TIC 370612437.01 (TYC 3968-348-1; $T = 9.80$, $T_{\rm eff} = 4380$ K, $R_\star = 0.66\,R_\odot$, $d = 48.4$ pc):

- $P = 3.564921^{+0.000022}_{-0.000018}$ d and $T_0 = 2458742.1409 \pm 0.0069$ BJD$_{\rm TDB}$, with 55 transit epochs covered in eight sectors (16, 17, 56, 57, 76, 77, 83, 84).
- A batman (L. Kreidberg 2015)/emcee (D. Foreman-Mackey et al. 2013) fit gives depth 210^{+45}_{-49} ppm, $T_{14} = 1.44^{+0.28}_{-0.14}$ h, $b = 0.73^{+0.08}_{-0.18}$ and $R_p = 1.04 \pm 0.15\,R_\oplus$. The fit integrates over exposures (D. M. Kipping 2010), uses A. Claret (2017) limb darkening, inflates the errors for red noise ($\beta = 1.18$) and applies a TIC density prior.
- The equilibrium temperature is $T_{\rm eq} \approx 850$ K.

TIC 252721481.01 (HIP 114417; $T = 9.73$, $T_{\rm eff} = 4301\,{\rm K}$, $R_\star = 0.61\,R_\odot$, $d = 41.1\,{\rm pc}$):

- $P = 4.933533^{+0.000019}_{-0.000021}$ d and $T_0 = 2458740.6616 \pm 0.0060\,{\rm BJD_{TDB}}$, with 21 transit epochs covered in five sectors (16, 17, 57, 77, 84).
- The fit gives depth 279^{+52}_{-48} ppm, $T_{14} = 1.59 \pm 0.15\,{\rm h}$, $b = 0.68^{+0.10}_{-0.14}$ and $R_p = 1.10^{+0.14}_{-0.13}\,R_\oplus$.
- The equilibrium temperature is $T_{\rm eq} \approx 720\,{\rm K}$.

Figure 1 shows the folded light curves.

3. VETTING

For both candidates:

- **Robustness.** The signals persist with transit-masked detrending and alternative detrending choices. Measured independently, the box depths are 208 ± 29 and 263 ± 34 ppm, with errors from matched-processing off-transit boxes.
- **Second reduction.** They appear in QLP, a second reduction of the same images: 225 ± 35 and 238 ± 71 ppm.
- **Retrospective season checks.** In leave-one-season-out tests, the dip appears at the predicted epoch in three of four held-out seasons for each candidate. For TIC 370612437, the 2022 season (sectors 56–57) is weaker, at about 105 ± 45 ppm. For HIP 114417, the data without 2019 are insufficient to recover the period.
- **Eclipsing-binary signatures.** Neither shows a secondary eclipse at phase 0.5, and the odd and even depths agree within 2σ.
- **Stellar density.** The density implied by the transit shape alone is weakly constrained but consistent with the TIC value.
- **Companions and prior reports.** Gaia DR3 (Gaia Collaboration et al. 2023) RUWE values (0.89, 1.03) and the Hipparcos–Gaia accelerations (T. D. Brandt 2021) show no sign of a bound companion. Neither star appears among the 341,351 SPOC TCEs, in the TESS-SPOC FFI DV manifests we checked, or in ExoFOP.

The main caveat is that the source is not localized. Difference images built from TESScut (C. E. Brasseur et al. 2019) show positive target-position amplitudes in every sector, but they cannot exclude faint neighbours within about one pixel (21″). TIC 370612437 lies at $b_{\rm gal} = -1.3°$ in a crowded field, where Gaia sources within 12″ would need eclipse depths of only 3–5%. A small detector-coordinate centroid shift in HIP 114417 is not significant once the spacecraft roll between sectors is taken into account. TRICERATOPS validation (S. Giacalone et al. 2021) could not be completed.

We therefore present both as unvalidated planet candidates. Seeing-limited photometry of nearby stars at the predicted epochs, together with high-resolution imaging, would test the blend scenarios. TESS is scheduled to reobserve TIC 370612437 in sectors 121, 122 and 129, and HIP 114417 in sectors 122 and 129. We published predicted transit times before these observations (P. Chopra 2026).

ACKNOWLEDGMENTS

This note uses TESS data from MAST (STScI) and Gaia data from ESA. The search and vetting code was written with the assistance of an AI coding agent (Claude Code, Anthropic), and independent code reviews were performed with OpenAI Codex. The author checked the analysis and is responsible for its content. Code, intermediate results and review reports are available on request.

Facilities: TESS, Gaia

Software: astropy (Astropy Collaboration et al. 2022), lightkurve (Lightkurve Collaboration 2018), wotan (M. Hippke et al. 2019), batman (L. Kreidberg 2015), emcee (D. Foreman-Mackey et al. 2013)

REFERENCES

Astropy Collaboration, Price-Whelan, A. M., et al. 2022, ApJ, 935, 167, doi: 10.3847/1538-4357/ac7c74

Brandt, T. D. 2021, ApJS, 254, 42, doi: 10.3847/1538-4365/abf93c




Figure 1. TESS-SPOC 200 s photometry from sectors 56–84 (gray), folded on each ephemeris, with 15-minute bins (black) and the best-fitting transit model (red).

Brasseur, C. E., Phillip, C., Fleming, S. W., Mullally, S. E., & White, R. L. 2019, Astrocut: Tools for creating cutouts of TESS images,, Astrophysics Source Code Library, ascl:1905.007

Caldwell, D. A., et al. 2020, Research Notes of the AAS, 4, 201, doi: 10.3847/2515-5172/abc9b3

Chopra, P. 2026, Predicted transit times for two TESS planet candidates: TIC 370612437 and HIP 114417,, Zenodo doi: 10.5281/zenodo.23257784

Claret, A. 2017, A&A, 600, A30, doi: 10.1051/0004-6361/201629705

Foreman-Mackey, D., Hogg, D. W., Lang, D., & Goodman, J. 2013, PASP, 125, 306, doi: 10.1086/670067

Gaia Collaboration, Smart, R. L., et al. 2021, A&A, 649, A6, doi: 10.1051/0004-6361/202039498

Gaia Collaboration, Vallenari, A., et al. 2023, A&A, 674, A1, doi: 10.1051/0004-6361/202243940

Giacalone, S., et al. 2021, AJ, 161, 24, doi: 10.3847/1538-3881/abc6af

Hippke, M., David, T. J., Mulders, G. D., & Heller, R. 2019, AJ, 158, 143, doi: 10.3847/1538-3881/ab3984

Huang, C. X., et al. 2020, Research Notes of the AAS, 4, 204, doi: 10.3847/2515-5172/abca2e

Kipping, D. M. 2010, MNRAS, 408, 1758, doi: 10.1111/j.1365-2966.2010.17242.x

Kovács, G., Zucker, S., & Mazeh, T. 2002, A&A, 391, 369, doi: 10.1051/0004-6361:20020802

Kreidberg, L. 2015, PASP, 127, 1161, doi: 10.1086/683602

Lightkurve Collaboration. 2018, Lightkurve: Kepler and TESS time series analysis in Python,, Astrophysics Source Code Library, ascl:1812.013

Ricker, G. R., et al. 2015, Journal of Astronomical Telescopes, Instruments, and Systems, 1, 014003, doi: 10.1117/1.JATIS.1.1.014003

Stassun, K. G., et al. 2019, AJ, 158, 138, doi: 10.3847/1538-3881/ab3467

Thompson, S. E., et al. 2018, ApJS, 235, 38, doi: 10.3847/1538-4365/aab4f9

Twicken, J. D., et al. 2018, PASP, 130, 064502, doi: 10.1088/1538-3873/aab694