


Chapter 5. Electron-Positron annihilation




QED Calculations

- How to calculate a cross section using QED (e.g. $e^+e^- \rightarrow \mu^+\mu^-$):


— Draw all possible Feynman Diagrams

- For $e^+e^- \rightarrow \mu^+\mu^-$ there is just one lowest order diagram



$$M \propto e^2 \propto \alpha_{em}$$

+ many **second order** diagrams + ...



For each diagram calculate the matrix element using Feynman rules derived in handout 4.

Sum the individual matrix elements (i.e. sum the amplitudes)



$$M_{fi} = M_1 + M_2 + M_3 + \dots$$

- Note: summing amplitudes therefore different diagrams for the same final state can interfere either positively or negatively!

and then square $|M_{fi}|^2 = (M_1 + M_2 + M_3 + \dots)(M_1^* + M_2^* + M_3^* + \dots)$

➡ this gives the full perturbation expansion in α_{em}

- For QED $\alpha_{em} \sim 1/137$ the lowest order diagram dominates and for most purposes it is sufficient to **neglect** higher order diagrams.



= Calculate decay rate/cross section using formulae from handout 1.

- e.g. for a decay

$$\Gamma = \frac{p^*}{32\pi^2 m_a^2} \int |M_{fi}|^2 d\Omega$$

- For scattering in the centre-of-mass frame

$$\frac{d\sigma}{d\Omega^*} = \frac{1}{64\pi^2 s} \frac{|\vec{p}_f^*|}{|\vec{p}_i^*|} |M_{fi}|^2 \quad (1)$$

- For scattering in lab. frame (neglecting mass of scattered particle)

$$\frac{d\sigma}{d\Omega} = \frac{1}{64\pi^2} \left(\frac{E_3}{ME_1} \right)^2 |M_{fi}|^2$$


Electron Positron Annihilation

★ Consider the process: $e^+e^- \rightarrow \mu^+\mu^-$


- Work in C.o.M. frame (this is appropriate for most e^+e^- colliders).

$$p_1 = (E, 0, 0, p) \quad p_2 = (E, 0, 0, -p)$$

$$p_3 = (E, \vec{p}_f) \quad p_4 = (E, -\vec{p}_f)$$



- Only consider the lowest order Feynman diagram:



♦ Feynman rules give:

$$-iM = [\bar{v}(p_2)ie\gamma^\mu u(p_1)] \frac{-ig_{\mu\nu}}{q^2} [\bar{u}(p_3)ie\gamma^\nu v(p_4)]$$

NOTE:

- Incoming anti-particle \bar{v}
- Incoming particle u
- Adjoint spinor written first

- In the C.o.M. frame have

$$\frac{d\sigma}{d\Omega} = \frac{1}{64\pi^2 s} \frac{|\vec{p}_f|}{|\vec{p}_i|} |M_{fi}|^2 \quad \text{with} \quad s = (p_1 + p_2)^2 = (E + E)^2 = 4E^2$$

Electron and Muon Currents

- Here $q^2 = (p_1 + p_2)^2 = s$ and matrix element

$$-iM = [\bar{v}(p_2)ie\gamma^\mu u(p_1)] \frac{-ig_{\mu\nu}}{q^2} [\bar{u}(p_3)ie\gamma^\nu v(p_4)]$$

→
$$M = -\frac{e^2}{s} g_{\mu\nu} [\bar{v}(p_2)\gamma^\mu u(p_1)][\bar{u}(p_3)\gamma^\nu v(p_4)]$$

- In handout 2 introduced the **four-vector** current

$$j^\mu = \bar{\psi}\gamma^\mu\psi$$

which has same form as the two terms in [] in the matrix element

- The matrix element can be written in terms of the electron and muon currents

$$(j_e)^\mu = \bar{v}(p_2)\gamma^\mu u(p_1) \quad \text{and} \quad (j_\mu)^\nu = \bar{u}(p_3)\gamma^\nu v(p_4)$$


→
$$M = -\frac{e^2}{s} g_{\mu\nu} (j_e)^\mu (j_\mu)^\nu$$

$$M = -\frac{e^2}{s} j_e \cdot j_\mu$$

- Matrix element is a four-vector scalar product – confirming it is **Lorentz Invariant**

Spin in e^+e^- Annihilation

- In general the electron and positron will not be polarized, i.e. there will be equal numbers of positive and negative helicity states
- There are four possible combinations of spins in the **initial state** !



- Similarly there are four possible helicity combinations in the final state
- In total there are **16** combinations e.g. **RL**→**RR**, **RL**→**RL**, ...
- To account for these states we need to **sum over all 16 possible helicity combinations** and then **average** over the number of **initial helicity states**:

$$\langle |M|^2 \rangle = \frac{1}{4} \sum_{\text{spins}} |M_i|^2 = \frac{1}{4} (|M_{LL \rightarrow LL}|^2 + |M_{LL \rightarrow LR}|^2 + \dots)$$

- ★ i.e. need to evaluate:

$$M = -\frac{e^2}{s} j_e \cdot j_\mu$$

for all 16 helicity combinations !


- ★ Fortunately, in the limit $E \gg m_\mu$ only 4 helicity combinations give non-zero matrix elements – we will see that this is an important feature of QED/QCD

- In the C.o.M. frame in the limit $E \gg m$

$$p_1 = (E, 0, 0, E); \quad p_2 = (E, 0, 0, -E)$$

$$p_3 = (E, E \sin \theta, 0, E \cos \theta);$$

$$p_4 = (E, -E \sin \theta, 0, -E \cos \theta)$$



- Left- and right-handed helicity spinors (handout 3) for particles/anti-particles are:

$$u_{\uparrow} = N \begin{pmatrix} c \\ e^{i\phi} s \\ \frac{|\vec{p}|}{E+m} c \\ \frac{|\vec{p}|}{E+m} e^{i\phi} s \end{pmatrix} \quad u_{\downarrow} = N \begin{pmatrix} -s \\ e^{i\phi} c \\ \frac{|\vec{p}|}{E+m} s \\ -\frac{|\vec{p}|}{E+m} e^{i\phi} c \end{pmatrix} \quad v_{\uparrow} = N \begin{pmatrix} \frac{|\vec{p}|}{E+m} s \\ -\frac{|\vec{p}|}{E+m} e^{i\phi} c \\ -s \\ e^{i\phi} c \end{pmatrix} \quad v_{\downarrow} = N \begin{pmatrix} \frac{|\vec{p}|}{E+m} c \\ \frac{|\vec{p}|}{E+m} e^{i\phi} s \\ c \\ e^{i\phi} s \end{pmatrix}$$

where $s = \sin \frac{\theta}{2}$; $c = \cos \frac{\theta}{2}$ and $N = \sqrt{E+m}$

- In the limit $E \gg m$ these become:

$$u_{\uparrow} = \sqrt{E} \begin{pmatrix} c \\ se^{i\phi} \\ c \\ se^{i\phi} \end{pmatrix}; \quad u_{\downarrow} = \sqrt{E} \begin{pmatrix} -s \\ ce^{i\phi} \\ s \\ -ce^{i\phi} \end{pmatrix}; \quad v_{\uparrow} = \sqrt{E} \begin{pmatrix} s \\ -ce^{i\phi} \\ -s \\ ce^{i\phi} \end{pmatrix}; \quad v_{\downarrow} = \sqrt{E} \begin{pmatrix} c \\ se^{i\phi} \\ c \\ se^{i\phi} \end{pmatrix}$$

- The initial-state electron can either be in a left- or right-handed helicity state


$$u_{\uparrow}(p_1) = \sqrt{E} \begin{pmatrix} 1 \\ 0 \\ 1 \\ 0 \end{pmatrix}; \quad u_{\downarrow}(p_1) = \sqrt{E} \begin{pmatrix} 0 \\ 1 \\ 0 \\ -1 \end{pmatrix};$$

- For the initial state positron ($\theta = \pi$) can have either:

$$v_{\uparrow}(p_2) = \sqrt{E} \begin{pmatrix} 1 \\ 0 \\ -1 \\ 0 \end{pmatrix}; v_{\downarrow}(p_2) = \sqrt{E} \begin{pmatrix} 0 \\ 1 \\ 0 \\ 1 \end{pmatrix}$$

- Similarly for the final state μ^- which has polar angle θ and choosing $\phi = 0$

$$u_{\uparrow}(p_3) = \sqrt{E} \begin{pmatrix} c \\ s \\ c \\ s \end{pmatrix}; u_{\downarrow}(p_3) = \sqrt{E} \begin{pmatrix} -s \\ c \\ s \\ -c \end{pmatrix};$$




- And for the final state μ^+ replacing $\theta \rightarrow \pi - \theta$; $\phi \rightarrow \pi$ obtain

$$v_{\uparrow}(p_4) = \sqrt{E} \begin{pmatrix} c \\ s \\ -c \\ -s \end{pmatrix}; v_{\downarrow}(p_4) = \sqrt{E} \begin{pmatrix} s \\ -c \\ s \\ -c \end{pmatrix}; \quad \left\{ \begin{array}{l} \text{using} \\ \sin\left(\frac{\pi-\theta}{2}\right) = \cos \frac{\theta}{2} \\ \cos\left(\frac{\pi-\theta}{2}\right) = \sin \frac{\theta}{2} \\ e^{i\pi} = -1 \end{array} \right.$$

- Wish to calculate the matrix element $M = -\frac{e^2}{s} j_e \cdot j_{\mu}$

- ★ first consider the muon current j_{μ} for 4 possible helicity combinations



The Muon Current

- Want to evaluate $(j_\mu)^\nu = \bar{u}(p_3)\gamma^\nu v(p_4)$ for all four helicity combinations
- For arbitrary spinors ψ, ϕ with it is straightforward to show that the components of $\bar{\psi}\gamma^\mu\phi$ are

$$\bar{\psi}\gamma^0\phi = \psi^\dagger\gamma^0\gamma^0\phi = \psi_1^*\phi_1 + \psi_2^*\phi_2 + \psi_3^*\phi_3 + \psi_4^*\phi_4 \quad (3)$$

$$\bar{\psi}\gamma^1\phi = \psi^\dagger\gamma^0\gamma^1\phi = \psi_1^*\phi_4 + \psi_2^*\phi_3 + \psi_3^*\phi_2 + \psi_4^*\phi_1 \quad (4)$$

$$\bar{\psi}\gamma^2\phi = \psi^\dagger\gamma^0\gamma^2\phi = -i(\psi_1^*\phi_4 - \psi_2^*\phi_3 + \psi_3^*\phi_2 - \psi_4^*\phi_1) \quad (5)$$

$$\bar{\psi}\gamma^3\phi = \psi^\dagger\gamma^0\gamma^3\phi = \psi_1^*\phi_3 - \psi_2^*\phi_4 + \psi_3^*\phi_1 - \psi_4^*\phi_2 \quad (6)$$

- Consider the $\mu_R^- \mu_L^+$ combination using $\psi = u_\uparrow$ $\phi = v_\downarrow$

with $v_\downarrow = \sqrt{E} \begin{pmatrix} s \\ -c \\ s \\ -c \end{pmatrix}; u_\uparrow = \sqrt{E} \begin{pmatrix} c \\ s \\ c \\ s \end{pmatrix};$

$$\bar{u}_\uparrow(p_3)\gamma^0 v_\downarrow(p_4) = E(cs - sc + cs - sc) = 0$$

$$\bar{u}_\uparrow(p_3)\gamma^1 v_\downarrow(p_4) = E(-c^2 + s^2 - c^2 + s^2) = 2E(s^2 - c^2) = -2E \cos \theta$$

$$\bar{u}_\uparrow(p_3)\gamma^2 v_\downarrow(p_4) = -iE(-c^2 - s^2 - c^2 - s^2) = 2iE$$


$$\bar{u}_\uparrow(p_3)\gamma^3 v_\downarrow(p_4) = E(cs + sc + cs + sc) = 4Esc = 2E \sin \theta$$



- Hence the four-vector muon current for the **RL** combination is

$$\bar{u}_{\uparrow}(p_3)\gamma^{\nu}v_{\downarrow}(p_4) = 2E(0, -\cos\theta, i, \sin\theta)$$

- The results for the 4 helicity combinations (obtained in the same manner) are:



$$\bar{u}_{\uparrow}(p_3)\gamma^{\nu}v_{\downarrow}(p_4) = 2E(0, -\cos\theta, i, \sin\theta) \quad \text{RL}$$

$$\bar{u}_{\uparrow}(p_3)\gamma^{\nu}v_{\uparrow}(p_4) = (0, 0, 0, 0) \quad \text{RR}$$

$$\bar{u}_{\downarrow}(p_3)\gamma^{\nu}v_{\downarrow}(p_4) = (0, 0, 0, 0) \quad \text{LL}$$


$$\bar{u}_{\downarrow}(p_3)\gamma^{\nu}v_{\uparrow}(p_4) = 2E(0, -\cos\theta, -i, \sin\theta) \quad \text{LR}$$

★ **IN THE LIMIT $E \gg m$ only two helicity combinations are non-zero !**

- This is an important feature of QED. It applies equally to QCD.
- In the Weak interaction only one helicity combination contributes.
- The origin of this will be discussed in the last part of this lecture
- But as a consequence of the 16 possible helicity combinations only four given non-zero matrix elements

Electron Positron Annihilation cont.

- Previously we derived the muon currents for the allowed helicities:



$$\begin{aligned} \mu_R^- \mu_L^+ : \quad & \bar{u}_\uparrow(p_3) \gamma^\nu v_\downarrow(p_4) = 2E(0, -\cos \theta, i, \sin \theta) \\ \mu_L^- \mu_R^+ : \quad & \bar{u}_\downarrow(p_3) \gamma^\nu v_\uparrow(p_4) = 2E(0, -\cos \theta, -i, \sin \theta) \end{aligned}$$

- Now need to consider the electron current

The Electron Current

- The incoming electron and positron spinors (L and R helicities) are:

$$u_{\uparrow} = \sqrt{E} \begin{pmatrix} 1 \\ 0 \\ 1 \\ 0 \end{pmatrix}; \quad u_{\downarrow} = \sqrt{E} \begin{pmatrix} 0 \\ 1 \\ 0 \\ -1 \end{pmatrix}; \quad v_{\uparrow} = \sqrt{E} \begin{pmatrix} 1 \\ 0 \\ -1 \\ 0 \end{pmatrix}; \quad v_{\downarrow} = \sqrt{E} \begin{pmatrix} 0 \\ 1 \\ 0 \\ 1 \end{pmatrix}$$

- The electron current can either be obtained from equations (3)-(6) as before or it can be obtained directly from the expressions for the muon current.

$$(j_e)^\mu = \bar{v}(p_2) \gamma^\mu u(p_1) \qquad (j_\mu)^\mu = \bar{u}(p_3) \gamma^\mu v(p_4)$$

- Taking the Hermitian conjugate of the muon current gives

$$\begin{aligned} [\bar{u}(p_3) \gamma^\mu v(p_4)]^\dagger &= [u(p_3)^\dagger \gamma^0 \gamma^\mu v(p_4)]^\dagger \\ &= v(p_4)^\dagger \gamma^{\mu\dagger} \gamma^{0\dagger} u(p_3) & (AB)^\dagger &= B^\dagger A^\dagger \\ &= v(p_4)^\dagger \gamma^{\mu\dagger} \gamma^0 u(p_3) & \gamma^{0\dagger} &= \gamma^0 \\ &= v(p_4)^\dagger \gamma^0 \gamma^\mu u(p_3) & \gamma^{\mu\dagger} \gamma^0 &= \gamma^0 \gamma^\mu \\ &= \bar{v}(p_4) \gamma^\mu u(p_3) \end{aligned}$$

- Taking the complex conjugate of the muon currents for the two non-zero helicity configurations:

$$\bar{\nu}_{\downarrow}(p_4)\gamma^{\mu}u_{\uparrow}(p_3) = [\bar{u}_{\uparrow}(p_3)\gamma^{\nu}\nu_{\downarrow}(p_4)]^* = 2E(0, -\cos\theta, -i, \sin\theta)$$

$$\bar{\nu}_{\uparrow}(p_4)\gamma^{\mu}u_{\downarrow}(p_3) = [\bar{u}_{\downarrow}(p_3)\gamma^{\nu}\nu_{\uparrow}(p_4)]^* = 2E(0, -\cos\theta, i, \sin\theta)$$

To obtain the electron currents we simply need to set $\theta = 0$

$$e^{-} \xrightarrow{\text{red arrow}} \xleftarrow{\text{red arrow}} e^{+}$$


$$e^{-} \xleftarrow{\text{red arrow}} \xrightarrow{\text{red arrow}} e^{+}$$

$e_R^{-} e_L^{+} :$	$\bar{\nu}_{\downarrow}(p_2)\gamma^{\nu}u_{\uparrow}(p_1) = 2E(0, -1, -i, 0)$
$e_L^{-} e_R^{+} :$	$\bar{\nu}_{\uparrow}(p_2)\gamma^{\nu}u_{\downarrow}(p_1) = 2E(0, -1, i, 0)$


Electron Positron Annihilation cont.

★ For $e^+e^- \rightarrow \mu^+\mu^-$ now only have to consider the 4 matrix elements:


M_{RR}




M_{RL}



M_{LR}



M_{LL}




$$\begin{aligned} RR &= RL^\dagger RL \\ RL &= RL^\dagger LR \\ LL &= LR^\dagger RL \\ RR &= LR^\dagger LR \end{aligned}$$

Here the first subscript refers to the helicity of the e^- and the second to the helicity of the μ^- . Don't need to specify other helicities due to "helicity conservation", only certain **chiral** combinations are non-zero.

Matrix Element Calculation

- We can now calculate $M = -\frac{e^2}{s} j_e \cdot j_\mu$ for the four possible helicity combinations.

e.g. the matrix element for $e_R^- e_L^+ \rightarrow \mu_R^- \mu_L^+$ which will denote M_{RR}




★ Using: $e_R^- e_L^+ :$ $(j_e)^\mu = \bar{v}_\downarrow(p_2) \gamma^\mu u_\uparrow(p_1) = 2E(0, -1, -i, 0)$
 $\mu_R^- \mu_L^+ :$ $(j_\mu)^\nu = \bar{u}_\uparrow(p_3) \gamma^\nu v_\downarrow(p_4) = 2E(0, -\cos \theta, i, \sin \theta)$

gives $M_{RR} = -\frac{e^2}{s} [2E(0, -1, -i, 0)] \cdot [2E(0, -\cos \theta, i, \sin \theta)]$
 $= -e^2(1 + \cos \theta)$
 $= -4\pi\alpha(1 + \cos \theta) \quad \text{where} \quad \alpha = e^2/4\pi \approx 1/137$

Similarly

$$|M_{RR}|^2 = |M_{LL}|^2 = (4\pi\alpha)^2(1 + \cos\theta)^2$$

$$|M_{RL}|^2 = |M_{LR}|^2 = (4\pi\alpha)^2(1 - \cos\theta)^2$$



- Assuming that the incoming electrons and positrons are **unpolarized**, all 4 possible initial helicity states are equally likely.

Differential Cross Section

- The cross section is obtained by averaging over the initial spin states and summing over the final spin states:

$$\begin{aligned}\frac{d\sigma}{d\Omega} &= \frac{1}{4} \times \frac{1}{64\pi^2 s} (|M_{RR}|^2 + |M_{RL}|^2 + |M_{LR}|^2 + |M_{LL}|^2) \\ &= \frac{(4\pi\alpha)^2}{256\pi^2 s} (2(1 + \cos\theta)^2 + 2(1 - \cos\theta)^2)\end{aligned}$$





$$\frac{d\sigma}{d\Omega} = \frac{\alpha^2}{4s} (1 + \cos^2\theta)$$

Example:

$$e^+e^- \rightarrow \mu^+\mu^-$$

$$\sqrt{s} = 29 \text{ GeV}$$



..... pure QED, $O(\alpha^3)$
 — QED plus Z contribution

Angular distribution becomes slightly asymmetric in higher order QED or when Z contribution is included

- The total cross section is obtained by integrating over θ, ϕ using


$$\int (1 + \cos^2 \theta) d\Omega = 2\pi \int_{-1}^{+1} (1 + \cos^2 \theta) d\cos \theta = \frac{16\pi}{3}$$

giving the **QED** total cross-section for the process $e^+e^- \rightarrow \mu^+\mu^-$

$$\sigma = \frac{4\pi\alpha^2}{3s}$$

★ Lowest order cross section calculation provides a good description of the data !

This is an impressive result. From first principles we have arrived at an expression for the electron-positron annihilation cross section which is good to **1%**




$\sqrt{s}(\text{GeV})$

Lorentz Invariant form of Matrix Element

- Before concluding this discussion, note that the spin-averaged Matrix Element derived above is written in terms of the muon angle in the C.o.M. frame.

$$\begin{aligned}
 \langle |M_{fi}|^2 \rangle &= \frac{1}{4} \times (|M_{RR}|^2 + |M_{RL}|^2 + |M_{LR}|^2 + |M_{LL}|^2) \\
 &= \frac{1}{4} e^4 (2(1 + \cos \theta)^2 + 2(1 - \cos \theta)^2) \\
 &= e^4 (1 + \cos^2 \theta)
 \end{aligned}$$



- The matrix element is **Lorentz Invariant** (scalar product of 4-vector currents) and it is desirable to write it in a frame-independent form, i.e. express in terms of Lorentz Invariant 4-vector scalar products
- In the C.o.M. $p_1 = (E, 0, 0, E)$ $p_2 = (E, 0, 0, -E)$
 $p_3 = (E, E \sin \theta, 0, E \cos \theta)$ $p_4 = (E, -E \sin \theta, 0, -E \cos \theta)$
giving: $p_1 \cdot p_2 = 2E^2$; $p_1 \cdot p_3 = E^2(1 - \cos \theta)$; $p_1 \cdot p_4 = E^2(1 + \cos \theta)$
- Hence we can write

$$\langle |M_{fi}|^2 \rangle = 2e^4 \frac{(p_1 \cdot p_3)^2 + (p_1 \cdot p_4)^2}{(p_1 \cdot p_2)^2}$$

$$\equiv 2e^4 \left(\frac{t^2 + u^2}{s^2} \right)$$

★ Valid in any frame !

CHIRALITY

- The helicity eigenstates for a particle/anti-particle for $E \gg m$ are:

$$u_{\uparrow} = \sqrt{E} \begin{pmatrix} c \\ se^{i\phi} \\ c \\ se^{i\phi} \end{pmatrix}; \quad u_{\downarrow} = \sqrt{E} \begin{pmatrix} -s \\ ce^{i\phi} \\ s \\ -ce^{i\phi} \end{pmatrix}; \quad v_{\uparrow} = \sqrt{E} \begin{pmatrix} s \\ -ce^{i\phi} \\ -s \\ ce^{i\phi} \end{pmatrix}; \quad v_{\downarrow} = \sqrt{E} \begin{pmatrix} c \\ se^{i\phi} \\ c \\ se^{i\phi} \end{pmatrix}$$

where $s = \sin \frac{\theta}{2}$; $c = \cos \frac{\theta}{2}$

- Define the matrix

$$\gamma^5 \equiv i\gamma^0\gamma^1\gamma^2\gamma^3 = \begin{pmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \end{pmatrix} = \begin{pmatrix} 0 & I \\ I & 0 \end{pmatrix}$$

- In the limit $E \gg m$ the **helicity states** are also eigenstates of γ^5

$$\gamma^5 u_{\uparrow} = +u_{\uparrow}; \quad \gamma^5 u_{\downarrow} = -u_{\downarrow}; \quad \gamma^5 v_{\uparrow} = -v_{\uparrow}; \quad \gamma^5 v_{\downarrow} = +v_{\downarrow}$$

- ★ In general, define the eigenstates of γ^5 as **LEFT** and **RIGHT HANDED CHIRAL** states
 $u_R; \quad u_L; \quad v_R; \quad v_L$

i.e. $\gamma^5 u_R = +u_R; \quad \gamma^5 u_L = -u_L; \quad \gamma^5 v_R = -v_R; \quad \gamma^5 v_L = +v_L$

- In the LIMIT $E \gg m$ (and **ONLY IN THIS LIMIT**):

$$u_R \equiv u_{\uparrow}; \quad u_L \equiv u_{\downarrow}; \quad v_R \equiv v_{\uparrow}; \quad v_L \equiv v_{\downarrow}$$

- ★ This is a subtle but important point: in general the **HELICITY** and **CHIRAL** eigenstates **are not the same**. It is **only** in the **ultra-relativistic limit** that the chiral eigenstates correspond to the helicity eigenstates.
- ★ Chirality is an important concept in the structure of QED, and any interaction of the form $\bar{u}\gamma^\nu u$

- In general, the eigenstates of the chirality operator are:

$$\gamma^5 u_R = +u_R; \quad \gamma^5 u_L = -u_L; \quad \gamma^5 v_R = -v_R; \quad \gamma^5 v_L = +v_L$$

- Define the **projection operators**:

$$P_R = \frac{1}{2}(1 + \gamma^5); \quad P_L = \frac{1}{2}(1 - \gamma^5)$$

- The projection operators, project out the chiral eigenstates

$$\begin{aligned} P_R u_R &= u_R; & P_R u_L &= 0; & P_L u_R &= 0; & P_L u_L &= u_L \\ P_R v_R &= 0; & P_R v_L &= v_L; & P_L v_R &= v_R; & P_L v_L &= 0 \end{aligned}$$

- Note P_R projects out **right-handed particle states** and **left-handed anti-particle states**
- We can then write any spinor in terms of its left and right-handed chiral components:

$$\psi = \psi_R + \psi_L = \frac{1}{2}(1 + \gamma^5)\psi + \frac{1}{2}(1 - \gamma^5)\psi$$

Chirality in QED

- In QED the basic interaction between a fermion and photon is:

$$ie\bar{\psi}\gamma^\mu\phi$$

- Can decompose the spinors in terms of **Left** and **Right**-handed chiral components:

$$\begin{aligned} ie\bar{\psi}\gamma^\mu\phi &= ie(\bar{\psi}_L + \bar{\psi}_R)\gamma^\mu(\phi_R + \phi_L) \\ &= ie(\bar{\psi}_R\gamma^\mu\phi_R + \bar{\psi}_R\gamma^\mu\phi_L + \bar{\psi}_L\gamma^\mu\phi_R + \bar{\psi}_L\gamma^\mu\phi_L) \end{aligned}$$

- Using the properties of γ^5 (Q8 on examples sheet)

$$(\gamma^5)^2 = 1; \quad \gamma^{5\dagger} = \gamma^5; \quad \gamma^5\gamma^\mu = -\gamma^\mu\gamma^5$$

it is straightforward to show

(Q9 on examples sheet)

$$\bar{\psi}_R\gamma^\mu\phi_L = 0; \quad \bar{\psi}_L\gamma^\mu\phi_R = 0$$





- ★ Hence only certain combinations of **chiral** eigenstates contribute to the interaction. This statement is **ALWAYS** true.
- For $E \gg m$, the chiral and helicity eigenstates are equivalent. This implies that for $E \gg m$ only certain helicity combinations contribute to the QED vertex ! This is why previously we found that for two of the four helicity combinations for the muon current were zero

Allowed QED Helicity Combinations





- ♦ In the ultra-relativistic limit the helicity eigenstates \equiv chiral eigenstates
- ♦ In this limit, the only non-zero **helicity** combinations in QED are:

Scattering:

“Helicity conservation”




Annihilation:



Crossing Symmetry

- ♦ In the ultra-relativistic limit the helicity eigenstates \equiv chiral eigenstates
- ♦ In this limit, the only non-zero helicity combinations in QED are:


Annihilation:



$$(j_e)^\mu = \bar{v}(p_2) \gamma^\mu u(p_1)$$

$$(j_\mu)^\mu = \bar{u}(p_3) \gamma^\mu v(p_4)$$

Scattering:



$$j_e^\mu = \bar{u}(p_3) \gamma^\mu u(p_1)$$

$$j_f^\nu = \bar{u}(p_4) \gamma^\nu u(p_2).$$

$$p_1 \rightarrow p_1, p_2 \rightarrow p_3, p_3 \rightarrow p_4 \text{ and } p_4 \rightarrow p_2$$

Summary

- ★ In the centre-of-mass frame the $e^+e^- \rightarrow \mu^+\mu^-$ differential cross-section is

$$\frac{d\sigma}{d\Omega} = \frac{\alpha^2}{4s} (1 + \cos^2 \theta)$$

NOTE: neglected masses of the muons, i.e. assumed $E \gg m_\mu$

- ★ In QED only certain combinations of **LEFT-** and **RIGHT-HANDED CHIRAL** states give non-zero matrix elements
- ★ **CHIRAL** states defined by chiral projection operators

$$P_R = \frac{1}{2}(1 + \gamma^5); \quad P_L = \frac{1}{2}(1 - \gamma^5)$$

- ★ In limit $E \gg m$ the chiral eigenstates correspond to the **HELICITY** eigenstates and only certain **HELICITY** combinations give non-zero matrix elements

