

✔ Congratulations! You passed!

Grade received 100% To pass 80% or higher

Go to next item

1. Let T be a linear transformation in the plane represented by the following matrix:

1 / 1 point

$$\begin{bmatrix} 1 & 0 \\ 2 & 3 \end{bmatrix}$$

The rank of T is:

- ☐ 0

☐ 3

☒ 2

☐ 1

✔ Correct

In this point of the course you have several ways of finding this information. Applying what you've seen in the lecture [Singularity and rank of linear transformations](#) [↗](#), it is necessary to understand how T works in the vectors $(0, 1)$ and $(1, 0)$.

$$\begin{bmatrix} 1 & 0 \\ 2 & 3 \end{bmatrix} \cdot \begin{bmatrix} 0 \\ 1 \end{bmatrix} = \begin{bmatrix} 0 \\ 3 \end{bmatrix}$$

And,

$$\begin{bmatrix} 1 & 0 \\ 2 & 3 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ 0 \end{bmatrix} = \begin{bmatrix} 1 \\ 2 \end{bmatrix}$$

So, $(0, 1) \rightarrow (0, 3)$ and $(1, 0) \rightarrow (1, 2)$.

Note that the vectors $(0, 3)$ and $(1, 2)$ form a parallelogram in the plane, therefore the rank of T is 2.

2. Consider the linear transformation T that maps the vectors $(1,0)$ and $(0, 1)$ in the following manner:

1 / 1 point

$$\begin{aligned} T(0, 1) &= (2, 5) \\ T(1, 0) &= (3, 1) \end{aligned}$$

The area of the parallelogram spanned by transforming the vectors $(0, 1)$ and $(1, 0)$ is:

13

✔ Correct

The matrix associated with this linear transformation is given by

$$\begin{bmatrix} 3 & 2 \\ 1 & 5 \end{bmatrix}$$

The area of the parallelogram spanned by applying T on the vectors $(1, 0)$ and $(0, 1)$ is then given by the determinant of such matrix:

$$\det \left(\begin{bmatrix} 3 & 2 \\ 1 & 5 \end{bmatrix} \right) = 13$$

You may watch again the lectures on [Linear transformations as matrices](#) [↗](#) to review how to obtain the matrix associated with a linear transformation and the lecture on [Determinant as an area](#) [↗](#) to review why the determinant represents the area of such parallelogram.

3. Consider the following three matrices

1 / 1 point

$$M_1 = \begin{bmatrix} 2 & 1 \\ 3 & 1 \end{bmatrix}$$

$$M_2 = \begin{bmatrix} 3 & 5 \\ 1 & 1 \end{bmatrix}$$

$$M_3 = \begin{bmatrix} 2 & 3 \\ 4 & 5 \end{bmatrix}$$

The determinant of $M_1 \cdot M_2 \cdot M_3$ is equal to:

-4

✔ Correct

As you've seen in the lecture [Determinant of a product](#) [↗](#), the determinant os a product of matrices is the product of their determinant. Therefore:

$$\det(M_1 \cdot M_2 \cdot M_3) = \det(M_1) \cdot \det(M_2) \cdot \det(M_3).$$

Since

$$\det(M_1) = 2 - 3 = -1$$

$$\det(M_2) = 3 - 5 = -2$$

$$\det(M_3) = 10 - 12 = -2$$

$$\text{Then } \det(M_1 \cdot M_2 \cdot M_3) = (-1) \cdot (-2) \cdot (-2) = -4$$

4. Let M and N be two square matrices with the same size.

1 / 1 point

Check all statements that are true.

✔ If M and N are non-singular matrices, then so is $M \cdot N$.

✔ Correct

If M and N are non-singular, then $\det(M) \neq 0$ and $\det(N) \neq 0$. Since $\det(M \cdot N) = \det(M)\det(N)$, then $\det(M \cdot N) \neq 0$ as well.

✔ If M is singular, then $M \cdot N$ is singular for any matrix N .

✔ Correct

If M is singular, then $\det(M) = 0$. So, for every matrix N with the same size, $\det(M \cdot N) = \det(M) \cdot \det(N) = 0 \cdot \det(N) = 0$. Therefore, $M \cdot N$ is singular.

☐ $\det(M + N) = \det(M) + \det(N)$.

☐ If $M \cdot N$ is singular, then M **and** N are singular.

5. Let M be the following 3×3 matrix:

1 / 1 point

$$\begin{bmatrix} 0 & 0 & 1 \\ 2 & 2 & 1 \\ 1 & 0 & 0 \end{bmatrix}$$

Compute $\det(M^{-1})$. Please provide your solution in decimal notation not in fraction, using one decimal place.

-.5

✔ Correct

As you've seen in the lecture [Determinant of inverses](#) [↗](#), $\det(M^{-1}) = \det(M)^{-1} = \frac{1}{\det(M)}$, so there is no need to compute the inverse of M ! Computing the determinant for M :

$$\det(M) = (0 \cdot 2 \cdot 0 + 0 \cdot 1 \cdot 1 + 1 \cdot 0 \cdot 2) - (1 \cdot 2 \cdot 1 + 0 \cdot 2 \cdot 0 + 0 \cdot 0 \cdot 1) = -2$$

$$\text{Therefore, } \det(M^{-1}) = \frac{1}{-2} = -0.5.$$