

Expected Number of Coin Tosses to Get Two Consecutive Heads

1. Problem

We are given an unbiased coin and want to estimate the expected number of tosses required to observe two consecutive heads for the first time. The coin is fair, i.e., the probability of heads (H) or tails (T) in any toss is:

$$P(H) = \frac{1}{2}, \quad P(T) = \frac{1}{2}$$

We aim to either derive a formula analytically or simulate the problem using a Monte Carlo algorithm. Here, we first present an analytical method using expected value equations.

2. Solution

Let X be the expected number of tosses required to obtain two consecutive heads. We model the process by considering the different possible outcomes of the initial tosses.

We define three states:

1. S_0 : Initial state, no heads yet.
2. S_1 : Last toss was a head, but not two in a row.
3. S_2 : Terminal state, two heads in a row (stop here).

Let E_0 be the expected number of tosses starting from S_0 , and E_1 be the expected number from S_1 .

We construct the following recurrence relations:

From State S_0 :

From the initial state, we toss a coin:

1. With probability $\frac{1}{2}$, we get a tail and stay in S_0 (wasted one flip): $\Rightarrow E_0 + 1$
2. With probability $\frac{1}{2}$, we get a head and move to S_1 (one flip used): $\Rightarrow E_1 + 1$

$$E_0 = \frac{1}{2}(E_0 + 1) + \frac{1}{2}(E_1 + 1)$$

From State S_1 :

Now the last toss was a head:

1. With probability $\frac{1}{2}$, we get a head again and reach the terminal state (two flips total): $\Rightarrow 2$
2. With probability $\frac{1}{2}$, we get a tail and return to S_0 : $\Rightarrow E_0 + 2$

$$E_1 = \frac{1}{2}(2) + \frac{1}{2}(E_0 + 2)$$

Solving the Equations

Start with the second equation:

$$E_1 = \frac{1}{2}(2) + \frac{1}{2}(E_0 + 2) = 1 + \frac{1}{2}E_0 + 1 = \frac{1}{2}E_0 + 2$$

Now substitute E_1 into the first equation:

$$E_0 = \frac{1}{2}(E_0 + 1) + \frac{1}{2}\left(\frac{1}{2}E_0 + 2 + 1\right) = \frac{1}{2}E_0 + \frac{1}{2} + \frac{1}{4}E_0 + \frac{3}{2}$$

$$E_0 = \left(\frac{3}{4}E_0\right) + 2 \Rightarrow E_0 - \frac{3}{4}E_0 = 2 \Rightarrow \frac{1}{4}E_0 = 2 \Rightarrow E_0 = 8$$

Thus, the expected number of tosses required to get two consecutive heads is:

$$\boxed{8}$$

3. Alternative Monte Carlo Simulation

A Monte Carlo algorithm can also estimate the expected number of tosses empirically. Here's a high-level description:

1. Repeat the following simulation N times (e.g., $N = 10^6$):
 - (a) Initialize a counter for tosses and a memory of the last toss.
 - (b) Toss the coin until two heads appear in a row.
 - (c) Record the number of tosses.
2. Compute the average number of tosses over all trials.

Pseudocode:

```
total_flips = 0
N = 1000000

for i in range(N):
    flips = 0
```

```

last = None
while True:
    flip = random.choice(['H', 'T'])
    flips += 1
    if last == 'H' and flip == 'H':
        break
    last = flip
total_flips += flips

expected_flips = total_flips / N

```

Output (approx.): 8

4. Note

Both the analytical and simulation-based approaches confirm that the expected number of coin tosses required to observe two consecutive heads using an unbiased coin is:

8