

Multiphysics Reduced Order Modelling

Reduced Basis Method for Lead-Cooled Fast Reactor

Parikshit Bajpai

MCSC 6020G - Numerical Analysis
October 28, 2019

- 1 Motivation and Objectives
- 2 Background - Reactor Physics 00.01
- 3 Parameterised Multiphysics Model
- 4 Finite Element Discretisation
- 5 Reduced Basis Formulation
- 6 Expected Outcomes and Results

Motivation

Multiphysics
Reduced
Order
Modelling

Parikshit
Bajpai

Motivation
and Objectives

Background -
Reactor
Physics 00.01

Parameterised
Multiphysics
Model

Finite Element
Discretisation

Reduced Basis
Formulation

Expected
Outcomes and
Results

The full order simulation of nuclear reactors is extremely costly!

Can we reduce the computational time and cost associated with these simulations without compromising with the accuracy?

Objective

Multiphysics
Reduced
Order
Modelling

Parikshit
Bajpai

Motivation
and Objectives

Background -
Reactor
Physics 00.01

Parameterised
Multiphysics
Model

Finite Element
Discretisation

Reduced Basis
Formulation

Expected
Outcomes and
Results

Develop a Reduced Basis method, with basis functions sampled by a Proper Orthogonal Decomposition technique, to develop a reduced order model of a parametrised Lead-cooled Fast Reactor single-channel.

Lead-cooled Fast Reactor (LFR)

Multiphysics
Reduced
Order
Modelling

Parikshit
Bajpai

Motivation
and Objectives


Background -
Reactor
Physics 00.01

Parameterised
Multiphysics
Model

Finite Element
Discretisation

Reduced Basis
Formulation

Expected
Outcomes and
Results



Neutron Flux

Multiphysics
Reduced
Order
Modelling

Parikshit
Bajpai

Motivation
and Objectives

Background -
Reactor
Physics 00.01

Parameterised
Multiphysics
Model

Finite Element
Discretisation

Reduced Basis
Formulation

Expected
Outcomes and
Results

Neutron balance

$$\frac{\partial \phi}{\partial t} = -\nabla \cdot D \nabla \phi + \nu \Sigma_f \phi - (\Sigma_a + \Sigma_s) \phi$$

ϕ - Neutron Flux

D - Diffusion Coefficient

Σ_a - Macroscopic Absorption Cross Section

Σ_s - Macroscopic Scattering Cross Section

ν - Fissile Yield (No. of neutrons produced per fission)

Σ_f - Macroscopic Fission Cross Section

Neutron Flux

Multiphysics
Reduced
Order
Modelling

Parikshit
Bajpai

Motivation
and Objectives

Background -
Reactor
Physics 00.01


Parameterised
Multiphysics
Model

Finite Element
Discretisation

Reduced Basis
Formulation

Expected
Outcomes and
Results

Rate of change
of neutron flux


$$\frac{\partial \phi}{\partial t} = -\nabla \cdot D \nabla \phi + \nu \Sigma_f \phi - (\Sigma_a + \Sigma_s) \phi$$

Neutron Flux

Multiphysics
Reduced
Order
Modelling

Parikshit
Bajpai

Motivation
and Objectives

Background -
Reactor
Physics 00.01

Parameterised
Multiphysics
Model

Finite Element
Discretisation

Reduced Basis
Formulation

Expected
Outcomes and
Results

Rate of change
of neutron flux

$$\frac{\partial \phi}{\partial t} = -\nabla \cdot D \nabla \phi + \nu \Sigma_f \phi - (\Sigma_a + \Sigma_s) \phi$$

Diffusion of neutrons

Neutron Flux

Multiphysics
Reduced
Order
Modelling

Parikshit
Bajpai

Motivation
and Objectives

Background -
Reactor
Physics 00.01

Parameterised
Multiphysics
Model

Finite Element
Discretisation

Reduced Basis
Formulation

Expected
Outcomes and
Results

Rate of change
of neutron flux

Neutron production
through fission

$$\frac{\partial \phi}{\partial t} = -\nabla \cdot D \nabla \phi + \nu \Sigma_f \phi - (\Sigma_a + \Sigma_s) \phi$$

Diffusion of neutrons

Neutron Flux

Multiphysics
Reduced
Order
Modelling

Parikshit
Bajpai

Motivation
and Objectives

Background -
Reactor
Physics 00.01

Parameterised
Multiphysics
Model

Finite Element
Discretisation

Reduced Basis
Formulation

Expected
Outcomes and
Results

Rate of change
of neutron flux

Neutron production
through fission

$$\frac{\partial \phi}{\partial t} = -\nabla \cdot D \nabla \phi + \nu \Sigma_f \phi - (\Sigma_a + \Sigma_s) \phi$$

Diffusion of neutrons

Loss of neutrons
due to absorption
and scattering

Heat Transfer

Multiphysics
Reduced
Order
Modelling

Parikshit
Bajpai

Motivation
and Objectives

Background -
Reactor
Physics 00.01

Parameterised
Multiphysics
Model

Finite Element
Discretisation

Reduced Basis
Formulation

Expected
Outcomes and
Results

Energy Balance

$$\frac{\partial T}{\partial t} = \nabla \cdot [(K + K_T) \nabla T] - \rho C_p \mathbf{v} \cdot \nabla T + Q$$

T - Temperature

K - Thermal Conductivity

K_T - Turbulent Thermal Conductivity

C_p - Heat Capacity

\mathbf{v} - Velocity of fluid

Q - Source Term

Heat Transfer

Multiphysics
Reduced
Order
Modelling

Parikshit
Bajpai

Motivation
and Objectives

Background -
Reactor
Physics 00.01

Parameterised
Multiphysics
Model

Finite Element
Discretisation

Reduced Basis
Formulation

Expected
Outcomes and
Results

Rate of change
of temperature


$$\frac{\partial T}{\partial t} = \nabla \cdot [(K + K_T) \nabla T] - \rho C_p \mathbf{v} \cdot \nabla T + Q$$

Heat Transfer

Multiphysics
Reduced
Order
Modelling

Parikshit
Bajpai

Motivation
and Objectives

Background -
Reactor
Physics 00.01

Parameterised
Multiphysics
Model

Finite Element
Discretisation

Reduced Basis
Formulation

Expected
Outcomes and
Results

Rate of change
of temperature

$$\frac{\partial T}{\partial t} = \nabla \cdot [(K + K_T) \nabla T] - \rho C_p \mathbf{v} \cdot \nabla T + Q$$

Diffusion term

Heat Transfer

Multiphysics
Reduced
Order
Modelling

Parikshit
Bajpai

Motivation
and Objectives

Background -
Reactor
Physics 00.01

Parameterised
Multiphysics
Model

Finite Element
Discretisation

Reduced Basis
Formulation

Expected
Outcomes and
Results

Rate of change
of temperature

Advection term

$$\frac{\partial T}{\partial t} = \nabla \cdot [(K + K_T) \nabla T] - \rho C_p \mathbf{v} \cdot \nabla T + Q$$

Diffusion term

Heat Transfer

Multiphysics
Reduced
Order
Modelling

Parikshit
Bajpai

Motivation
and Objectives

Background -
Reactor
Physics 00.01

Parameterised
Multiphysics
Model

Finite Element
Discretisation

Reduced Basis
Formulation

Expected
Outcomes and
Results

Rate of change
of temperature

Advection term

$$\frac{\partial T}{\partial t} = \nabla \cdot [(K + K_T) \nabla T] - \rho C_p \mathbf{v} \cdot \nabla T + Q$$

Diffusion term

Source term

Multiphysics Model

Multiphysics
Reduced
Order
Modelling

Parikshit
Bajpai

Motivation
and Objectives


Background -
Reactor
Physics 00.01

Parameterised
Multiphysics
Model

Finite Element
Discretisation

Reduced Basis
Formulation

Expected
Outcomes and
Results



Parametrised Multiphysics Model

Multiphysics
Reduced
Order
Modelling

Parikshit
Bajpai

Motivation
and Objectives

Background -
Reactor
Physics 00.01


Parameterised
Multiphysics
Model

Finite Element
Discretisation

Reduced Basis
Formulation

Expected
Outcomes and
Results

- Parametrised LFR single channel model.
- Rotational symmetry along fuel channel (r-z model).
- Time-independent settings.
- Parameters:
 - Inner radius of fuel pellet
 $\mu_1 \in [0.1, 0.43]$ (cm)
 - Nominal coolant flow rate
 $\mu_2 \in [0.8, 1.6]$ (m/s)



Neutronics

Multiphysics
Reduced
Order
Modelling

Parikshit
Bajpai

Motivation
and Objectives

Background -
Reactor
Physics 00.01

Parameterised
Multiphysics
Model

Finite Element
Discretisation

Reduced Basis
Formulation

Expected
Outcomes and
Results

The multi-group neutron diffusion equation in stationary formulation is a generalised eigenvalue problem.

$$\left(-\nabla \cdot \overline{\overline{D}} \nabla + \overline{\overline{\Sigma}}_a + \overline{\overline{\Sigma}}_s\right) \overline{\Phi} = \lambda_{eff} \overline{\chi} \overline{F}' \overline{\Phi}$$

$$\begin{aligned} \overline{\Phi} &= \begin{bmatrix} \Phi_1(\mathbf{r}) \\ \vdots \\ \Phi_6(\mathbf{r}) \end{bmatrix} \quad \overline{\overline{D}} = \begin{bmatrix} D_1(\mathbf{r}) & \dots & 0 \\ & \ddots & \\ & & D_6(\mathbf{r}) \end{bmatrix} \quad \overline{\overline{\Sigma}}_a = \begin{bmatrix} \Sigma_a^1(\mathbf{r}) & \dots & 0 \\ & \ddots & \\ & & \Sigma_a^6(\mathbf{r}) \end{bmatrix} \\ \overline{\overline{\Sigma}}_s &= \begin{bmatrix} \sum_{g' \neq 1} \Sigma_s^{1 \rightarrow g'}(\mathbf{r}) & \dots & -\Sigma_s^{6 \rightarrow 1} \\ & \ddots & \\ -\Sigma_s^{1 \rightarrow 6} & \dots & \sum_{g' \neq 6} \Sigma_s^{6 \rightarrow g'} \end{bmatrix} \quad \overline{\chi} = \begin{bmatrix} \chi_1 \\ \vdots \\ \chi_6 \end{bmatrix} \quad \overline{F} = \begin{bmatrix} \nu \Sigma_f^1(\mathbf{r}) \\ \vdots \\ \nu \Sigma_f^6(\mathbf{r}) \end{bmatrix} \end{aligned}$$

Neutronics

Multiphysics
Reduced
Order
Modelling

Parikshit
Bajpai

Motivation
and Objectives

Background -
Reactor
Physics 00.01

Parameterised
Multiphysics
Model

Finite Element
Discretisation

Reduced Basis
Formulation

Expected
Outcomes and
Results

The multi-group neutron diffusion equation in stationary formulation is a generalised eigenvalue problem.

$$\left(-\nabla \cdot \overline{\overline{D}} \nabla + \overline{\overline{\Sigma}}_a + \overline{\overline{\Sigma}}_s\right) \overline{\Phi} = \lambda_{eff} \overline{\chi} \overline{F}' \overline{\Phi}$$

Fuel domain

$$\Sigma(T, \rho) = \frac{\rho}{\rho_0} \left[\Sigma_0 + \alpha \log \left(\frac{T}{T_0} \right) \right]$$

Lead domain

$$\Sigma(T, \rho) = \frac{\rho}{\rho_0} \Sigma_0$$

Neutronics

Multiphysics
Reduced
Order
Modelling

Parikshit
Bajpai

Motivation
and Objectives

Background -
Reactor
Physics 00.01

Parameterised
Multiphysics
Model

Finite Element
Discretisation

Reduced Basis
Formulation

Expected
Outcomes and
Results

The multi-group neutron diffusion equation in stationary formulation is a generalised eigenvalue problem.

$$\left(-\nabla \cdot \overline{\overline{D}} \nabla + \overline{\overline{\Sigma}_a} + \overline{\overline{\Sigma}_s} \right) \overline{\Phi} = \lambda_{eff} \overline{\chi} \overline{F}' \overline{\Phi}$$

Boundary conditions

- Albedo boundary conditions provide a good compromise between accuracy and computational requirements.

$$\overline{n} \cdot (D_g \nabla \Phi_g) = -\gamma_z \phi_g$$

$$\overline{n} \cdot (D_g \nabla \Phi_g) = -\gamma_r \phi_g$$

Heat Transfer

Multiphysics
Reduced
Order
Modelling

Parikshit
Bajpai

Motivation
and Objectives

Background -
Reactor
Physics 00.01

Parameterised
Multiphysics
Model

Finite Element
Discretisation

Reduced Basis
Formulation

Expected
Outcomes and
Results

Within the cladding and gap domains the heat transfer is purely conductive.

$$-\nabla \cdot (K \nabla T) = 0$$

Within the fuel domain the heat transfer is purely conductive but has a source term.

$$-\nabla \cdot (\nabla T) = \sum_{g=1}^6 \Sigma_g^f E_f \Phi_g$$

Heat Transfer

Multiphysics
Reduced
Order
Modelling

Parikshit
Bajpai

Motivation
and Objectives

Background -
Reactor
Physics 00.01

Parameterised
Multiphysics
Model

Finite Element
Discretisation

Reduced Basis
Formulation

Expected
Outcomes and
Results

Within the lead domain, the heat transfer is has an advective and a conductive term.

$$-\nabla \cdot [(K + K_T)\nabla T] = \rho C_p \mathbf{v} \cdot \nabla T$$

Boundary conditions

- Symmetry conditions at inner radius of the fluid domain and and outerradius of fluid domain.
- Homogeneous Neumann conditions on top and bottom of the domain.

Back to Objective

Multiphysics
Reduced
Order
Modelling

Parikshit
Bajpai

Motivation
and Objectives

Background -
Reactor
Physics 00.01

Parameterised
Multiphysics
Model

Finite Element
Discretisation

Reduced Basis
Formulation

Expected
Outcomes and
Results

The full-fidelity model can have $\mathcal{O}(10^6 - 10^9)$ degrees of freedom.

We want to capture the behaviour of the system without using a high-fidelity, computationally expensive discretisation scheme.

Offline/Online Strategy

Multiphysics
Reduced
Order
Modelling

Parikshit
Bajpai

Motivation
and Objectives


Background -
Reactor
Physics 00.01

Parameterised
Multiphysics
Model

Finite Element
Discretisation

Reduced Basis
Formulation

Expected
Outcomes and
Results



Finite Element Discretisation

Multiphysics
Reduced
Order
Modelling

Parikshit
Bajpai

Motivation
and Objectives

Background -
Reactor
Physics 00.01

Parameterised
Multiphysics
Model

Finite Element
Discretisation

Reduced Basis
Formulation

Expected
Outcomes and
Results

For a given $\mu \in \mathcal{D}$, find $\bar{\Phi}(\mu) \in V^{\mathcal{N}}$ and $T(\mu) \in W^{\mathcal{N}}$ such that

$$a(\bar{\Phi}, \bar{\nu}; \mu, \Sigma(T)) = \lambda_{\text{eff}} m(\bar{\Phi}, \bar{\nu}; \mu, \Sigma(T)) \quad \forall \bar{\nu} \in V^{\mathcal{N}}$$

$$d(T, w; \mu) = f(w; \mu, S(\bar{\Phi})) \quad \forall w \in W^{\mathcal{N}}$$

$$\begin{aligned} a(\bar{\Phi}, \bar{\nu}; \mu, \Sigma(T)) &= \int_{\Omega} \bar{D}(\nabla \bar{\Phi} \cdot \nabla \bar{\nu}) + (\bar{\Sigma}_a + \bar{\Sigma}_s) \bar{\Phi} \bar{\nu} \\ &\quad + \int_{\Gamma_{\text{side}}} \gamma_r \bar{\Phi} \bar{\nu} + \int_{\Gamma_{\text{top}} + \Gamma_{\text{bot}}} \gamma_z \bar{\Phi} \bar{\nu} \end{aligned}$$

$$m(\bar{\Phi}, \bar{\nu}; \mu, \Sigma(T)) = \int_{\Omega} \bar{\chi} \bar{F}' \bar{\Phi} \bar{\nu}$$

Finite Element Discretisation

Multiphysics
Reduced
Order
Modelling

Parikshit
Bajpai

Motivation
and Objectives

Background -
Reactor
Physics 00.01

Parameterised
Multiphysics
Model

Finite Element
Discretisation

Reduced Basis
Formulation

Expected
Outcomes and
Results

For a given $\mu \in \mathcal{D}$, find $\bar{\Phi}(\mu) \in V^{\mathcal{N}}$ and $T(\mu) \in W^{\mathcal{N}}$ such that

$$a(\bar{\Phi}, \bar{\nu}; \mu, \Sigma(T)) = \lambda_{\text{eff}} m(\bar{\Phi}, \bar{\nu}; \mu, \Sigma(T)) \quad \forall \bar{\nu} \in V^{\mathcal{N}}$$

$$d(T, w; \mu) = f(w; \mu, S(\bar{\Phi})) \quad \forall w \in W^{\mathcal{N}}$$

$$d(T, w; \mu) = \int_{\Omega} K \nabla T \nabla w + \int_{\Omega_{\text{lead}}} K_T \nabla T \nabla w$$

$$f(w; \mu, S(\bar{\Phi})) = \int_{\Omega_{\text{fuel}}} Q w + \int_{\Omega_{\text{lead}}} \rho C_p (\mathbf{v} \cdot \nabla T) w$$

Affine Parametric Dependence

Multiphysics
Reduced
Order
Modelling

Parikshit
Bajpai

$$\mathbf{a}(\bar{\Phi}, \bar{\nu}; \mu, \Sigma(T)) = \sum_{q=1}^{Q_a} \Theta_a^q(\mu) \mathbf{a}^q(\bar{\Phi}, \bar{\nu}; \Sigma(T))$$

Motivation
and Objectives

Background -
Reactor
Physics 00.01

Parameterised
Multiphysics
Model

Finite Element
Discretisation

Reduced Basis
Formulation

Expected
Outcomes and
Results

$$\mathbf{m}(\bar{\Phi}, \bar{\nu}; \mu, \Sigma(T)) = \sum_{q=1}^{Q_m} \Theta_m^q(\mu) \mathbf{m}^q(\bar{\Phi}, \bar{\nu}; \Sigma(T))$$

$$\mathbf{d}(T, w; \mu) = \sum_{q=1}^{Q_d} \Theta_d^q(\mu) \mathbf{d}^q(T, w)$$

$$\mathbf{f}(w; \mu, S(\bar{\phi})) = \sum_{q=1}^{Q_f} \Theta_f^q(\mu) \mathbf{f}^q(w; S(\bar{\phi}))$$

Reduced Basis Model

Multiphysics
Reduced
Order
Modelling

Parikshit
Bajpai

Motivation
and Objectives

Background -
Reactor
Physics 00.01

Parameterised
Multiphysics
Model

Finite Element
Discretisation

Reduced Basis
Formulation

Expected
Outcomes and
Results

The RB method is aimed at constructing reduced order solutions $\bar{\Phi}^{N_f}$ and T^{N_T} such that

$$a\left(\bar{\Phi}^{N_f}, \bar{\nu}; \mu, \Sigma(T^{N_T})\right) = \lambda_{eff} m\left(\bar{\Phi}^{N_f}, \bar{\nu}; \mu, \Sigma(T^{N_T})\right) \\ \forall \bar{\nu} \in V^{N_f} \subset V^{\mathcal{N}}$$

$$d\left(T^{N_T}, w; \mu\right) = f\left(w; \mu, S(\bar{\Phi}^{N_f})\right) \\ \forall w \in W^{N_T} \subset W^{\mathcal{N}}$$

Reduced Basis Model

The RB method is aimed at constructing reduced order solutions $\bar{\Phi}^{N_f}$ and T^{N_T} such that

$$\mathbf{a} \left(\bar{\Phi}^{N_f}, \bar{\nu}; \mu, \Sigma(T^{N_T}) \right) = \lambda_{eff} \mathbf{m} \left(\bar{\Phi}^{N_f}, \bar{\nu}; \mu, \Sigma(T^{N_T}) \right) \\ \forall \bar{\nu} \in V^{N_f} \subset V^{\mathcal{N}}$$

$$\mathbf{d} \left(T^{N_T}, w; \mu \right) = \mathbf{f} \left(w; \mu, S(\bar{\Phi}^{N_f}) \right) \\ \forall w \in W^{N_T} \subset W^{\mathcal{N}}$$

The reduced spaces V^{N_f} and W^{N_T} have dimensions N_f and N_T respectively and are defined as:

$$V^{N_f} = \text{span} \left\{ \xi_1^f, \dots, \xi_{N_f}^f \right\} \quad W^{N_T} = \text{span} \left\{ \xi_1^T, \dots, \xi_{N_T}^T \right\}$$

Reduced Basis Model

Multiphysics
Reduced
Order
Modelling

Parikshit
Bajpai

Motivation
and Objectives

Background -
Reactor
Physics 00.01

Parameterised
Multiphysics
Model

Finite Element
Discretisation

Reduced Basis
Formulation

Expected
Outcomes and
Results

$$\Phi^{N_f} = \sum_{i=1}^{N_f} \phi_{N,i} \xi_i^f \quad T^{N_T} = \sum_{j=1}^{N_T} T_{N,j} \xi_j^T$$

$$a(\mathcal{Z}_f \overline{\phi_N}, \overline{\nu}; \mu, \Sigma(\mathcal{Z}_T \overline{T_N})) = \lambda_{\text{eff}}^N m(\mathcal{Z}_f \overline{\phi_N}, \overline{\nu}; \mu, \Sigma(\mathcal{Z}_T \overline{T_N}))$$

$$d(\mathcal{Z}_f \overline{T_N}, w; \mu) = f(w; \mu, S(\mathcal{Z}_f \overline{\phi_N}))$$

$$\mathcal{Z}_f = [\xi_1^f | \dots | \xi_{N_f}^f] \quad \mathcal{Z}_T = [\xi_1^T | \dots | \xi_{N_T}^T]$$

Offline Phase

Multiphysics
Reduced
Order
Modelling

Parikshit
Bajpai

Motivation
and Objectives

Background -
Reactor
Physics 00.01

Parameterised
Multiphysics
Model

Finite Element
Discretisation

Reduced Basis
Formulation

Expected
Outcomes and
Results

Offline

- Run high fidelity computations to get snapshots.
- Construct matrices \mathcal{Z}_f and \mathcal{Z}_T required to build the matrices of ROM
- Matrices corresponding to the bilinear form of affine decompositions.

Online

- Multiply reduced matrices by proper coefficients and sum together.
- Solve the ROM.

Expected Outcomes and Results

Multiphysics
Reduced
Order
Modelling

Parikshit
Bajpai

Motivation
and Objectives

Background -
Reactor
Physics 00.01

Parameterised
Multiphysics
Model

Finite Element
Discretisation

Reduced Basis
Formulation

Expected
Outcomes and
Results

- Evaluate the errors between the reactivity provided by ROM and FE solution and error vs. number of basis functions.

$$\epsilon_{\lambda}(\mu) = |\lambda_N(\mu) - \lambda_{\mathcal{N}}(\mu)|$$

- Evaluate the error between the neutron flux provided by the two models.

$$\epsilon_f = \int_{\Omega} (\Phi^{N_f} - \bar{\Phi})^2$$

- Evaluate the error between the temperature field provided by the two models.

$$\epsilon_T = \int_{\Omega} (T^{N_f} - T)^2$$

Expected Outcomes and Results

Multiphysics
Reduced
Order
Modelling

Parikshit
Bajpai

Motivation
and Objectives

Background -
Reactor
Physics 00.01

Parameterised
Multiphysics
Model

Finite Element
Discretisation

Reduced Basis
Formulation

Expected
Outcomes and
Results

- Comparison of computational cost of ROM and FE model.

$$\text{Speed-up} = \frac{\text{Wall time of ROM calculation}}{\text{Wall time of FE simulation}}$$

- Computational break-even.

$$\text{Break-even} = \frac{\text{Total Offline time}}{\text{Time for 1 FE simulation}}$$

Sample Results

Multiphysics
Reduced
Order
Modelling

Parikshit
Bajpai

Motivation
and Objectives


Background -
Reactor
Physics 00.01

Parameterised
Multiphysics
Model


Finite Element
Discretisation

Reduced Basis
Formulation

Expected
Outcomes and
Results



(a) "Truth"



(b) ROM

Sample Results

Multiphysics
Reduced
Order
Modelling

Parikshit
Bajpai

Motivation
and Objectives

Background -
Reactor
Physics 00.01

Parameterised
Multiphysics
Model

Finite Element
Discretisation

Reduced Basis
Formulation

Expected
Outcomes and
Results

