

# Discrete random variables — Expectation and variance

MTH382 Probability Theory for Finance and Actuarial Science

---

Parin Chaipunya

KMUTT

↳ Mathematics @ Faculty of Science

This lecture is bears the most important tools in probability, which are expectation and variance of a r.v. Expectation could be thought of as the *mean* (or *average*) of all possible values produced by a random variable.

# Expectation

---

## Definition

Always consider a probability space  $(\Omega, \mathcal{F}, P)$ .

### Definition 1.

Let  $X$  be a discrete r.v. with values in  $E \subset \mathbb{R}$  with the condition that

$$\text{either } E \subset \mathbb{R}_+ \quad \text{or} \quad \sum_{x \in E} |x| P(X = x) < \infty. \quad (1)$$

Then the **expectation** (or **expected value**) of  $X$ , denoted with  $\mathbb{E}X$  (or  $\mathbb{E}[X]$ ), is defined by

$$\mathbb{E}X := \sum_{x \in E} x P(X = x).$$



### Remark.

The condition (1) is only to guarantee that the summation in the definition of  $\mathbb{E}X$  is well-defined. Note that an expectation could take an infinite value.

# Examples

We begin with the simplest examples.

## *Example 2.*

Compute  $\mathbb{E}X$  where  $X$  represents the following situations.

- (1)  $X$  is the result of tossing a fair dice.
- (2)  $X$  is the result of tossing an unfair dice with

$$P(\square) = 0.2, \quad P(\blacksquare) = 0.1, \quad P(\blacklozenge) = 0.2, \quad P(\blacksquare) = 0.1, \quad P(\blacksquare) = 0.3, \quad P(\blacksquare) = 0.1.$$

## Example

---

Now we move to a more complicate example.

***Example 3.***

Consider  $S_n$  which is the number of heads appearing in tossing a coin  $n$  times.

Calculate the expected value  $\mathbb{E}[S_n]$ .

## Example

The next example shows that even a r.v.  $X$  has finite values, its expectation could be infinite.

**Example 4.**

Let  $X$  be a r.v. taking values in  $E = \mathbb{N}$  with the distribution

$$P(X = n) = \frac{1}{cn^2}$$

for each  $n \in \mathbb{N}$ , where  $c = \sum_{n=1}^{\infty} \frac{1}{n^2} < \infty$ .

- (a) Show that  $P$  is a probability measure.
- (b) Show that  $\mathbb{E}X = +\infty$

# Expectation

---

Properties of  $\mathbb{E}$



## Properties of $\mathbb{E}$

### Theorem 5.

*The following properties hold.*

(a) *The expectation is linear, i.e.*

$$\mathbb{E}[\lambda_1 X_1 + \cdots + \lambda_n X_n] = \lambda_1 \mathbb{E}X_1 + \cdots + \lambda_n \mathbb{E}X_n$$

*for any  $\lambda_1, \dots, \lambda_n \in \mathbb{R}$ , where  $X_1, \dots, X_n : \Omega \rightarrow \mathbb{R}$  are discrete r.v.s.*

(b) *The expectation is monotone, i.e. for any discrete r.v.s  $X_1, X_2 : \Omega \rightarrow \mathbb{R}$  such that  $X_1 \leq X_2$ , then*

$$\mathbb{E}X_1 \leq \mathbb{E}X_2.$$

(c) *The expectation satisfies the triangle inequality, i.e.*

$$|\mathbb{E}X| \leq \mathbb{E}|X|$$

# Variance

---

## Definition

Take a discrete r.v.  $X$  taking values in  $E \subset \mathbb{R}$ . We say that it is **integrable** if  $\mathbb{E}[|X|] < \infty$  and **square integrable** if  $\mathbb{E}[X^2] < \infty$ .

### Definition 6.

Let  $X$  be a discrete r.v. with values in  $E \subset \mathbb{R}$  which is square integrable, and let  $\mu := \mathbb{E}X$ . The **variance** of  $X$  is defined by

$$\text{var}(X) := \mathbb{E}[(X - \mu)^2] = \sum_{x \in E} (x - \mu)^2 P(X = x).$$

A lot of times, we use the notation  $\sigma^2 := \text{var}(X)$ .

**Proposition 7.**

*The variance also has the following expression*

$$\sigma^2 = \mathbb{E}[X^2] - \mu^2.$$

## Examples

### *Example 8.*

Consider two r.v.s  $X$  and  $Y$ , where we express them with their distribution functions

$$\pi_X(x) := \begin{cases} 0.5 & \text{if } x = 10, \\ 0.5 & \text{if } x = -10, \\ 0 & \text{otherwise,} \end{cases} \quad \pi_Y(y) := \begin{cases} 1 & \text{if } y = 0, \\ 0 & \text{otherwise.} \end{cases}$$

Compare and explain between their expectations  $\mathbb{E}X$  and  $\mathbb{E}Y$  and also their variances  $\sigma_X^2$  and  $\sigma_Y^2$ .

### *Example 9.*

Compute the variances of r.v.s in the previous examples.

# Variance

---

Propoerties of var

### Theorem 10.

Let  $X$  be a discrete r.v. and  $a, b \in \mathbb{R}$ , then

$$\text{var}(aX + b) = a^2 \text{var}(X).$$

### Theorem 11.

If  $X_1, \dots, X_n$  are independent discrete r.v.s, then

$$\text{var}(X_1 + \dots + X_n) = \text{var}(X_1) + \dots + \text{var}(X_n).$$

***Example 12.***

Consider repeatedly tossing a weighted coin with  $P(H) = p$  and  $P(T) = 1 - p$ , with  $p \in (0, 1)$ . Find the expected value and variance of  $S_n$  representing the number of heads appearing in the  $n$  tosses.



# Moments

---

# Moments

We have studied  $\mathbb{E}X$  and  $\text{var}(X)$ , which is related to  $\mathbb{E}[X^2]$ . One might be curious about  $\mathbb{E}[X^k]$  for larger  $k \in \mathbb{N}$  and in fact, some of them has special meanings. For examples,  $\mathbb{E}[X^3]$  is the **skewness** (the lack of symmetry) and  $\mathbb{E}[X^4]$  is the **kurtosis** (the fatness of the tail).

In fact,  $\mathbb{E}[X^k]$  is called the  $k^{\text{th}}$  **moment** of  $X$ .

One could also consider the **moment generating function**  $M_X(s) := e^{sX}$ , since this function captures *all* the moments of  $X$  from the fact that

$$\left. \frac{d^k M_X}{ds^k} \right|_{s=0} = \mathbb{E}[X^k]$$

for all  $k \in \mathbb{N}$ .

## Takeaways

---

# Takeaways

---

- $\mathbb{E}$  is the **mean** and  $\text{var}$  is the **variance** of the data encoded by a given random variable.
- $\mathbb{E}$  is linear but  $\text{var}$  is not.
- $\text{var}$  is additive only for independent random variables.

